


Solutions for College Algebra 11th Edition by Sullivan

[CLICK HERE TO ACCESS COMPLETE Solutions](#)



Solutions

Chapter R

Review

Section R.1

1. rational
2. $4 + 5 \cdot 6 - 3 = 4 + 30 - 3 = 31$
3. Distributive
4. c
5. a
6. b
7. True
8. False; The Zero-Product Property states that if a product equals 0, then at least one of the factors must equal 0.
9. False; 6 is the Greatest Common Factor of 12 and 18. The Least Common Multiple is the smallest value that both numbers will divide evenly. The LCM for 12 and 18 is 36.
10. True
11. $A \cup B = \{1, 3, 4, 5, 9\} \cup \{2, 4, 6, 7, 8\}$
 $= \{1, 2, 3, 4, 5, 6, 7, 8, 9\}$
12. $A \cup C = \{1, 3, 4, 5, 9\} \cup \{1, 3, 4, 6\}$
 $= \{1, 3, 4, 5, 6, 9\}$
13. $A \cap B = \{1, 3, 4, 5, 9\} \cap \{2, 4, 6, 7, 8\} = \{4\}$
14. $A \cap C = \{1, 3, 4, 5, 9\} \cap \{1, 3, 4, 6\} = \{1, 3, 4\}$
15. $(A \cup B) \cap C$
 $= (\{1, 3, 4, 5, 9\} \cup \{2, 4, 6, 7, 8\}) \cap \{1, 3, 4, 6\}$
 $= \{1, 2, 3, 4, 5, 6, 7, 8, 9\} \cap \{1, 3, 4, 6\}$
 $= \{1, 3, 4, 6\}$
16. $(A \cap B) \cup C$
 $= (\{1, 3, 4, 5, 9\} \cap \{2, 4, 6, 7, 8\}) \cup \{1, 3, 4, 6\}$
 $= \{4\} \cup \{1, 3, 4, 6\}$
 $= \{1, 3, 4, 6\}$
17. $\overline{A} = \{0, 2, 6, 7, 8\}$
18. $\overline{C} = \{0, 2, 5, 7, 8, 9\}$
19. $\overline{A \cap B} = \overline{\{1, 3, 4, 5, 9\} \cap \{2, 4, 6, 7, 8\}}$
 $= \overline{\{4\}} = \{0, 1, 2, 3, 5, 6, 7, 8, 9\}$
20. $\overline{B \cup C} = \overline{\{2, 4, 6, 7, 8\} \cup \{1, 3, 4, 6\}}$
 $= \overline{\{1, 2, 3, 4, 6, 7, 8\}} = \{0, 5, 9\}$
21. $\overline{A} \cup \overline{B} = \{0, 2, 6, 7, 8\} \cup \{0, 1, 3, 5, 9\}$
 $= \{0, 1, 2, 3, 5, 6, 7, 8, 9\}$
22. $\overline{B} \cap \overline{C} = \{0, 1, 3, 5, 9\} \cap \{0, 2, 5, 7, 8, 9\}$
 $= \{0, 5, 9\}$
23. a. $\{2, 5\}$
b. $\{-6, 2, 5\}$
c. $\left\{-6, \frac{1}{2}, -1.333\ldots = -1.\overline{3}, 2, 5\right\}$
d. $\{\pi\}$
e. $\left\{-6, \frac{1}{2}, -1.333\ldots = -1.\overline{3}, \pi, 2, 5\right\}$
24. a. $\{1\}$
b. $\{0, 1\}$
c. $\left\{-\frac{5}{3}, 2.060606\ldots = 2.\overline{06}, 1.25, 0, 1\right\}$
d. $\{\sqrt{5}\}$

Chapter R: Review

- e. $\left\{-\frac{5}{3}, 2.060606\dots = 2.\overline{06}, 1.25, 0, 1, \sqrt{5}\right\}$
25. a. $\{1\}$
 b. $\{0, 1\}$
 c. $\left\{0, 1, \frac{1}{2}, \frac{1}{3}, \frac{1}{4}\right\}$
 d. None
 e. $\left\{0, 1, \frac{1}{2}, \frac{1}{3}, \frac{1}{4}\right\}$
26. a. None
 b. $\{-1\}$
 c. $\{-1.3, -1.2, -1.1, -1\}$
 d. None
 e. $\{-1.3, -1.2, -1.1, -1\}$
27. a. None
 b. None
 c. None
 d. $\left\{\sqrt{2}, \pi, \sqrt{2}+1, \pi+\frac{1}{2}\right\}$
 e. $\left\{\sqrt{2}, \pi, \sqrt{2}+1, \pi+\frac{1}{2}\right\}$
28. a. None
 b. None
 c. $\left\{\frac{1}{2}+10.3\right\}$
 d. $\left\{-\sqrt{2}, \pi+\sqrt{2}\right\}$
 e. $\left\{-\sqrt{2}, \pi+\sqrt{2}, \frac{1}{2}+10.3\right\}$
29. a. 18.953 b. 18.952
30. a. 25.861 b. 25.861
31. a. 28.653 b. 28.653
32. a. 99.052 b. 99.052
33. a. 0.063 b. 0.062
34. a. 0.054 b. 0.053
35. a. 9.999 b. 9.998
36. a. 1.001 b. 1.000
37. a. 0.429 b. 0.428
38. a. 0.556 b. 0.555
39. a. 34.733 b. 34.733
40. a. 16.200 b. 16.200
41. $3+2=5$
42. $5\cdot 2=10$
43. $x+2=3\cdot 4$
44. $3+y=2+2$
45. $3y=1+2$
46. $2x=4\cdot 6$
47. $x-2=6$
48. $2-y=6$
49. $\frac{x}{2}=6$
50. $\frac{2}{x}=6$
51. $9-4+2=5+2=7$
52. $6-4+3=2+3=5$
53. $-6+4\cdot 3=-6+12=6$
54. $8-4\cdot 2=8-8=0$
55. $18-5\cdot 2=18-10=8$
56. $100-10\cdot 2=100-20=80$
57. $4+\frac{1}{3}=\frac{12+1}{3}=\frac{13}{3}$
58. $2-\frac{1}{2}=\frac{4-1}{2}=\frac{3}{2}$

Section R.1: Real Numbers

$$\begin{aligned} 59. \quad 6 - [3 \cdot 5 + 2 \cdot (3 - 2)] &= 6 - [15 + 2 \cdot (1)] \\ &= 6 - 17 \\ &= -11 \end{aligned}$$

$$\begin{aligned} 60. \quad 2 \cdot [8 - 3 \cdot (4 + 2)] - 3 &= 2 \cdot [8 - 3 \cdot (6)] - 3 \\ &= 2 \cdot [8 - 18] - 3 \\ &= 2 \cdot [-10] - 3 \\ &= -20 - 3 \\ &= -23 \end{aligned}$$

$$\begin{aligned} 61. \quad 4 \cdot (9 + 5) - 6 \cdot 7 + 3 &= 4 \cdot (14) - 42 + 3 \\ &= 56 - 42 + 3 \\ &= 14 + 3 \\ &= 17 \end{aligned}$$

$$\begin{aligned} 62. \quad 1 - (4 \cdot 3 - 2 + 2) &= 1 - (12 - 2 + 2) \\ &= 1 - 12 \\ &= -11 \end{aligned}$$

$$\begin{aligned} 63. \quad 10 - [6 - 2 \cdot 2 + (8 - 3)] \cdot 2 &= 10 - [6 - 4 + 5] \cdot 2 \\ &= 10 - [2 + 5] \cdot 2 \\ &= 10 - [7] \cdot 2 \\ &= 10 - 14 \\ &= -4 \end{aligned}$$

$$\begin{aligned} 64. \quad 2 - 5 \cdot 4 - [6 \cdot (3 - 4)] &= 2 - 20 - [6 \cdot (-1)] \\ &= -18 - [-6] \\ &= -18 + 6 \\ &= -12 \end{aligned}$$

$$65. \quad (5 - 3) \frac{1}{2} = (2) \frac{1}{2} = 1$$

$$66. \quad (5 + 4) \frac{1}{3} = (9) \frac{1}{3} = 3$$

$$67. \quad \frac{4 + 8}{5 - 3} = \frac{12}{2} = 6$$

$$68. \quad \frac{2 - 4}{5 - 3} = \frac{-2}{2} = -1$$

$$69. \quad \frac{3}{5} \cdot \frac{10}{21} = \frac{3 \cdot 2 \cdot 5}{5 \cdot 3 \cdot 7} = \frac{\cancel{3} \cdot 2 \cdot \cancel{5}}{\cancel{5} \cdot \cancel{3} \cdot 7} = \frac{2}{7}$$

$$70. \quad \frac{5}{9} \cdot \frac{3}{10} = \frac{5 \cdot 3}{3 \cdot 3 \cdot 5 \cdot 2} = \frac{\cancel{5} \cdot \cancel{3}}{\cancel{3} \cdot 3 \cdot \cancel{5} \cdot 2} = \frac{1}{6}$$

$$71. \quad \frac{6}{25} \cdot \frac{10}{27} = \frac{2 \cdot 3 \cdot 5 \cdot 2}{5 \cdot 5 \cdot 3 \cdot 9} = \frac{2 \cdot \cancel{3} \cdot \cancel{5} \cdot 2}{\cancel{5} \cdot 5 \cdot \cancel{3} \cdot 9} = \frac{4}{45}$$

$$72. \quad \frac{21}{25} \cdot \frac{100}{3} = \frac{3 \cdot 7 \cdot 4 \cdot 25}{25 \cdot 3} = \frac{\cancel{3} \cdot 7 \cdot 4 \cdot \cancel{25}}{\cancel{25} \cdot \cancel{3}} = 28$$

$$73. \quad \frac{3}{4} + \frac{2}{5} = \frac{15 + 8}{20} = \frac{23}{20}$$

$$74. \quad \frac{4}{3} + \frac{1}{2} = \frac{8 + 3}{6} = \frac{11}{6}$$

$$75. \quad \frac{7}{8} + \frac{4}{7} = \frac{49 + 32}{56} = \frac{81}{56}$$

$$76. \quad \frac{8}{9} + \frac{15}{2} = \frac{16 + 135}{18} = \frac{151}{18}$$

$$77. \quad \frac{5}{18} + \frac{1}{12} = \frac{10 + 3}{36} = \frac{13}{36}$$

$$78. \quad \frac{2}{15} + \frac{8}{9} = \frac{6 + 40}{45} = \frac{46}{45}$$

$$79. \quad \frac{5}{24} - \frac{8}{15} = \frac{25 - 64}{120} = -\frac{39}{120} = -\frac{13}{40}$$

$$80. \quad \frac{3}{14} - \frac{2}{21} = \frac{9 - 4}{42} = \frac{5}{42}$$

$$81. \quad \frac{3}{20} - \frac{2}{15} = \frac{9 - 8}{60} = \frac{1}{60}$$

$$82. \quad \frac{6}{35} - \frac{3}{14} = \frac{12 - 15}{70} = -\frac{3}{70}$$

$$83. \quad \left(\frac{5}{18} \right) = \frac{5}{18} \cdot \frac{27}{11} = \frac{5 \cdot 9 \cdot 3}{9 \cdot 2 \cdot 11} = \frac{5 \cdot \cancel{9} \cdot 3}{\cancel{9} \cdot 2 \cdot 11} = \frac{15}{22}$$

$$84. \quad \left(\frac{5}{21} \right) = \frac{5}{21} \cdot \frac{35}{2} = \frac{5 \cdot 7 \cdot 5}{7 \cdot 3 \cdot 2} = \frac{5 \cdot \cancel{7} \cdot 5}{\cancel{7} \cdot 3 \cdot 2} = \frac{25}{6}$$

Chapter R: Review

$$85. \frac{1}{3} \cdot \frac{4}{7} + \frac{17}{21} = \frac{4}{21} + \frac{17}{21} = \frac{21}{21} = 1$$

$$86. \frac{2}{3} + \frac{4}{5} \cdot \frac{1}{6} = \frac{2}{3} + \frac{2 \cdot 2}{5 \cdot 3 \cdot 2} = \frac{2}{3} + \frac{2 \cdot \cancel{2}}{5 \cdot 3 \cdot \cancel{2}} = \frac{2}{3} + \frac{2}{15}$$

$$= \frac{2}{3} \cdot \frac{5}{5} + \frac{2}{15} = \frac{10}{15} + \frac{2}{15} = \frac{10+2}{15} = \frac{12}{15}$$

$$= \frac{4 \cdot 3}{5 \cdot 3} = \frac{4 \cdot \cancel{3}}{5 \cdot \cancel{3}} = \frac{4}{5}$$

$$87. 2 \cdot \frac{3}{4} + \frac{3}{8} = \frac{2}{1} \cdot \frac{3}{4} + \frac{3}{8} = \frac{6}{4} + \frac{3}{8} = \frac{6}{4} \cdot \frac{2}{2} + \frac{3}{8}$$

$$= \frac{12}{8} + \frac{3}{8} = \frac{12+3}{8} = \frac{15}{8}$$

$$88. 3 \cdot \frac{5}{6} - \frac{1}{2} = \frac{3}{1} \cdot \frac{5}{6} - \frac{1}{2} = \frac{3 \cdot 5}{3 \cdot 2} - \frac{1}{2} = \frac{\cancel{3} \cdot 5}{\cancel{3} \cdot 2} - \frac{1}{2}$$

$$= \frac{5}{2} - \frac{1}{2} = \frac{5-1}{2} = \frac{4}{2} = 2$$

$$89. 6(x+4) = 6x+24$$

$$90. 4(2x-1) = 8x-4$$

$$91. x(x-4) = x^2-4x$$

$$92. 4x(x+3) = 4x^2+12x$$

$$93. 2\left(\frac{3}{4}x - \frac{1}{2}\right) = 2 \cdot \frac{3}{4}x - 2 \cdot \frac{1}{2} = \frac{2 \cdot 3x}{2 \cdot 2} - \frac{2}{2}$$

$$= \frac{\cancel{2} \cdot 3x}{\cancel{2} \cdot 2} - \frac{2}{2} = \frac{3}{2}x - 1$$

$$94. 3\left(\frac{2}{3}x + \frac{1}{6}\right) = 3 \cdot \frac{2}{3}x + 3 \cdot \frac{1}{6} = \frac{3 \cdot 2x}{3} + \frac{3}{3 \cdot 2}$$

$$= \frac{\cancel{3} \cdot 2x}{\cancel{3}} + \frac{\cancel{3}}{\cancel{3} \cdot 2} = 2x + \frac{1}{2}$$

$$95. (x+2)(x+4) = x^2+4x+2x+8$$

$$= x^2+6x+8$$

$$96. (x+5)(x+1) = x^2+x+5x+5$$

$$= x^2+6x+5$$

$$97. (x+9)(2x-7) = 2x^2-7x+18x-63$$

$$= 2x^2+11x-63$$

$$98. (3x-1)(x+5) = 3x^2+15x-x-5$$

$$= 3x^2+14x-5$$

$$99. (x-8)(x-2) = x^2-2x-8x+16$$

$$= x^2-10x+16$$

$$100. (x-4)(x-2) = x^2-2x-4x+8$$

$$= x^2-6x+8$$

$$101. 3x(x-5k) = 3x^2-60x$$

$$3x^2-15xk = 3x^2-60x$$

$$-15xk = -60x$$

$$k = 4$$

$$102. (x-k)(x+3k) = x^2+4x-12$$

$$x^2+3kx-kx-3k^2 = x^2+4x-12$$

$$x^2+x(3k-k)-3k^2 = x^2+4x-12$$

$$x^2+x(3k-k)-3k^2 = x^2+4x-12$$

$$x^2+x(2k)-3k^2 = x^2+4x-12$$

$$2k = 4$$

$$k = 2$$

$$103. 2x+3x = 2 \cdot x+3 \cdot x$$

$$= (2+3) \cdot x$$

$$= (5) \cdot x$$

$$= 5x$$

$$104. 2+3 \cdot 4 = 2+12 = 14$$

since multiplication comes before addition in the order of operations for real numbers.

$$(2+3) \cdot 4 = 5 \cdot 4 = 20$$

since operations inside parentheses come before multiplication in the order of operations for real numbers.

$$105. 2(3 \cdot 4) = 2(12) = 24$$

$$(2 \cdot 3) \cdot (2 \cdot 4) = (6)(8) = 48$$

$$106. \frac{4+3}{2+5} = \frac{7}{7} = 1, \text{ but}$$


$$\frac{4}{2} + \frac{3}{5} = \frac{4 \cdot 5 + 3 \cdot 2}{10} = \frac{20+6}{10} = \frac{26}{10} = \frac{13}{5} = 2.6$$


$$107. \text{ Subtraction is not commutative; for example: } 2-3 = -1 \neq 1 = 3-2.$$

Section R.2: Algebra Essentials

108. Subtraction is not associative; for example: $(5 - 2) - 1 = 2 \neq 4 = 5 - (2 - 1)$.
109. Division is not commutative; for example:
 $\frac{2}{3} \neq \frac{3}{2}$.
110. Division is not associative; for example: $(12 \div 2) \div 2 = 6 \div 2 = 3$, but $12 \div (2 \div 2) = 12 \div 1 = 12$.
111. The Symmetric Property implies that if $2 = x$, then $x = 2$.
112. From the *principle of substitution*, if $x = 5$, then
 $(x)(x) = (5)(5)$
 $\Rightarrow x^2 = 25$
 $\Rightarrow x^2 + x = 25 + 5$
 $\Rightarrow x^2 + x = 30$
113. There are no real numbers that are both rational and irrational, since an irrational number, by definition, is a number that cannot be expressed as the ratio of two integers; that is, not a rational number
 Every real number is either a rational number or an irrational number, since the decimal form of a real number either involves an infinitely repeating pattern of digits or an infinite, non-repeating string of digits.
114. The sum of an irrational number and a rational number must be irrational. Otherwise, the irrational number would then be the difference of two rational numbers, and therefore would have to be rational.
115. Answers will vary.
116. Since 1 day = 24 hours, we compute
 $\frac{12997}{24} = 541.541\bar{6}$.
 Now we only need to consider the decimal part of the answer in terms of a 24 hour day. That is,
 $(0.541\bar{6})(24) \approx 13$ hours. So it must be 13 hours later than 12 noon, which makes the time 1 AM CST.
117. Answers will vary.

Section R.2

- variable
- origin
- strict
- base; exponent (or power)
- 1.2345678×10^3
- d
- a
- b
- True
- False; the absolute value of a real number is nonnegative. $|0| = 0$ which is not a positive number.
- False; a number in scientific notation is expressed as the product of a number, x , $1 \leq x < 10$ or $-10 < x \leq -1$, and a power of 10.
- True
- 

13. A number line with tick marks at intervals of 0.5. Points are marked at -2.5, -1, 0, 0.25, $\frac{3}{4}$, 1, and $\frac{5}{2}$. Arrows point from the labels to the corresponding points on the line.
- 

14. A number line with tick marks at intervals of 0.5. Points are marked at -2, -1.5, 0, $\frac{1}{3}$, $\frac{2}{3}$, $\frac{3}{2}$, and 2. Arrows point from the labels to the corresponding points on the line.
- $\frac{1}{2} > 0$
- $5 < 6$
- $-1 > -2$
- $-3 < -\frac{5}{2}$
- $\pi > 3.14$
- $\sqrt{2} > 1.41$

Chapter R: Review

21. $\frac{1}{2} = 0.5$

22. $\frac{1}{3} > 0.33$

23. $\frac{2}{3} < 0.67$

24. $\frac{1}{4} = 0.25$

25. $x > 0$

26. $z < 0$

27. $x < 2$

28. $y > -5$

29. $x \leq 1$

30. $x \geq 2$

31. Graph on the number line: $x \geq -2$



32. Graph on the number line: $x < 4$



33. Graph on the number line: $x > -1$



34. Graph on the number line: $x \leq 7$



35. $d(C, D) = d(0, 1) = |1 - 0| = |1| = 1$

36. $d(C, A) = d(0, -3) = |-3 - 0| = |-3| = 3$

37. $d(D, E) = d(1, 3) = |3 - 1| = |2| = 2$

38. $d(C, E) = d(0, 3) = |3 - 0| = |3| = 3$

39. $d(A, E) = d(-3, 3) = |3 - (-3)| = |6| = 6$

40. $d(D, B) = d(1, -1) = |-1 - 1| = |-2| = 2$

41. $x + 2y = -2 + 2 \cdot 3 = -2 + 6 = 4$

42. $3x + y = 3(-2) + 3 = -6 + 3 = -3$

43. $5xy + 2 = 5(-2)(3) + 2 = -30 + 2 = -28$

44. $-2x + xy = -2(-2) + (-2)(3) = 4 - 6 = -2$

45. $\frac{2x}{x-y} = \frac{2(-2)}{-2-3} = \frac{-4}{-5} = \frac{4}{5}$

46. $\frac{x+y}{x-y} = \frac{-2+3}{-2-3} = \frac{1}{-5} = -\frac{1}{5}$

47. $\frac{3x+2y}{2+y} = \frac{3(-2)+2(3)}{2+3} = \frac{-6+6}{5} = \frac{0}{5} = 0$

48. $\frac{2x-3}{y} = \frac{2(-2)-3}{3} = \frac{-4-3}{3} = -\frac{7}{3}$

49. $|x+y| = |3+(-2)| = |1| = 1$

50. $|x-y| = |3-(-2)| = |5| = 5$

51. $|x|+|y| = |3|+|-2| = 3+2 = 5$

52. $|x|-|y| = |3|-|-2| = 3-2 = 1$

53. $\frac{|x|}{x} = \frac{|3|}{3} = \frac{3}{3} = 1$

54. $\frac{|y|}{y} = \frac{|-2|}{-2} = \frac{2}{-2} = -1$

55. $|4x-5y| = |4(3)-5(-2)|$
 $= |12+10|$
 $= |22|$
 $= 22$

56. $|3x+2y| = |3(3)+2(-2)| = |9-4| = |5| = 5$

57. $||4x|-|5y|| = ||4(3)|-|5(-2)||$
 $= ||12|-|10||$
 $= |12-10|$
 $= |2|$
 $= 2$

Section R.2: Algebra Essentials

$$\begin{aligned} 58. \quad 3|x| + 2|y| &= 3|3| + 2|-2| \\ &= 3 \cdot 3 + 2 \cdot 2 \\ &= 9 + 4 \\ &= 13 \end{aligned}$$

$$59. \quad \frac{x^2 - 1}{x}$$

Part (c) must be excluded. The value $x = 0$ must be excluded from the domain because it causes division by 0.

$$60. \quad \frac{x^2 + 1}{x}$$

Part (c) must be excluded. The value $x = 0$ must be excluded from the domain because it causes division by 0.

$$61. \quad \frac{x}{x^2 - 9} = \frac{x}{(x-3)(x+3)}$$

Part (a), $x = 3$, must be excluded because it causes the denominator to be 0.

$$62. \quad \frac{x}{x^2 + 9}$$

None of the given values are excluded. The domain is all real numbers.

$$63. \quad \frac{x^2}{x^2 + 1}$$

None of the given values are excluded. The domain is all real numbers.

$$64. \quad \frac{x^3}{x^2 - 1} = \frac{x^3}{(x-1)(x+1)}$$

Parts (b) and (d) must be excluded. The values $x = 1$, and $x = -1$ must be excluded from the domain because they cause division by 0.

$$65. \quad \frac{x^2 + 5x - 10}{x^3 - x} = \frac{x^2 + 5x - 10}{x(x-1)(x+1)}$$

Parts (b), (c), and (d) must be excluded. The values $x = 0$, $x = 1$, and $x = -1$ must be excluded from the domain because they cause division by 0.

$$66. \quad \frac{-9x^2 - x + 1}{x^3 + x} = \frac{-9x^2 - x + 1}{x(x^2 + 1)}$$

Part (c) must be excluded. The value $x = 0$ must

be excluded from the domain because it causes division by 0.

$$67. \quad \frac{4}{x-5}$$

$x = 5$ must be excluded because it makes the denominator equal 0.
Domain = $\{x | x \neq 5\}$

$$68. \quad \frac{-6}{x+4}$$

$x = -4$ must be excluded sine it makes the denominator equal 0.
Domain = $\{x | x \neq -4\}$

$$69. \quad \frac{x}{x+4}$$

$x = -4$ must be excluded sine it makes the denominator equal 0.
Domain = $\{x | x \neq -4\}$

$$70. \quad \frac{x-2}{x-6}$$

$x = 6$ must be excluded sine it makes the denominator equal 0.
Domain = $\{x | x \neq 6\}$

$$71. \quad C = \frac{5}{9}(F - 32) = \frac{5}{9}(32 - 32) = \frac{5}{9}(0) = 0^\circ\text{C}$$

$$72. \quad C = \frac{5}{9}(F - 32) = \frac{5}{9}(212 - 32) = \frac{5}{9}(180) = 100^\circ\text{C}$$

$$73. \quad C = \frac{5}{9}(F - 32) = \frac{5}{9}(77 - 32) = \frac{5}{9}(45) = 25^\circ\text{C}$$

$$\begin{aligned} 74. \quad C &= \frac{5}{9}(F - 32) = \frac{5}{9}(-4 - 32) \\ &= \frac{5}{9}(-36) \\ &= -20^\circ\text{C} \end{aligned}$$

$$75. \quad (-9)^2 = (-9)(-9) = 81$$

$$76. \quad -4^2 = -(4)^2 = -16$$

$$77. \quad 4^{-2} = \frac{1}{4^2} = \frac{1}{16}$$

Chapter R: Review

$$78. -4^{-2} = -\frac{1}{4^2} = -\frac{1}{16}$$

$$79. 3^{-6} \cdot 3^4 = 3^{-6+4} = 3^{-2} = \frac{1}{3^2} = \frac{1}{9}$$

$$80. 4^{-2} \cdot 4^3 = 4^{-2+3} = 4^1 = 4$$

$$81. (4^{-3})^{-1} = 4^{(-3)(-1)} = 4^3 = 64$$

$$82. (2^{-1})^{-3} = 2^{(-1)(-3)} = 2^3 = 8$$

$$83. \sqrt{100} = \sqrt{10^2} = 10$$

$$84. \sqrt{36} = \sqrt{6^2} = 6$$

$$85. \sqrt{(-4)^2} = |-4| = 4$$

$$86. \sqrt{(-3)^2} = |-3| = 3$$

$$87. (9x^4)^2 = 9^2 (x^4)^2 = 81x^8$$

$$88. (-4x^2)^{-1} = \frac{1}{-4x^2} = -\frac{1}{4x^2}$$

$$89. (x^2 y^{-1})^2 = (x^2)^2 \cdot (y^{-1})^2 = x^4 y^{-2} = \frac{x^4}{y^2}$$

$$90. (x^{-1} y)^3 = (x^{-1})^3 \cdot y^3 = x^{-3} y^3 = \frac{y^3}{x^3}$$

$$91. \frac{x^2 y^5}{x^3 y^4} = x^{2-3} y^{5-4} = x^{-1} y^1 = \frac{y}{x}$$

$$92. \frac{x^{-2} y}{x y^2} = x^{-2-1} y^{1-2} = x^{-3} y^{-1} = \frac{1}{x^3 y}$$

$$\begin{aligned} 93. \frac{(-4)^2 y^5 (xz)^3}{(-3)^3 x y^7 z^2} &= \frac{16 y^5 x^3 z^3}{-27 x y^7 z^2} \\ &= -\frac{16}{27} x^{3-1} y^{5-7} z^{3-2} \\ &= -\frac{16}{27} x^2 y^{-2} z^1 \\ &= -\frac{16x^2 z}{27y^2} \end{aligned}$$

$$\begin{aligned} 94. \frac{4x^{-2} (yz)^{-1}}{2^3 x^4 y} &= \frac{4x^{-2} y^{-1} z^{-1}}{8x^4 y} \\ &= \frac{4}{8} x^{-2-4} y^{-1-1} z^{-1} \\ &= \frac{1}{2} x^{-6} y^{-2} z^{-1} \\ &= \frac{1}{2x^6 y^2 z} \end{aligned}$$

$$95. \left(\frac{2x^{-3}}{3y^{-1}} \right)^{-2} = \left(\frac{3y}{2x^3} \right)^{-2} = \left(\frac{3x^3}{2y} \right)^2 = \frac{3^2 x^6}{2^2 y^2} = \frac{9x^6}{4y^2}$$

$$\begin{aligned} 96. \left(\frac{5x^{-2}}{6y^{-2}} \right)^{-3} &= \left(\frac{5y^2}{6x^2} \right)^{-3} = \left(\frac{6x^2}{5y^2} \right)^3 \\ &= \frac{6^3 (x^2)^3}{5^3 (y^2)^3} = \frac{216x^6}{125y^6} \end{aligned}$$

$$97. 2xy^{-1} = \frac{2x}{y} = \frac{2(2)}{(-1)} = -4$$

$$98. -3x^{-1}y = \frac{-3y}{x} = \frac{-3(-1)}{(2)} = \frac{3}{2}$$

$$99. x^2 + y^2 = (2)^2 + (-1)^2 = 4 + 1 = 5$$

$$100. x^2 y^2 = (2)^2 (-1)^2 = 4 \cdot 1 = 4$$

$$101. (xy)^2 = (2 \cdot (-1))^2 = (-2)^2 = 4$$

$$102. (x+y)^2 = (2+(-1))^2 = (1)^2 = 1$$

$$103. \sqrt{x^2} = |x| = |2| = 2$$

$$104. (\sqrt{x})^2 = x = 2$$

$$105. \sqrt{x^2 + y^2} = \sqrt{(2)^2 + (-1)^2} = \sqrt{4+1} = \sqrt{5}$$

$$106. \sqrt{x^2} + \sqrt{y^2} = |x| + |y| = |2| + |-1| = 2 + 1 = 3$$

$$107. x^y = 2^{-1} = \frac{1}{2}$$

Section R.2: Algebra Essentials

108. $y^x = (-1)^2 = 1$

109. If $x = 2$,

$$2x^3 - 3x^2 + 5x - 4 = 2 \cdot 2^3 - 3 \cdot 2^2 + 5 \cdot 2 - 4$$

$$= 16 - 12 + 10 - 4$$

$$= 10$$

If $x = 1$,

$$2x^3 - 3x^2 + 5x - 4 = 2 \cdot 1^3 - 3 \cdot 1^2 + 5 \cdot 1 - 4$$

$$= 2 - 3 + 5 - 4$$

$$= 0$$

110. If $x = 1$,

$$4x^3 + 3x^2 - x + 2 = 4 \cdot 1^3 + 3 \cdot 1^2 - 1 + 2$$

$$= 4 + 3 - 1 + 2$$

$$= 8$$

If $x = 2$,

$$4x^3 + 3x^2 - x + 2 = 4 \cdot 2^3 + 3 \cdot 2^2 - 2 + 2$$

$$= 32 + 12 - 2 + 2$$

$$= 44$$

111. $\frac{(666)^4}{(222)^4} = \left(\frac{666}{222}\right)^4 = 3^4 = 81$

112. $(0.1)^3(20)^3 = \left(\frac{1}{10}\right)^3 \cdot (2 \cdot 10)^3$

$$= \frac{1}{10^3} \cdot 2^3 \cdot 10^3$$

$$= 2^3 = 8$$

113. $(8.2)^6 \approx 304,006.671$

114. $(3.7)^5 \approx 693.440$

115. $(6.1)^{-3} \approx 0.004$

116. $(2.2)^{-5} \approx 0.019$

117. $(-2.8)^6 \approx 481.890$

118. $-(2.8)^6 \approx -481.890$

119. $(-8.11)^{-4} \approx 0.000$

120. $-(8.11)^{-4} \approx -0.000$

121. $454.2 = 4.542 \times 10^2$

122. $32.14 = 3.214 \times 10^1$

123. $0.013 = 1.3 \times 10^{-2}$

124. $0.00421 = 4.21 \times 10^{-3}$

125. $32,155 = 3.2155 \times 10^4$

126. $21,210 = 2.121 \times 10^4$

127. $0.000423 = 4.23 \times 10^{-4}$

128. $0.0514 = 5.14 \times 10^{-2}$

129. $6.15 \times 10^4 = 61,500$

130. $9.7 \times 10^3 = 9700$

131. $1.214 \times 10^{-3} = 0.001214$

132. $9.88 \times 10^{-4} = 0.000988$

133. $1.1 \times 10^8 = 110,000,000$

134. $4.112 \times 10^2 = 411.2$

135. $8.1 \times 10^{-2} = 0.081$

136. $6.453 \times 10^{-1} = 0.6453$

137. $A = lw$

138. $P = 2(l + w)$

139. $C = \pi d$

140. $A = \frac{1}{2}bh$

141. $A = \frac{\sqrt{3}}{4}x^2$

142. $P = 3x$

143. $V = \frac{4}{3}\pi r^3$

Chapter R: Review

144. $S = 4\pi r^2$

145. $V = x^3$

146. $S = 6x^2$

147. a. If $x = 1000$,
 $C = 4000 + 2x$
 $= 4000 + 2(1000)$
 $= 4000 + 2000$
 $= \$6000$
 The cost of producing 1000 watches is \$6000.

b. If $x = 2000$,
 $C = 4000 + 2x$
 $= 4000 + 2(2000)$
 $= 4000 + 4000$
 $= \$8000$
 The cost of producing 2000 watches is \$8000.

148. $210 + 80 - 120 + 25 - 60 - 32 - 5 = \98
 His balance at the end of the month was \$98.

149. We want the difference between x and 4 to be at least 6 units. Since we don't care whether the value for x is larger or smaller than 4, we take the absolute value of the difference. We want the inequality to be non-strict since we are dealing with an 'at least' situation. Thus, we have
 $|x - 4| \geq 6$

150. We want the difference between x and 2 to be more than 5 units. Since we don't care whether the value for x is larger or smaller than 2, we take the absolute value of the difference. We want the inequality to be strict since we are dealing with a 'more than' situation. Thus, we have
 $|x - 2| > 5$

151. a. $|x - 110| = |108 - 110| = |-2| = 2 \leq 5$
 108 volts is acceptable.

b. $|x - 110| = |104 - 110| = |-6| = 6 > 5$
 104 volts is not acceptable.

152. a. $|x - 220| = |214 - 220| = |-6| = 6 \leq 8$
 214 volts is acceptable.

b. $|x - 220| = |209 - 220| = |-11| = 11 > 8$
 209 volts is not acceptable.

153. a. $|x - 3| = |2.999 - 3|$
 $= |-0.001|$
 $= 0.001 \leq 0.01$
 A radius of 2.999 centimeters is acceptable.

b. $|x - 3| = |2.89 - 3|$
 $= |-0.11|$
 $= 0.11 \not\leq 0.01$
 A radius of 2.89 centimeters is not acceptable.

154. a. $|x - 98.6| = |97 - 98.6|$
 $= |-1.6|$
 $= 1.6 \geq 1.5$
 97°F is unhealthy.

b. $|x - 98.6| = |100 - 98.6|$
 $= |1.4|$
 $= 1.4 < 1.5$
 100°F is not unhealthy.

155. The distance from Earth to the Moon is about
 $4 \times 10^8 = 400,000,000$ meters.

156. The height of Mt. Everest is about
 $8848 = 8.848 \times 10^3$ meters.

157. The wavelength of visible light is about
 $5 \times 10^{-7} = 0.0000005$ meters.

158. The diameter of an atom is about
 $1 \times 10^{-10} = 0.0000000001$ meters.

159. The diameter is about $0.0403 = 4.03 \times 10^{-2}$ inches.

160. The tiniest motor is less than $0.00004 = 4 \times 10^{-5}$ millimeters tall.

Section R.3: Geometry Essentials

161. $186,000 \cdot 60 \cdot 60 \cdot 24 \cdot 365$
 $= (1.86 \times 10^5) (6 \times 10^1)^2 (2.4 \times 10^1) (3.65 \times 10^2)$
 $= 586.5696 \times 10^{10} = 5.865696 \times 10^{12}$
 There are about 5.9×10^{12} miles in one light-year.

162. $\frac{93,000,000}{186,000} = \frac{9.3 \times 10^7}{1.86 \times 10^5} = 5 \times 10^2$
 $= 500$ seconds $\approx$ 8 min. 20 sec.
 It takes about 8 minutes 20 seconds for a beam of light to reach Earth from the Sun.

163. $\frac{1}{3} = 0.333333 \dots > 0.333$
 $\frac{1}{3}$ is larger by approximately 0.0003333 ...

164. $\frac{2}{3} = 0.666666 \dots > 0.666$
 $\frac{2}{3}$ is larger by approximately 0.000666 ...

165. $(5.24 \times 10^6) (6.5 \times 10^{13}) = 34.06 \times 10^{19}$
 $= 3.406 \times 10^{20}$

166. $\frac{1.62 \times 10^{-4}}{4.5 \times 10^{-10}} = \frac{1.62}{4.5} \times \frac{10^{-4}}{10^{-10}} = 0.36 \times 10^6$
 $= 3.6 \times 10^5$

167. No. For any positive number a , the value $\frac{a}{2}$ is smaller and therefore closer to 0.

168. We are given that $1 < x^2 < 10$. This implies that $1 < x < \sqrt{10}$. Since $x < \sqrt{10} \approx 3.162$ and $x > \pi \approx 3.142$, the number could be 3.15 or 3.16 (which are between 1 and 10 as required). The number could also be 3.14 since numbers such as 3.146 which lie between π and $\sqrt{10}$ would equal 3.14 when truncated to two decimal places.

169. Answers will vary.

170. Answers will vary.
 $5 < 8$ is a true statement because 5 is further to the left than 8 on a real number line.

2. $A = \frac{1}{2}bh$

3. $C = 2\pi r$

4. similar

5. c

6. b

7. True.

8. True. $6^2 + 8^2 = 36 + 64 = 100 = 10^2$

9. False; the surface area of a sphere of radius r is given by $V = 4\pi r^2$.

10. True. The lengths of the corresponding sides are equal.

11. True. Two corresponding angles are equal.

12. False. The sides are not proportional.

13. $a = 5, b = 12,$
 $c^2 = a^2 + b^2$
 $= 5^2 + 12^2$
 $= 25 + 144$
 $= 169 \Rightarrow c = 13$

14. $a = 6, b = 8,$
 $c^2 = a^2 + b^2$
 $= 6^2 + 8^2$
 $= 36 + 64$
 $= 100 \Rightarrow c = 10$

15. $a = 10, b = 24,$
 $c^2 = a^2 + b^2$
 $= 10^2 + 24^2$
 $= 100 + 576$
 $= 676 \Rightarrow c = 26$

16. $a = 4, b = 3,$
 $c^2 = a^2 + b^2$
 $= 4^2 + 3^2$
 $= 16 + 9$
 $= 25 \Rightarrow c = 5$

Section R.3

1. right; hypotenuse

Chapter R: Review

17. $a = 7, b = 24,$

$$c^2 = a^2 + b^2$$

$$= 7^2 + 24^2$$

$$= 49 + 576$$

$$= 625 \Rightarrow c = 25$$

18. $a = 14, b = 48,$

$$c^2 = a^2 + b^2$$

$$= 14^2 + 48^2$$

$$= 196 + 2304$$

$$= 2500 \Rightarrow c = 50$$

19. $5^2 = 3^2 + 4^2$

$$25 = 9 + 16$$

$$25 = 25$$

The given triangle is a right triangle. The hypotenuse is 5.

20. $10^2 = 6^2 + 8^2$

$$100 = 36 + 64$$

$$100 = 100$$

The given triangle is a right triangle. The hypotenuse is 10.

21. $6^2 = 4^2 + 5^2$

$$36 = 16 + 25$$

$$36 = 41 \text{ false}$$

The given triangle is not a right triangle.

22. $3^2 = 2^2 + 2^2$

$$9 = 4 + 4$$

$$9 = 8 \text{ false}$$

The given triangle is not a right triangle.

23. $25^2 = 7^2 + 24^2$

$$625 = 49 + 576$$

$$625 = 625$$

The given triangle is a right triangle. The hypotenuse is 25.

24. $26^2 = 10^2 + 24^2$

$$676 = 100 + 576$$

$$676 = 676$$

The given triangle is a right triangle. The hypotenuse is 26.

25. $6^2 = 3^2 + 4^2$

$$36 = 9 + 16$$

$$36 = 25 \text{ false}$$

The given triangle is not a right triangle.

26. $7^2 = 5^2 + 4^2$

$$49 = 25 + 16$$

$$49 = 41 \text{ false}$$

The given triangle is not a right triangle.

27. $A = l \cdot w = 6 \cdot 7 = 42 \text{ in}^2$

28. $A = l \cdot w = 9 \cdot 4 = 36 \text{ cm}^2$

29. $A = \frac{1}{2}b \cdot h = \frac{1}{2}(14)(4) = 28 \text{ in}^2$

30. $A = \frac{1}{2}b \cdot h = \frac{1}{2}(4)(9) = 18 \text{ cm}^2$

31. $A = \pi r^2 = \pi(5)^2 = 25\pi \text{ m}^2$

$$C = 2\pi r = 2\pi(5) = 10\pi \text{ m}$$

32. $A = \pi r^2 = \pi(2)^2 = 4\pi \text{ ft}^2$

$$C = 2\pi r = 2\pi(2) = 4\pi \text{ ft}$$

33. $V = lwh = 6 \cdot 8 \cdot 5 = 240 \text{ ft}^3$

$$S = 2lw + 2lh + 2wh$$

$$= 2(6)(8) + 2(6)(5) + 2(8)(5)$$

$$= 96 + 60 + 80$$

$$= 236 \text{ ft}^2$$

34. $V = lwh = 9 \cdot 4 \cdot 8 = 288 \text{ in}^3$

$$S = 2lw + 2lh + 2wh$$

$$= 2(9)(4) + 2(9)(8) + 2(4)(8)$$

$$= 72 + 144 + 64$$

$$= 280 \text{ in}^2$$

35. $V = \frac{4}{3}\pi r^3 = \frac{4}{3}\pi \cdot 5^3 = \frac{500}{3}\pi \text{ cm}^3$

$$S = 4\pi r^2 = 4\pi \cdot 5^2 = 100\pi \text{ cm}^2$$

36. $V = \frac{4}{3}\pi r^3 = \frac{4}{3}\pi \cdot 3^3 = 36\pi \text{ ft}^3$

$$S = 4\pi r^2 = 4\pi \cdot 3^2 = 36\pi \text{ ft}^2$$

Section R.3: Geometry Essentials

37. $V = \pi r^2 h = \pi(9)^2(8) = 648\pi \text{ in}^3$

$$S = 2\pi r^2 + 2\pi rh$$

$$= 2\pi(9)^2 + 2\pi(9)(8)$$

$$= 162\pi + 144\pi$$

$$= 306\pi \text{ in}^2$$

38. $V = \pi r^2 h = \pi(8)^2(9) = 576\pi \text{ in}^3$

$$S = 2\pi r^2 + 2\pi rh$$

$$= 2\pi(8)^2 + 2\pi(8)(9)$$

$$= 128\pi + 144\pi$$

$$= 272\pi \text{ in}^2$$

39. The diameter of the circle is 2, so its radius is 1.

$$A = \pi r^2 = \pi(1)^2 = \pi \text{ square units}$$

40. The diameter of the circle is 2, so its radius is 1.

$$A = 2^2 - \pi(1)^2 = 4 - \pi \text{ square units}$$

41. The diameter of the circle is the length of the diagonal of the square.

$$d^2 = 2^2 + 2^2$$

$$= 4 + 4$$

$$= 8$$

$$d = \sqrt{8} = 2\sqrt{2}$$

$$r = \frac{d}{2} = \frac{2\sqrt{2}}{2} = \sqrt{2}$$

The area of the circle is:

$$A = \pi r^2 = \pi(\sqrt{2})^2 = 2\pi \text{ square units}$$

42. The diameter of the circle is the length of the diagonal of the square.

$$d^2 = 2^2 + 2^2$$

$$= 4 + 4$$

$$= 8$$

$$d = \sqrt{8} = 2\sqrt{2}$$

$$r = \frac{d}{2} = \frac{2\sqrt{2}}{2} = \sqrt{2}$$

The area is:

$$A = \pi(\sqrt{2})^2 - 2^2 = 2\pi - 4 \text{ square units}$$

43. Since the triangles are similar, the lengths of corresponding sides are proportional. Therefore, we get

$$\frac{8}{4} = \frac{x}{2}$$

$$\frac{8 \cdot 2}{4} = x$$

$$4 = x$$

In addition, corresponding angles must have the same angle measure. Therefore, we have

$$A = 90^\circ, B = 60^\circ, \text{ and } C = 30^\circ.$$

44. Since the triangles are similar, the lengths of corresponding sides are proportional. Therefore, we get

$$\frac{6}{12} = \frac{x}{16}$$

$$\frac{6 \cdot 16}{12} = x$$

$$8 = x$$

In addition, corresponding angles must have the same angle measure. Therefore, we have

$$A = 30^\circ, B = 75^\circ, \text{ and } C = 75^\circ.$$

45. Since the triangles are similar, the lengths of corresponding sides are proportional. Therefore, we get

$$\frac{30}{20} = \frac{x}{45}$$

$$\frac{30 \cdot 45}{20} = x$$

$$\frac{135}{2} = x \text{ or } x = 67.5$$

In addition, corresponding angles must have the same angle measure. Therefore, we have

$$A = 60^\circ, B = 95^\circ, \text{ and } C = 25^\circ.$$

46. Since the triangles are similar, the lengths of corresponding sides are proportional. Therefore, we get

$$\frac{8}{10} = \frac{x}{50}$$

$$\frac{8 \cdot 50}{10} = x$$

$$40 = x$$

In addition, corresponding angles must have the same angle measure. Therefore, we have

$$A = 50^\circ, B = 125^\circ, \text{ and } C = 5^\circ.$$

Chapter R: Review

47. The total distance traveled is 4 times the circumference of the wheel.
 Total Distance = $4C = 4(\pi d) = 4\pi \cdot 16$
 $= 64\pi \approx 201.1$ inches ≈ 16.8 feet

48. The distance traveled in one revolution is the circumference of the disk 4π .
 The number of revolutions =
 $\frac{\text{dist. traveled}}{\text{circumference}} = \frac{20}{4\pi} = \frac{5}{\pi} \approx 1.6$ revolutions

49. Area of the border = area of EFGH – area of ABCD
 $= 10^2 - 6^2 = 100 - 36 = 64$ ft²

50. FG = 4 feet; BG = 4 feet and BC = 10 feet, so CG = 6 feet. The area of the triangle CGF is:
 $A = \frac{1}{2} \cdot (4)(6) = 12$ ft²

51. Area of the window = area of the rectangle + area of the semicircle.

$$A = (6)(4) + \frac{1}{2} \cdot \pi \cdot 2^2 = 24 + 2\pi \approx 30.28$$
 ft²

Perimeter of the window = 2 heights + width + one-half the circumference.

$$P = 2(6) + 4 + \frac{1}{2} \cdot \pi(4) = 12 + 4 + 2\pi$$

 $= 16 + 2\pi \approx 22.28$ feet

52. Area of the deck = area of the pool and deck – area of the pool.


$$A = \pi(13)^2 - \pi(10)^2 = 169\pi - 100\pi$$

 $= 69\pi$ ft² ≈ 216.77 ft²

The amount of fence is the circumference of the circle with radius 13 feet.

$$C = 2\pi(13) = 26\pi$$
 ft ≈ 81.68 ft

53. We can form similar triangles using the Great Pyramid's height/shadow and Thales' height/shadow:



This allows us to write

$$\frac{h}{240} = \frac{2}{3}$$

$$h = \frac{2 \cdot 240}{3} = 160$$

The height of the Great Pyramid is 160 paces.

54. Let x = the approximate distance from San Juan to Hamilton and y = the approximate distance from Hamilton to Fort Lauderdale. Using similar triangles, we get

$$\frac{1046}{58} = \frac{x}{53.5} \qquad \frac{1046}{58} = \frac{y}{57}$$

$$\frac{1046 \cdot 53.5}{58} = x \qquad \frac{1046 \cdot 57}{58} = y$$

$$964.8 \approx x \qquad 1028.0 \approx y$$


The approximate distance between San Juan and Hamilton is 965 miles and the approximate distance between Hamilton and Fort Lauderdale is 1028 miles.

55. Convert 20 feet to miles, and solve the Pythagorean Theorem to find the distance:

$$20 \text{ feet} = 20 \text{ feet} \cdot \frac{1 \text{ mile}}{5280 \text{ feet}} = 0.003788 \text{ miles}$$

$$d^2 = (3960 + 0.003788)^2 - 3960^2 = 30 \text{ sq. miles}$$

$$d \approx 5.477 \text{ miles}$$




56. Convert 6 feet to miles, and solve the Pythagorean Theorem to find the distance:

$$6 \text{ feet} = 6 \text{ feet} \cdot \frac{1 \text{ mile}}{5280 \text{ feet}} = 0.001136 \text{ miles}$$

$$d^2 = (3960 + 0.001136)^2 - 3960^2 = 9 \text{ sq. miles}$$

$$d \approx 3 \text{ miles}$$



Section R.3: Geometry Essentials

57. Convert 100 feet to miles, and solve the Pythagorean Theorem to find the distance:

$$100 \text{ feet} = 100 \text{ feet} \cdot \frac{1 \text{ mile}}{5280 \text{ feet}} = 0.018939 \text{ miles}$$

$$d^2 = (3960 + 0.018939)^2 - 3960^2 \approx 150 \text{ sq. miles}$$

$$d \approx 12.2 \text{ miles}$$

Convert 150 feet to miles, and solve the Pythagorean Theorem to find the distance:

$$150 \text{ feet} = 150 \text{ feet} \cdot \frac{1 \text{ mile}}{5280 \text{ feet}} = 0.028409 \text{ miles}$$

$$d^2 = (3960 + 0.028409)^2 - 3960^2 \approx 225 \text{ sq. miles}$$

$$d \approx 15.0 \text{ miles}$$

58. Given $m > 0$, $n > 0$ and $m > n$,

if $a = m^2 - n^2$, $b = 2mn$ and $c = m^2 + n^2$, then

$$\begin{aligned} a^2 + b^2 &= (m^2 - n^2)^2 + (2mn)^2 \\ &= m^4 - 2m^2n^2 + n^4 + 4m^2n^2 \\ &= m^4 + 2m^2n^2 + n^4 \end{aligned}$$

$$\text{and } c^2 = (m^2 + n^2)^2 = m^4 + 2m^2n^2 + n^4$$

$\therefore a^2 + b^2 = c^2 \rightarrow a, b$ and c represent the sides of a right triangle.

59. $V = \pi r^2 h$

$$= \pi(10)^2(4.5)$$

$$= 450\pi \text{ ft}^3$$

So, $1 \text{ ft}^3 \approx 7.48052 \text{ gal}$ so

$$(450\pi \text{ ft}^3)(7.48052 \text{ gal/ft}^3) \approx 10,575 \text{ gal}$$

60. $10000(5.61458) = 56145.8 \text{ ft}^3$

$$V = \pi r^2 h$$

$$56145.8 = \pi(25)^2 h$$

$$h = \frac{56145.8}{625\pi} = 28.6 \text{ ft}$$

61. $A = \pi r^2$

$$A_2 = \pi(2r)^2$$

$$= \pi 4r^2$$

$$= 4\pi r^2 = 4A$$

If you double the radius, the area is four times the original area.

62. $V = \frac{4}{3}\pi r^3$

$$V_2 = \frac{4}{3}\pi(2r)^3$$

$$= \frac{4}{3}\pi \cdot 8r^3$$

$$= 8 \cdot \frac{4}{3}\pi r^3 = 8V$$

If you double the radius the volume is 8 times the original volume.

63. Let l = length of the rectangle and w = width of the rectangle.

Notice that

$$\begin{aligned} (l+w)^2 - (l-w)^2 &= [(l+w) + (l-w)][(l+w) - (l-w)] \\ &= (2l)(2w) = 4lw = 4A \end{aligned}$$

$$\text{So } A = \frac{1}{4}[(l+w)^2 - (l-w)^2]$$

Since $(l-w)^2 \geq 0$, the largest area will occur when $l-w=0$ or $l=w$; that is, when the rectangle is a square. But

$$1000 = 2l + 2w = 2(l+w)$$

$$500 = l + w = 2l$$

$$250 = l = w$$

The largest possible area is $250^2 = 62500 \text{ sq ft}$.

A circular pool with circumference = 1000 feet

yields the equation: $2\pi r = 1000 \Rightarrow r = \frac{500}{\pi}$

The area enclosed by the circular pool is:

$$A = \pi r^2 = \pi \left(\frac{500}{\pi} \right)^2 = \frac{500^2}{\pi} \approx 79577.47 \text{ ft}^2$$

Thus, a circular pool will enclose the most area.

64. Consider the diagram showing the lighthouse at point L, relative to the center of Earth, using the radius of Earth as 3960 miles. Let P refer to the furthest point on the horizon from which the light is visible. Note also that

$$362 \text{ feet} = \frac{362}{5280} \text{ miles.}$$

Section R.4: Polynomials

9. $2x^3$ Monomial; Variable: x ;
Coefficient: 2; Degree: 3
10. $-4x^2$ Monomial; Variable: x ; Coefficient:
 -4 ; Degree: 2
11. $\frac{8}{x} = 8x^{-1}$ Not a monomial; when written in
the form ax^k , the variable has a negative
exponent.
12. $-2x^{-3}$ Not a monomial; when written in the
form ax^k , the variable has a negative exponent.
13. $-2xy^2$ Monomial; Variables: x, y ;
Coefficient: -2 ; Degree: 3
14. $5x^2y^3$ Monomial; Variables: x, y ;
Coefficient: 5; Degree: 5
15. $\frac{8x}{y} = 8xy^{-1}$ Not a monomial; when written
in the form $ax^n y^m$, the exponent on the variable
 y is negative.
16. $-\frac{2x^2}{y^3} = -2x^2 y^{-3}$ Not a monomial; when
written in the form $ax^n y^m$, the exponent on the
variable y is negative.
17. $x^2 + y^2$ Not a monomial; the expression
contains more than one term. This expression is
a binomial.
18. $3x^2 + 4$ Not a monomial; the expression
contains more than one term. This expression is
a binomial.
19. $3x^2 - 5$ Polynomial; Degree: 2
20. $1 - 4x$ Polynomial; Degree: 1
21. 5 Polynomial; Degree: 0
22. $-\pi$ Polynomial; Degree: 0
23. $3x^2 - \frac{5}{x}$ Not a polynomial; the variable in the
denominator results in an exponent that is not a
nonnegative integer.
24. $\frac{3}{x} + 2$ Not a polynomial; the variable in the
denominator results in an exponent that is not a
nonnegative integer.
25. $2y^3 - \sqrt{2}$ Polynomial; Degree: 3
26. $10z^2 + z$ Polynomial; Degree: 2
27. $\frac{x^2 + 5}{x^3 - 1}$ Not a polynomial; the polynomial in
the denominator has a degree greater than 0.
28. $\frac{3x^3 + 2x - 1}{x^2 + x + 1}$ Not a polynomial; the
polynomial in the denominator has a degree
greater than 0.
29. $(x^2 + 6x + 8) + (3x^2 - 4x + 7)$
 $= (x^2 + 3x^2) + (6x - 4x) + (8 + 7)$
 $= 4x^2 + 2x + 15$
30. $(x^3 + 3x^2 + 2) + (x^2 - 4x + 4)$
 $= x^3 + (3x^2 + x^2) + (-4x) + (2 + 4)$
 $= x^3 + 4x^2 - 4x + 6$
31. $(x^3 - 2x^2 + 5x + 10) - (2x^2 - 4x + 3)$
 $= x^3 - 2x^2 + 5x + 10 - 2x^2 + 4x - 3$
 $= x^3 + (-2x^2 - 2x^2) + (5x + 4x) + (10 - 3)$
 $= x^3 - 4x^2 + 9x + 7$
32. $(x^2 - 3x - 4) - (x^3 - 3x^2 + x + 5)$
 $= x^2 - 3x - 4 - x^3 + 3x^2 - x - 5$
 $= -x^3 + (x^2 + 3x^2) + (-3x - x) + (-4 - 5)$
 $= -x^3 + 4x^2 - 4x - 9$
33. $(6x^5 + x^3 + x) + (5x^4 - x^3 + 3x^2)$
 $= 6x^5 + 5x^4 + 3x^2 + x$
34. $(10x^5 - 8x^2) + (3x^3 - 2x^2 + 6)$
 $= 10x^5 + 3x^3 - 10x^2 + 6$
35. $(x^2 - 6x + 4) + 3(2x^2 + x - 5)$
 $= x^2 - 6x + 4 + 6x^2 + 3x - 15$
 $= 7x^2 - 3x - 11$

Chapter R: Review

$$\begin{aligned} 36. \quad & -2(x^2 + x + 1) + (-5x^2 - x + 2) \\ & = -2x^2 - 2x - 2 - 5x^2 - x + 2 \\ & = -7x^2 - 3x \end{aligned}$$

$$\begin{aligned} 37. \quad & 6(x^3 + x^2 - 3) - 4(2x^3 - 3x^2) \\ & = 6x^3 + 6x^2 - 18 - 8x^3 + 12x^2 \\ & = -2x^3 + 18x^2 - 18 \end{aligned}$$

$$\begin{aligned} 38. \quad & 8(4x^3 - 3x^2 - 1) - 6(4x^3 + 8x - 2) \\ & = 32x^3 - 24x^2 - 8 - 24x^3 - 48x + 12 \\ & = 8x^3 - 24x^2 - 48x + 4 \end{aligned}$$

$$\begin{aligned} 39. \quad & (x^2 - x + 2) + (2x^2 - 3x + 5) - (x^2 + 1) \\ & = x^2 - x + 2 + 2x^2 - 3x + 5 - x^2 - 1 \\ & = 2x^2 - 4x + 6 \end{aligned}$$

$$\begin{aligned} 40. \quad & (x^2 + 1) - (4x^2 + 5) + (x^2 + x - 2) \\ & = x^2 + 1 - 4x^2 - 5 + x^2 + x - 2 \\ & = -2x^2 + x - 6 \end{aligned}$$

$$\begin{aligned} 41. \quad & 7(y^2 - 5y + 3) - 4(3 - y^2) \\ & = 7y^2 - 35y + 21 - 12 + 4y^2 \\ & = 11y^2 - 35y + 9 \end{aligned}$$

$$\begin{aligned} 42. \quad & 8(1 - y^3) + 4(1 + y + y^2 + y^3) \\ & = 8 - 8y^3 + 4 + 4y + 4y^2 + 4y^3 \\ & = -4y^3 + 4y^2 + 4y + 12 \end{aligned}$$

$$43. \quad x^2(x^2 + 2x - 5) = x^4 + 2x^3 - 5x^2$$

$$44. \quad 4x^2(x^3 - x + 2) = 4x^5 - 4x^3 + 8x^2$$

$$45. \quad -2x^2(4x^3 + 5) = -8x^5 - 10x^2$$

$$46. \quad 5x^3(3x - 4) = 15x^4 - 20x^3$$

$$\begin{aligned} 47. \quad & (x + 1)(x^2 + 2x - 4) \\ & = x(x^2 + 2x - 4) + 1(x^2 + 2x - 4) \\ & = x^3 + 2x^2 - 4x + x^2 + 2x - 4 \\ & = x^3 + 3x^2 - 2x - 4 \end{aligned}$$

$$\begin{aligned} 48. \quad & (2x - 3)(x^2 + x + 1) \\ & = 2x(x^2 + x + 1) - 3(x^2 + x + 1) \\ & = 2x^3 + 2x^2 + 2x - 3x^2 - 3x - 3 \\ & = 2x^3 - x^2 - x - 3 \end{aligned}$$

$$\begin{aligned} 49. \quad & (x + 2)(x + 4) = x^2 + 4x + 2x + 8 \\ & = x^2 + 6x + 8 \end{aligned}$$

$$\begin{aligned} 50. \quad & (x + 3)(x + 5) = x^2 + 5x + 3x + 15 \\ & = x^2 + 8x + 15 \end{aligned}$$

$$\begin{aligned} 51. \quad & (2x + 7)(x + 5) = 2x^2 + 7x + 10x + 35 \\ & = 2x^2 + 17x + 35 \end{aligned}$$

$$\begin{aligned} 52. \quad & (3x + 1)(2x + 1) = 6x^2 + 3x + 2x + 1 \\ & = 6x^2 + 5x + 1 \end{aligned}$$

$$\begin{aligned} 53. \quad & (x - 4)(x + 2) = x^2 + 2x - 4x - 8 \\ & = x^2 - 2x - 8 \end{aligned}$$

$$\begin{aligned} 54. \quad & (x + 4)(x - 2) = x^2 - 2x + 4x - 8 \\ & = x^2 + 2x - 8 \end{aligned}$$

$$\begin{aligned} 55. \quad & (x - 6)(x - 3) = x^2 - 6x - 3x + 18 \\ & = x^2 - 9x + 18 \end{aligned}$$

$$\begin{aligned} 56. \quad & (x - 5)(x - 1) = x^2 - x - 5x + 5 \\ & = x^2 - 6x + 5 \end{aligned}$$

$$\begin{aligned} 57. \quad & (2x + 3)(x - 2) = 2x^2 - 4x + 3x - 6 \\ & = 2x^2 - x - 6 \end{aligned}$$

$$\begin{aligned} 58. \quad & (2x - 4)(3x + 1) = 6x^2 + 2x - 12x - 4 \\ & = 6x^2 - 10x - 4 \end{aligned}$$

$$\begin{aligned} 59. \quad & (-3x + 4)(x - 2) = -3x^2 + 4x + 6x - 8 \\ & = -3x^2 + 10x - 8 \end{aligned}$$

$$\begin{aligned} 60. \quad & (-3x - 1)(x + 1) = -3x^2 - 3x - x - 1 \\ & = -3x^2 - 4x - 1 \end{aligned}$$

$$\begin{aligned} 61. \quad & (-x - 5)(-2x - 7) = 2x^2 + 10x + 7x + 35 \\ & = 2x^2 + 17x + 35 \end{aligned}$$

Section R.4: Polynomials

$$\begin{aligned} 62. \quad (-2x-3)(3-x) &= -6x + 2x^2 - 9 + 3x \\ &= 2x^2 - 3x - 9 \end{aligned}$$

$$\begin{aligned} 63. \quad (x-2y)(x+y) &= x^2 + xy - 2xy - 2y^2 \\ &= x^2 - xy - 2y^2 \end{aligned}$$

$$\begin{aligned} 64. \quad (2x+3y)(x-y) &= 2x^2 - 2xy + 3xy - 3y^2 \\ &= 2x^2 + xy - 3y^2 \end{aligned}$$

$$\begin{aligned} 65. \quad (-2x-3y)(3x+2y) &= -6x^2 - 4xy - 9xy - 6y^2 \\ &= -6x^2 - 13xy - 6y^2 \end{aligned}$$

$$\begin{aligned} 66. \quad (x-3y)(-2x+y) &= -2x^2 + xy + 6xy - 3y^2 \\ &= -2x^2 + 7xy - 3y^2 \end{aligned}$$

$$67. \quad (x-7)(x+7) = x^2 - 7^2 = x^2 - 49$$

$$68. \quad (x-1)(x+1) = x^2 - 1^2 = x^2 - 1$$

$$69. \quad (2x+3)(2x-3) = (2x)^2 - 3^2 = 4x^2 - 9$$

$$70. \quad (3x+2)(3x-2) = (3x)^2 - 2^2 = 9x^2 - 4$$

$$71. \quad (x+4)^2 = x^2 + 2 \cdot x \cdot 4 + 4^2 = x^2 + 8x + 16$$

$$72. \quad (x+5)^2 = x^2 + 2 \cdot x \cdot 5 + 5^2 = x^2 + 10x + 25$$

$$73. \quad (x-4)^2 = x^2 - 2 \cdot x \cdot 4 + 4^2 = x^2 - 8x + 16$$

$$74. \quad (x-5)^2 = x^2 - 2 \cdot x \cdot 5 + 5^2 = x^2 - 10x + 25$$

$$75. \quad (3x+4)(3x-4) = (3x)^2 - 4^2 = 9x^2 - 16$$

$$76. \quad (5x-3)(5x+3) = (5x)^2 - 3^2 = 25x^2 - 9$$

$$\begin{aligned} 77. \quad (2x-3)^2 &= (2x)^2 - 2(2x)(3) + 3^2 \\ &= 4x^2 - 12x + 9 \end{aligned}$$

$$\begin{aligned} 78. \quad (3x-4)^2 &= (3x)^2 - 2(3x)(4) + 4^2 \\ &= 9x^2 - 24x + 16 \end{aligned}$$

$$79. \quad (x+y)(x-y) = (x)^2 - (y)^2 = x^2 - y^2$$

$$80. \quad (x+3y)(x-3y) = (x)^2 - (3y)^2 = x^2 - 9y^2$$

$$81. \quad (3x+y)(3x-y) = (3x)^2 - (y)^2 = 9x^2 - y^2$$

$$82. \quad (3x+4y)(3x-4y) = (3x)^2 - (4y)^2 = 9x^2 - 16y^2$$

$$83. \quad (x+y)^2 = x^2 + 2xy + y^2$$

$$84. \quad (x-y)^2 = x^2 - 2xy + y^2$$

$$\begin{aligned} 85. \quad (x-2y)^2 &= x^2 + 2(x \cdot (-2y)) + (-2y)^2 \\ &= x^2 - 4xy + 4y^2 \end{aligned}$$

$$\begin{aligned} 86. \quad (2x+3y)^2 &= (2x)^2 + 2(2x \cdot 3y) + (3y)^2 \\ &= 4x^2 + 12xy + 9y^2 \end{aligned}$$

$$\begin{aligned} 87. \quad (x-2)^3 &= x^3 - 3 \cdot x^2 \cdot 2 + 3 \cdot x \cdot 2^2 - 2^3 \\ &= x^3 - 6x^2 + 12x - 8 \end{aligned}$$

$$\begin{aligned} 88. \quad (x+1)^3 &= x^3 + 3 \cdot x^2 \cdot 1 + 3 \cdot x \cdot 1^2 + 1^3 \\ &= x^3 + 3x^2 + 3x + 1 \end{aligned}$$

$$\begin{aligned} 89. \quad (2x+1)^3 &= (2x)^3 + 3(2x)^2(1) + 3(2x) \cdot 1^2 + 1^3 \\ &= 8x^3 + 12x^2 + 6x + 1 \end{aligned}$$

$$\begin{aligned} 90. \quad (3x-2)^3 &= (3x)^3 - 3(3x)^2(2) + 3(3x) \cdot 2^2 - 2^3 \\ &= 27x^3 - 54x^2 + 36x - 8 \end{aligned}$$

$$\begin{array}{r} 4x^2 - 11x + 23 \\ 91. \quad x+2 \overline{) 4x^3 - 3x^2 + x + 1} \\ \underline{4x^3 + 8x^2} \\ -11x^2 + x \\ \underline{-11x^2 - 22x} \\ 23x + 1 \\ \underline{23x + 46} \\ -45 \end{array}$$

Check:

$$\begin{aligned} (x+2)(4x^2 - 11x + 23) &+ (-45) \\ &= 4x^3 - 11x^2 + 23x + 8x^2 - 22x + 46 - 45 \\ &= 4x^3 - 3x^2 + x + 1 \end{aligned}$$

The quotient is $4x^2 - 11x + 23$; the remainder is -45 .

Chapter R: Review

$$\begin{array}{r}
 3x^2 - 7x + 15 \\
 92. \quad x + 2 \overline{) 3x^3 - x^2 + x - 2} \\
 \underline{3x^3 + 6x^2} \\
 -7x^2 + x \\
 \underline{-7x^2 - 14x} \\
 15x - 2 \\
 \underline{15x + 30} \\
 -32
 \end{array}$$

Check:

$$\begin{aligned}
 (x+2)(3x^2-7x+15) + (-32) \\
 = 3x^3 - 7x^2 + 15x + 6x^2 - 14x + 30 - 32 \\
 = 3x^3 - x^2 + x - 2
 \end{aligned}$$

The quotient is $3x^2 - 7x + 15$; the remainder is -32 .

$$\begin{array}{r}
 4x - 3 \\
 93. \quad x^2 \overline{) 4x^3 - 3x^2 + x + 1} \\
 \underline{4x^3} \\
 -3x^2 + x + 1 \\
 \underline{-3x^2} \\
 x + 1
 \end{array}$$

Check:

$$(x^2)(4x-3) + (x+1) = 4x^3 - 3x^2 + x + 1$$

The quotient is $4x - 3$; the remainder is $x + 1$.

$$\begin{array}{r}
 3x - 1 \\
 94. \quad x^2 \overline{) 3x^3 - x^2 + x - 2} \\
 \underline{3x^3} \\
 -x^2 + x - 2 \\
 \underline{-x^2} \\
 x - 2
 \end{array}$$

Check:

$$(x^2)(3x-1) + (x-2) = 3x^3 - x^2 + x - 2$$

The quotient is $3x - 1$; the remainder is $x - 2$.

$$\begin{array}{r}
 5x^2 - 13 \\
 95. \quad x^2 + 2 \overline{) 5x^4 + 0x^3 - 3x^2 + x + 1} \\
 \underline{5x^4} + 10x^2 \\
 -13x^2 + x + 1 \\
 \underline{-13x^2} - 26 \\
 x + 27
 \end{array}$$

Check:

$$\begin{aligned}
 (x^2+2)(5x^2-13) + (x+27) \\
 = 5x^4 + 10x^2 - 13x^2 - 26 + x + 27 \\
 = 5x^4 - 3x^2 + x + 1
 \end{aligned}$$

The quotient is $5x^2 - 13$; the remainder is $x + 27$.

$$\begin{array}{r}
 5x^2 - 11 \\
 96. \quad x^2 + 2 \overline{) 5x^4 + 0x^3 - x^2 + x - 2} \\
 \underline{5x^4} + 10x^2 \\
 -11x^2 + x - 2 \\
 \underline{-11x^2} - 22 \\
 x + 20
 \end{array}$$

Check:

$$\begin{aligned}
 (x^2+2)(5x^2-11) + (x+20) \\
 = 5x^4 + 10x^2 - 11x^2 - 22 + x + 20 \\
 = 5x^4 - x^2 + x - 2
 \end{aligned}$$

The quotient is $5x^2 - 11$; the remainder is $x + 20$.

$$\begin{array}{r}
 2x^2 \\
 97. \quad 2x^3 - 1 \overline{) 4x^5 + 0x^4 + 0x^3 - 3x^2 + x + 1} \\
 \underline{4x^5} - 2x^2 \\
 -x^2 + x + 1
 \end{array}$$

Check:

$$\begin{aligned}
 (2x^3-1)(2x^2) + (-x^2+x+1) \\
 = 4x^5 - 2x^2 - x^2 + x + 1 = 4x^5 - 3x^2 + x + 1
 \end{aligned}$$

The quotient is $2x^2$; the remainder is $-x^2 + x + 1$.

Section R.4: Polynomials

$$98. \quad 3x^3 - 1 \overline{) 3x^5 + 0x^4 + 0x^3 - x^2 + x - 2}$$

$$\underline{3x^5} -x^2 $$

$$ x - 2$$

Check:

$$(3x^3 - 1)(x^2) + (x - 2) = 3x^5 - x^2 + x - 2$$

The quotient is x^2 ; the remainder is $x - 2$.

$$99. \quad 2x^2 + x + 1 \overline{) 2x^4 - 3x^3 + 0x^2 + x + 1}$$

$$\underline{2x^4 + x^3 + x^2} $$

$$ -4x^3 - x^2 + x $$

$$\underline{-4x^3 - 2x^2 - 2x} $$

$$ x^2 + 3x + 1$$

$$ \underline{x^2 + \frac{1}{2}x + \frac{1}{2}}$$

$$ \frac{5}{2}x + \frac{1}{2}$$

Check:

$$(2x^2 + x + 1)(x^2 - 2x + \frac{1}{2}) + \frac{5}{2}x + \frac{1}{2}$$

$$= 2x^4 - 4x^3 + x^2 + x^3 - 2x^2 + \frac{1}{2}x$$

$$+ x^2 - 2x + \frac{1}{2} + \frac{5}{2}x + \frac{1}{2}$$

$$= 2x^4 - 3x^3 + x + 1$$

The quotient is $x^2 - 2x + \frac{1}{2}$; the remainder is

$$\frac{5}{2}x + \frac{1}{2}.$$

$$100. \quad 3x^2 + x + 1 \overline{) 3x^4 - x^3 + 0x^2 + x - 2}$$

$$\underline{3x^4 + x^3 + x^2} $$

$$ -2x^3 - x^2 + x $$

$$\underline{-2x^3 - \frac{2}{3}x^2 - \frac{2}{3}x} $$

$$ -\frac{1}{3}x^2 + \frac{5}{3}x - 2$$

$$\underline{-\frac{1}{3}x^2 - \frac{1}{9}x - \frac{1}{9}}$$

$$ \phantom{-\frac{1}{3}x^2 +} \frac{16}{9}x - \frac{17}{9}$$

Check:

$$(3x^2 + x + 1)(x^2 - \frac{2}{3}x - \frac{1}{9}) + (\frac{16}{9}x - \frac{17}{9})$$

$$= 3x^4 + x^3 + x^2 - 2x^3 - \frac{2}{3}x^2 - \frac{2}{3}x$$

$$- \frac{1}{3}x^2 - \frac{1}{9}x - \frac{1}{9} + \frac{16}{9}x - \frac{17}{9}$$

$$= 3x^4 - x^3 + x - 2$$

The quotient is $x^2 - \frac{2}{3}x - \frac{1}{9}$; the remainder is

$$\frac{16}{9}x - \frac{17}{9}.$$

$$101. \quad x - 1 \overline{) -4x^2 - 3x - 3}$$

$$\underline{-4x^3 + x^2 + 0x - 4}$$

$$ -4x^3 + 4x^2$$

$$ \underline{-3x^2}$$

$$ -3x^2 + 3x$$

$$ \underline{-3x - 4}$$

$$ \underline{-3x + 3}$$

$$ -7$$

Check:

$$(x - 1)(-4x^2 - 3x - 3) + (-7)$$

$$= -4x^3 - 3x^2 - 3x + 4x^2 + 3x + 3 - 7$$

$$= -4x^3 + x^2 - 4$$

The quotient is $-4x^2 - 3x - 3$; the remainder is -7 .

$$102. \quad x - 1 \overline{) -3x^3 - 3x^2 - 3x - 5}$$

$$\underline{-3x^4 + 3x^3}$$

$$ -3x^3$$

$$\underline{-3x^3 + 3x^2}$$

$$ -3x^2 - 2x$$

$$\underline{-3x^2 + 3x}$$

$$ -5x - 1$$

$$\underline{-5x + 5}$$

$$ -6$$

Chapter R: Review

Check:

$$\begin{aligned} & (x-1)(-3x^3-3x^2-3x-5)+(-6) \\ &= -3x^4-3x^3-3x^2-5x+3x^3+3x^2 \\ & \quad +3x+5-6 \\ &= -3x^4-2x-1 \end{aligned}$$

The quotient is $-3x^3-3x^2-3x-5$; the remainder is -6 .

$$\begin{array}{r} 103. \quad x^2+x+1 \overline{) x^4+0x^3-x^2+0x+1} \\ \underline{x^4+x^3+x^2} \\ -x^3-2x^2 \\ \underline{-x^3-x^2-x} \\ -x^2+x+1 \\ \underline{-x^2-x-1} \\ 2x+2 \end{array}$$

Check:

$$\begin{aligned} & (x^2+x+1)(x^2-x-1)+2x+2 \\ &= x^4+x^3+x^2-x^3-x^2-x-x^2-x \\ & \quad -1+2x+2 \\ &= x^4-x^2+1 \end{aligned}$$

The quotient is x^2-x-1 ; the remainder is $2x+2$.

$$\begin{array}{r} 104. \quad x^2-x+1 \overline{) x^4+0x^3-x^2+0x+1} \\ \underline{x^4-x^3+x^2} \\ x^3-2x^2 \\ \underline{x^3-x^2+x} \\ -x^2-x+1 \\ \underline{-x^2+x-1} \\ -2x+2 \end{array}$$

Check:

$$\begin{aligned} & (x^2-x+1)(x^2+x-1)+(-2x+2) \\ &= x^4+x^3-x^2-x^3-x^2+x+x^2+x \\ & \quad -1-2x+2 \\ &= x^4-x^2+1 \end{aligned}$$

The quotient is x^2+x-1 ; the remainder is $-2x+2$.

$$\begin{array}{r} 105. \quad x-a \overline{) x^3+0x^2+0x-a^3} \\ \underline{x^3-ax^2} \\ ax^2 \\ \underline{ax^2-a^2x} \\ a^2x-a^3 \\ \underline{a^2x-a^3} \\ 0 \end{array}$$

Check:

$$\begin{aligned} & (x-a)(x^2+ax+a^2)+0 \\ &= x^3+ax^2+a^2x-ax^2-a^2x-a^3 \\ &= x^3-a^3 \end{aligned}$$

The quotient is x^2+ax+a^2 ; the remainder is 0 .

$$\begin{array}{r} 106. \quad x-a \overline{) x^5+0x^4+0x^3+0x^2+0x-a^5} \\ \underline{x^5-ax^4} \\ ax^4 \\ \underline{ax^4-a^2x^3} \\ a^2x^3 \\ \underline{a^2x^3-a^3x^2} \\ a^3x^2 \\ \underline{a^3x^2-a^4x} \\ a^4x-a^5 \\ \underline{a^4x-a^5} \\ 0 \end{array}$$

Section R.5: Factoring Polynomials

Check:

$$\begin{aligned}(x-a)(x^4+ax^3+a^2x^2+a^3x+a^4)+0 \\&= x^5+ax^4+a^2x^3+a^3x^2+a^4x-ax^4 \\&\quad -a^2x^3-a^3x^2-a^4x-a^5 \\&= x^5-a^5\end{aligned}$$

The quotient is $x^4+ax^3+a^2x^2+a^3x+a^4$; the remainder is 0.

- 107.** $(3x-2k)(4x+3k)=12x^2+kx-96$
 $12x^2-8kx+9kx-6k^2=12x^2+kx-96$
 $kx-6k^2=kx-96$
 $-6k^2=-96$
 $k^2=16$
 $k=\pm 4$
- 108.** The products $(x+y)(x-y)$ and $(z+w)(z-w)$ will each result in a binomial that is the difference of squares. The product of those resulting binomials will have 4 terms.
- 109.** When we multiply polynomials $p_1(x)$ and $p_2(x)$, each term of $p_1(x)$ will be multiplied by each term of $p_2(x)$. So when the highest-powered term of $p_1(x)$ multiplies by the highest powered term of $p_2(x)$, the exponents on the variables in those terms will add according to the basic rules of exponents. Therefore, the highest powered term of the product polynomial will have degree equal to the sum of the degrees of $p_1(x)$ and $p_2(x)$.
- 110.** When we add two polynomials $p_1(x)$ and $p_2(x)$, where the degree of $p_1(x) \neq$ the degree of $p_2(x)$, each term of $p_1(x)$ will be added to each term of $p_2(x)$. Since only the terms with equal degrees will combine via addition, the degree of the sum polynomial will be the degree of the highest powered term overall, that is, the degree of the polynomial that had the higher degree.
- 111.** When we add two polynomials $p_1(x)$ and $p_2(x)$, where the degree of $p_1(x) =$ the degree

of $p_2(x)$, the new polynomial will have degree $\leq$ the degree of $p_1(x)$ and $p_2(x)$.

112. Answers will vary.

113. Answers will vary.

Section R.5

1. $3x(x-2)(x+2)$
2. prime
3. c
4. b
5. d
6. c
7. True; x^2+4 is prime over the set of real numbers.
8. False; $3x^3-2x^2-6x+4=(3x-2)(x^2-2)$
9. $3x+6=3(x+2)$
10. $7x-14=7(x-2)$
11. $ax^2+a=a(x^2+1)$
12. $ax-a=a(x-1)$
13. $x^3+x^2+x=x(x^2+x+1)$
14. $x^3-x^2+x=x(x^2-x+1)$
15. $2x^2-2x=2x(x-1)$
16. $3x^2-3x=3x(x-1)$
17. $3x^2y-6xy^2+12xy=3xy(x-2y+4)$
18. $60x^2y-48xy^2+72x^3y=12xy(5x-4y+6x^2)$
19. $x^2-1=x^2-1^2=(x-1)(x+1)$

Chapter R: Review

$$20. \quad x^2 - 4 = x^2 - 2^2 = (x-2)(x+2)$$

$$21. \quad 4x^2 - 1 = (2x)^2 - 1^2 = (2x-1)(2x+1)$$

$$22. \quad 9x^2 - 1 = (3x)^2 - 1^2 = (3x-1)(3x+1)$$

$$23. \quad x^2 - 16 = x^2 - 4^2 = (x-4)(x+4)$$

$$24. \quad x^2 - 25 = x^2 - 5^2 = (x-5)(x+5)$$

$$25. \quad 25x^2 - 4 = (5x-2)(5x+2)$$

$$26. \quad 36x^2 - 9 = 9(4x^2 - 1) = 9(2x-1)(2x+1)$$

$$27. \quad x^2 + 2x + 1 = (x+1)^2$$

$$28. \quad x^2 - 4x + 4 = (x-2)^2$$

$$29. \quad x^2 + 4x + 4 = (x+2)^2$$

$$30. \quad x^2 - 2x + 1 = (x-1)^2$$

$$31. \quad x^2 - 10x + 25 = (x-5)^2$$

$$32. \quad x^2 + 10x + 25 = (x+5)^2$$

$$33. \quad 4x^2 + 4x + 1 = (2x+1)^2$$

$$34. \quad 9x^2 + 6x + 1 = (3x+1)^2$$

$$35. \quad 16x^2 + 8x + 1 = (4x+1)^2$$

$$36. \quad 25x^2 + 10x + 1 = (5x+1)^2$$

$$37. \quad x^3 - 27 = x^3 - 3^3 = (x-3)(x^2 + 3x + 9)$$

$$38. \quad x^3 + 125 = x^3 + 5^3 = (x+5)(x^2 - 5x + 25)$$

$$39. \quad x^3 + 27 = x^3 + 3^3 = (x+3)(x^2 - 3x + 9)$$

$$\begin{aligned} 40. \quad 27 - 8x^3 &= 3^3 - (2x)^3 \\ &= (3-2x)(9+6x+4x^2) \\ &= -(2x-3)(4x^2+6x+9) \end{aligned}$$

$$\begin{aligned} 41. \quad 8x^3 + 27 &= (2x)^3 + 3^3 \\ &= (2x+3)(4x^2 - 6x + 9) \end{aligned}$$

$$\begin{aligned} 42. \quad 64 - 27x^3 &= 4^3 - (3x)^3 \\ &= (4-3x)(16+12x+9x^2) \\ &= -(3x-4)(9x^2+12x+16) \end{aligned}$$

$$43. \quad x^2 + 5x + 6 = (x+2)(x+3)$$

$$44. \quad x^2 + 6x + 8 = (x+2)(x+4)$$

$$45. \quad x^2 + 7x + 6 = (x+6)(x+1)$$

$$46. \quad x^2 + 9x + 8 = (x+8)(x+1)$$

$$47. \quad x^2 + 7x + 10 = (x+2)(x+5)$$

$$48. \quad x^2 + 11x + 10 = (x+10)(x+1)$$

$$49. \quad x^2 - 10x + 16 = (x-2)(x-8)$$

$$50. \quad x^2 - 17x + 16 = (x-16)(x-1)$$

$$51. \quad x^2 - 7x - 8 = (x+1)(x-8)$$

$$52. \quad x^2 - 2x - 8 = (x+2)(x-4)$$

$$53. \quad x^2 + 7x - 8 = (x+8)(x-1)$$

$$54. \quad x^2 + 2x - 8 = (x+4)(x-2)$$

$$\begin{aligned} 55. \quad 2x^2 + 4x + 3x + 6 &= 2x(x+2) + 3(x+2) \\ &= (x+2)(2x+3) \end{aligned}$$

$$\begin{aligned} 56. \quad 3x^2 - 3x + 2x - 2 &= 3x(x-1) + 2(x-1) \\ &= (x-1)(3x+2) \end{aligned}$$

$$\begin{aligned} 57. \quad 5x^2 - 15x + x - 3 &= 5x(x-3) + 1(x-3) \\ &= (x-3)(5x+1) \end{aligned}$$

$$\begin{aligned} 58. \quad 3x^2 + 6x - x - 2 &= 3x(x+2) - 1(x+2) \\ &= (x+2)(3x-1) \end{aligned}$$

Section R.5: Factoring Polynomials

$$\begin{aligned} 59. \quad 6x^2 + 21x + 8x + 28 &= 3x(2x + 7) + 4(2x + 7) \\ &= (2x + 7)(3x + 4) \end{aligned}$$

$$\begin{aligned} 60. \quad 9x^2 - 6x + 3x - 2 &= 3x(3x - 2) + 1(3x - 2) \\ &= (3x - 2)(3x + 1) \end{aligned}$$

$$61. \quad 3x^2 + 4x + 1 = (3x + 1)(x + 1)$$

$$62. \quad 2x^2 + 3x + 1 = (2x + 1)(x + 1)$$

$$63. \quad 2z^2 + 9z + 7 = (2z + 7)(z + 1)$$

$$64. \quad 6z^2 + 5z + 1 = (3z + 1)(2z + 1)$$

$$65. \quad 5x^2 + 6x - 8 = (5x - 4)(x + 2)$$

$$66. \quad 3x^2 + 10x + 8 = (3x + 4)(x + 2)$$

$$67. \quad 5x^2 - 6x - 8 = (5x + 4)(x - 2)$$

$$68. \quad 3x^2 - 10x + 8 = (3x - 4)(x - 2)$$

$$69. \quad 5x^2 + 22x + 8 = (5x + 2)(x + 4)$$

$$70. \quad 3x^2 - 14x + 8 = (3x - 2)(x - 4)$$

$$71. \quad 5x^2 + 18x - 8 = (5x - 2)(x + 4)$$

$$72. \quad 3x^2 - 10x - 8 = (3x + 2)(x - 4)$$

$$73. \quad \text{Since } b \text{ is } 10 \text{ then we need half of } 10 \text{ squared to be the last term in our trinomial. Thus}$$

$$\frac{1}{2}(10) = 5; (5)^2 = 25$$

$$x^2 + 10x + 25 = (x + 5)^2$$

$$74. \quad \text{Since } b \text{ is } 14 \text{ then we need half of } 14 \text{ squared to be the last term in our trinomial. Thus}$$

$$\frac{1}{2}(14) = 7; (7)^2 = 49$$

$$p^2 + 14p + 49 = (p + 7)^2$$

$$75. \quad \text{Since } b \text{ is } -6 \text{ then we need half of } -6 \text{ squared to be the last term in our trinomial. Thus}$$

$$\frac{1}{2}(-6) = -3; (-3)^2 = 9$$

$$y^2 - 6y + 9 = (y - 3)^2$$

$$76. \quad \text{Since } b \text{ is } -4 \text{ then we need half of } -4 \text{ squared to be the last term in our trinomial. Thus}$$

$$\frac{1}{2}(-4) = -2; (-2)^2 = 4$$

$$x^2 - 4x + 4 = (x - 2)^2$$

$$77. \quad \text{Since } b \text{ is } -\frac{1}{2} \text{ then we need half of } -\frac{1}{2} \text{ squared to be the last term in our trinomial. Thus}$$

$$\frac{1}{2}\left(-\frac{1}{2}\right) = -\frac{1}{4}; \left(-\frac{1}{4}\right)^2 = \frac{1}{16}$$

$$x^2 - \frac{1}{2}x + \frac{1}{16} = \left(x - \frac{1}{4}\right)^2$$

$$78. \quad \text{Since } b \text{ is } \frac{1}{3} \text{ then we need half of } \frac{1}{3} \text{ squared to be the last term in our trinomial. Thus}$$

$$\frac{1}{2}\left(\frac{1}{3}\right) = \frac{1}{6}; \left(\frac{1}{6}\right)^2 = \frac{1}{36}$$

$$x^2 + \frac{1}{3}x + \frac{1}{36} = \left(x + \frac{1}{6}\right)^2$$

$$79. \quad x^2 - 36 = (x - 6)(x + 6)$$

$$80. \quad x^2 - 9 = (x - 3)(x + 3)$$

$$81. \quad 2 - 8x^2 = 2(1 - 4x^2) = 2(1 - 2x)(1 + 2x)$$

$$82. \quad 3 - 27x^2 = 3(1 - 9x^2) = 3(1 - 3x)(1 + 3x)$$

$$83. \quad x^2 + 11x + 10 = (x + 1)(x + 10)$$

$$84. \quad x^2 + 5x + 4 = (x + 4)(x + 1)$$

$$85. \quad x^2 - 10x + 21 = (x - 7)(x - 3)$$

$$86. \quad x^2 - 6x + 8 = (x - 2)(x - 4)$$

$$87. \quad 4x^2 - 8x + 32 = 4(x^2 - 2x + 8)$$

$$88. \quad 3x^2 - 12x + 15 = 3(x^2 - 4x + 5)$$

$$89. \quad x^2 + 4x + 16 \text{ is prime over the reals because there are no factors of } 16 \text{ whose sum is } 4.$$

$$90. \quad x^2 + 12x + 36 = (x + 6)^2$$

$$91. \quad 15 + 2x - x^2 = -(x^2 - 2x - 15) = -(x - 5)(x + 3)$$

Chapter R: Review

92. $14 + 6x - x^2 = -(x^2 - 6x - 14)$ is prime over the integers because there are no factors of -14 whose sum is -6 .
93. $3x^2 - 12x - 36 = 3(x^2 - 4x - 12)$
 $= 3(x - 6)(x + 2)$
94. $x^3 + 8x^2 - 20x = x(x^2 + 8x - 20)$
 $= x(x + 10)(x - 2)$
95. $y^4 + 11y^3 + 30y^2 = y^2(y^2 + 11y + 30)$
 $= y^2(y + 5)(y + 6)$
96. $3y^3 - 18y^2 - 48y = 3y(y^2 - 6y - 16)$
 $= 3y(y + 2)(y - 8)$
97. $4x^2 + 12x + 9 = (2x + 3)^2$
98. $9x^2 - 12x + 4 = (3x - 2)^2$
99. $6x^2 + 8x + 2 = 2(3x^2 + 4x + 1)$
 $= 2(3x + 1)(x + 1)$
100. $8x^2 + 6x - 2 = 2(4x^2 + 3x - 1)$
 $= 2(4x - 1)(x + 1)$
101. $x^4 - 81 = (x^2)^2 - 9^2 = (x^2 - 9)(x^2 + 9)$
 $= (x - 3)(x + 3)(x^2 + 9)$
102. $x^4 - 1 = (x^2)^2 - 1^2 = (x^2 - 1)(x^2 + 1)$
 $= (x - 1)(x + 1)(x^2 + 1)$
103. $x^6 - 2x^3 + 1 = (x^3 - 1)^2$
 $= [(x - 1)(x^2 + x + 1)]^2$
 $= (x - 1)^2(x^2 + x + 1)^2$
104. $x^6 + 2x^3 + 1 = (x^3 + 1)^2$
 $= [(x + 1)(x^2 - x + 1)]^2$
 $= (x + 1)^2(x^2 - x + 1)^2$
105. $x^7 - x^5 = x^5(x^2 - 1) = x^5(x - 1)(x + 1)$
106. $x^8 - x^5 = x^5(x^3 - 1) = x^5(x - 1)(x^2 + x + 1)$
107. $16x^2 + 24x + 9 = (4x + 3)^2$
108. $9x^2 - 24x + 16 = (3x - 4)^2$
109. $5 + 16x - 16x^2 = -(16x^2 - 16x - 5)$
 $= -(4x - 5)(4x + 1)$
110. $5 + 11x - 16x^2 = -(16x^2 - 11x - 5)$
 $= -(16x + 5)(x - 1)$
111. $4y^2 - 16y + 15 = (2y - 5)(2y - 3)$
112. $9y^2 + 9y - 4 = (3y + 4)(3y - 1)$
113. $1 - 8x^2 - 9x^4 = -(9x^4 + 8x^2 - 1)$
 $= -(9x^2 - 1)(x^2 + 1)$
 $= -(3x - 1)(3x + 1)(x^2 + 1)$
114. $4 - 14x^2 - 8x^4 = -2(4x^4 + 7x^2 - 2)$
 $= -2(4x^2 - 1)(x^2 + 2)$
 $= -2(2x - 1)(2x + 1)(x^2 + 2)$
115. $x(x + 3) - 6(x + 3) = (x + 3)(x - 6)$
116. $5(3x - 7) + x(3x - 7) = (3x - 7)(x + 5)$
117. $(x + 2)^2 - 5(x + 2) = (x + 2)[(x + 2) - 5]$
 $= (x + 2)(x - 3)$
118. $(x - 1)^2 - 2(x - 1) = (x - 1)[(x - 1) - 2]$
 $= (x - 1)(x - 3)$
119. $(3x - 2)^3 - 27$
 $= (3x - 2)^3 - 3^3$
 $= [(3x - 2) - 3][(3x - 2)^2 + 3(3x - 2) + 9]$
 $= (3x - 5)(9x^2 - 12x + 4 + 9x - 6 + 9)$
 $= (3x - 5)(9x^2 - 3x + 7)$

Section R.5: Factoring Polynomials

$$\begin{aligned} 120. \quad & (5x+1)^3 - 1 \\ &= (5x+1)^3 - 1^3 \\ &= [(5x+1)-1][(5x+1)^2 + (1)(5x+1)+1] \\ &= 5x(25x^2 + 10x + 1 + 5x + 1 + 1) \\ &= 5x(25x^2 + 15x + 3) \end{aligned}$$

$$\begin{aligned} 121. \quad & 3(x^2 + 10x + 25) - 4(x+5) \\ &= 3(x+5)^2 - 4(x+5) \\ &= (x+5)[3(x+5) - 4] \\ &= (x+5)(3x+15-4) \\ &= (x+5)(3x+11) \end{aligned}$$

$$\begin{aligned} 122. \quad & 7(x^2 - 6x + 9) + 5(x-3) \\ &= 7(x-3)^2 + 5(x-3) \\ &= (x-3)[7(x-3) + 5] \\ &= (x-3)(7x-21+5) \\ &= (x-3)(7x-16) \end{aligned}$$

$$\begin{aligned} 123. \quad & x^3 + 2x^2 - x - 2 = x^2(x+2) - 1(x+2) \\ &= (x+2)(x^2 - 1) \\ &= (x+2)(x-1)(x+1) \end{aligned}$$

$$\begin{aligned} 124. \quad & x^3 - 3x^2 - x + 3 = x^2(x-3) - 1(x-3) \\ &= (x-3)(x^2 - 1) \\ &= (x-3)(x-1)(x+1) \end{aligned}$$

$$\begin{aligned} 125. \quad & x^4 - x^3 + x - 1 = x^3(x-1) + 1(x-1) \\ &= (x-1)(x^3 + 1) \\ &= (x-1)(x+1)(x^2 - x + 1) \end{aligned}$$

$$\begin{aligned} 126. \quad & x^4 + x^3 + x + 1 = x^3(x+1) + 1(x+1) \\ &= (x+1)(x^3 + 1) \\ &= (x+1)(x+1)(x^2 - x + 1) \\ &= (x+1)^2(x^2 - x + 1) \end{aligned}$$

$$\begin{aligned} 127. \quad & 2(3x+4)^2 + (2x+3) \cdot 2(3x+4) \cdot 3 \\ &= 2(3x+4)((3x+4) + (2x+3) \cdot 3) \\ &= 2(3x+4)(3x+4+6x+9) \\ &= 2(3x+4)(9x+13) \end{aligned}$$

$$\begin{aligned} 128. \quad & 5(2x+1)^2 + (5x-6) \cdot 2(2x+1) \cdot 2 \\ &= (2x+1)(5(2x+1) + (5x-6) \cdot 4) \\ &= (2x+1)(10x+5+20x-24) \\ &= (2x+1)(30x-19) \end{aligned}$$

$$\begin{aligned} 129. \quad & 2x(2x+5) + x^2 \cdot 2 = 2x((2x+5) + x) \\ &= 2x(2x+5+x) \\ &= 2x(3x+5) \end{aligned}$$

$$\begin{aligned} 130. \quad & 3x^2(8x-3) + x^3 \cdot 8 = x^2(3(8x-3) + 8x) \\ &= x^2(24x-9+8x) \\ &= x^2(32x-9) \end{aligned}$$

$$\begin{aligned} 131. \quad & 2(x+3)(x-2)^3 + (x+3)^2 \cdot 3(x-2)^2 \\ &= (x+3)(x-2)^2(2(x-2) + (x+3) \cdot 3) \\ &= (x+3)(x-2)^2(2x-4+3x+9) \\ &= (x+3)(x-2)^2(5x+5) \\ &= 5(x+3)(x-2)^2(x+1) \end{aligned}$$

$$\begin{aligned} 132. \quad & 4(x+5)^3(x-1)^2 + (x+5)^4 \cdot 2(x-1) \\ &= 2(x+5)^3(x-1)(2(x-1) + (x+5)) \\ &= 2(x+5)^3(x-1)(2x-2+x+5) \\ &= 2(x+5)^3(x-1)(3x+3) \\ &= 2 \cdot 3(x+5)^3(x-1)(x+1) \\ &= 6(x+5)^3(x-1)(x+1) \end{aligned}$$

$$\begin{aligned} 133. \quad & (4x-3)^2 + x \cdot 2(4x-3) \cdot 4 \\ &= (4x-3)((4x-3) + 8x) \\ &= (4x-3)(4x-3+8x) \\ &= (4x-3)(12x-3) \\ &= 3(4x-3)(4x-1) \end{aligned}$$

Chapter R: Review

- 134.** $3x^2(3x+4)^2 + x^3 \cdot 2(3x+4) \cdot 3$
 $= 3x^2(3x+4)((3x+4)+2x)$
 $= 3x^2(3x+4)(3x+4+2x)$
 $= 3x^2(3x+4)(5x+4)$
- 135.** $2(3x-5) \cdot 3(2x+1)^3 + (3x-5)^2 \cdot 3(2x+1)^2 \cdot 2$
 $= 6(3x-5)(2x+1)^2((2x+1)+(3x-5))$
 $= 6(3x-5)(2x+1)^2(2x+1+3x-5)$
 $= 6(3x-5)(2x+1)^2(5x-4)$
- 136.** $3(4x+5)^2 \cdot 4(5x+1)^2 + (4x+5)^3 \cdot 2(5x+1) \cdot 5$
 $= 2(4x+5)^2(5x+1)(6(5x+1)+5(4x+5))$
 $= 2(4x+5)^2(5x+1)(30x+6+20x+25)$
 $= 2(4x+5)^2(5x+1)(50x+31)$
- 137.** The possible factorizations are
 $(x \pm 1)(x \pm 4) = x^2 \pm 5x + 4$ or
 $(x \pm 2)(x \pm 2) = x^2 \pm 4x + 4$, none of which
equals $x^2 + 4$.
- 138.** The possible factorizations are
 $(x \pm 1)^2 = x^2 \pm 2x + 1$, neither of which equals
 $x^2 + x + 1$.
- 139.** Answers will vary.
- 140.** Answers will vary.

Section R.6

1. quotient; divisor; remainder

$$2. \begin{array}{r} -3 \overline{) 2 \ 0 \ -5 \ 1} \end{array}$$

3. d

4. a

5. True

6. True

$$7. \begin{array}{r} 2 \overline{) 1 \ -7 \ 5 \ 10} \\ \underline{2 \ -10 \ -10} \\ 1 \ -5 \ -5 \ 0 \end{array}$$

Quotient: $x^2 - 5x - 5$

Remainder: 0

$$8. \begin{array}{r} -1 \overline{) 1 \ 2 \ -3 \ 1} \\ \underline{-1 \ -1 \ 4} \\ 1 \ 1 \ -4 \ 5 \end{array}$$

Quotient: $x^2 + x - 4$

Remainder: 5

$$9. \begin{array}{r} 3 \overline{) 3 \ 2 \ -1 \ 3} \\ \underline{9 \ 33 \ 96} \\ 3 \ 11 \ 32 \ 99 \end{array}$$

Quotient: $3x^2 + 11x + 32$

Remainder: 99

$$10. \begin{array}{r} -2 \overline{) -4 \ 2 \ -1 \ 1} \\ \underline{8 \ -20 \ 42} \\ -4 \ 10 \ -21 \ 43 \end{array}$$

Quotient: $-4x^2 + 10x - 21$

Remainder: 43

$$11. \begin{array}{r} -3 \overline{) 1 \ 0 \ -4 \ 0 \ 1 \ 0} \\ \underline{-3 \ 9 \ -15 \ 45 \ -138} \\ 1 \ -3 \ 5 \ -15 \ 46 \ -138 \end{array}$$

Quotient: $x^4 - 3x^3 + 5x^2 - 15x + 46$

Remainder: -138

Section R.6: Synthetic Division

$$\begin{array}{r} 12. \quad 2 \overline{) 1 \ 0 \ 1 \ 0 \ 2} \\ \underline{2 \ 4 \ 10 \ 20} \\ 1 \ 2 \ 5 \ 10 \ 22 \end{array}$$

Quotient: $x^3 + 2x^2 + 5x + 10$
Remainder: 22

$$\begin{array}{r} 13. \quad 1 \overline{) 4 \ 0 \ -3 \ 0 \ 1 \ 0 \ 5} \\ \underline{4 \ 4 \ 1 \ 1 \ 2 \ 2} \\ 4 \ 4 \ 1 \ 1 \ 2 \ 2 \ 7 \end{array}$$

Quotient: $4x^5 + 4x^4 + x^3 + x^2 + 2x + 2$
Remainder: 7

$$\begin{array}{r} 14. \quad -1 \overline{) 1 \ 0 \ 5 \ 0 \ 0 \ -10} \\ \underline{-1 \ 1 \ -6 \ 6 \ -6} \\ 1 \ -1 \ 6 \ -6 \ 6 \ -16 \end{array}$$

Quotient: $x^4 - x^3 + 6x^2 - 6x + 6$
Remainder: -16

$$\begin{array}{r} 15. \quad -1.1 \overline{) 0.1 \ 0 \ 0.2 \ 0} \\ \underline{-0.11 \ 0.121 \ -0.3531} \\ 0.1 \ -0.11 \ 0.321 \ -0.3531 \end{array}$$

Quotient: $0.1x^2 - 0.11x + 0.321$
Remainder: -0.3531

$$\begin{array}{r} 16. \quad -2.1 \overline{) 0.1 \ 0 \ -0.2} \\ \underline{-0.21 \ 0.441} \\ 0.1 \ -0.21 \ 0.241 \end{array}$$

Quotient: $0.1x - 0.21$
Remainder: 0.241

$$\begin{array}{r} 17. \quad 2 \overline{) 1 \ 0 \ 0 \ 0 \ 0 \ -32} \\ \underline{2 \ 4 \ 8 \ 16 \ 32} \\ 1 \ 2 \ 4 \ 8 \ 16 \ 0 \end{array}$$

Quotient: $x^4 + 2x^3 + 4x^2 + 8x + 16$
Remainder: 0

$$\begin{array}{r} 18. \quad -1 \overline{) 1 \ 0 \ 0 \ 0 \ 0 \ 1} \\ \underline{-1 \ 1 \ -1 \ 1 \ -1} \\ 1 \ -1 \ 1 \ -1 \ 1 \ 0 \end{array}$$

Quotient: $x^4 - x^3 + x^2 - x + 1$
Remainder: 0

$$\begin{array}{r} 19. \quad 2 \overline{) 4 \ -3 \ -8 \ 4} \\ \underline{8 \ 10 \ 4} \\ 4 \ 5 \ 2 \ 8 \end{array}$$

Remainder = $8 \neq 0$. Therefore, $x - 2$ is not a factor of $4x^3 - 3x^2 - 8x + 4$.

$$\begin{array}{r} 20. \quad -3 \overline{) -4 \ 5 \ 0 \ 8} \\ \underline{12 \ -51 \ 153} \\ -4 \ 17 \ -51 \ 161 \end{array}$$

Remainder = $161 \neq 0$. Therefore, $x + 3$ is not a factor of $-4x^3 + 5x^2 + 8$.

$$\begin{array}{r} 21. \quad 3 \overline{) 2 \ -6 \ 0 \ -7 \ 21} \\ \underline{6 \ 0 \ 0 \ -21} \\ 2 \ 0 \ 0 \ -7 \ 0 \end{array}$$

Remainder = 0. Therefore, $x - 3$ is a factor of $2x^4 - 6x^3 - 7x + 21$.

$$\begin{array}{r} 22. \quad 2 \overline{) 4 \ 0 \ -15 \ 0 \ -4} \\ \underline{8 \ 16 \ 2 \ 4} \\ 4 \ 8 \ 1 \ 2 \ 0 \end{array}$$

Remainder = 0. Therefore, $x - 2$ is a factor of $4x^4 - 15x^2 - 4$.

$$\begin{array}{r} 23. \quad -2 \overline{) 5 \ 0 \ 0 \ 43 \ 0 \ 0 \ 24} \\ \underline{-10 \ 20 \ -40 \ -6 \ 12 \ -24} \\ 3 \ -10 \ 20 \ 3 \ -6 \ 12 \ 0 \end{array}$$

Remainder = 0. Therefore, $x + 3$ is a factor of $5x^6 + 43x^3 + 24$.

$$\begin{array}{r} 24. \quad -3 \overline{) 2 \ 0 \ -18 \ 0 \ 1 \ 0 \ -9} \\ \underline{-6 \ 18 \ 0 \ 0 \ -3 \ 9} \\ 2 \ -6 \ 0 \ 0 \ 1 \ -3 \ 0 \end{array}$$

Remainder = 0. Therefore, $x + 2$ is a factor of $2x^6 - 18x^4 + x^2 - 9$.

Chapter R: Review

$$\begin{array}{r} 25. \quad -4 \overline{) 1 \quad 0 \quad -16 \quad -1 \quad 0 \quad 19} \\ \underline{-4 \quad 16 \quad 0 \quad 4 \quad -16} \\ 1 \quad -4 \quad 0 \quad -1 \quad 4 \quad 3 \end{array}$$

Remainder = $1 \neq 0$. Therefore, $x+4$ is not a factor of $x^5 - 16x^3 - x^2 + 19$.

$$\begin{array}{r} 26. \quad -4 \overline{) 1 \quad 0 \quad -16 \quad 0 \quad 1 \quad 0 \quad -16} \\ \underline{-4 \quad 16 \quad 0 \quad 0 \quad -4 \quad 16} \\ 1 \quad -4 \quad 0 \quad 0 \quad 1 \quad -4 \quad 0 \end{array}$$

Remainder = 0. Therefore, $x+4$ is a factor of $x^6 - 16x^4 + x^2 - 16$.

$$\begin{array}{r} 27. \quad \frac{1}{3} \overline{) 3 \quad -1 \quad 0 \quad 6 \quad -2} \\ \underline{1 \quad 0 \quad 0 \quad 2} \\ 3 \quad 0 \quad 0 \quad 6 \quad 0 \end{array}$$

Remainder = 0; therefore $x - \frac{1}{3}$ is a factor of $3x^4 - x^3 + 6x - 2$.

$$\begin{array}{r} 28. \quad -\frac{1}{3} \overline{) 3 \quad 1 \quad 0 \quad -3 \quad 1} \\ \underline{-1 \quad 0 \quad 0 \quad 1} \\ 3 \quad 0 \quad 0 \quad -3 \quad 2 \end{array}$$

Remainder = $2 \neq 0$; therefore $x + \frac{1}{3}$ is not a factor of $3x^4 + x^3 - 3x + 1$.

$$\begin{array}{r} 29. \quad -2 \overline{) 1 \quad -2 \quad 3 \quad 5} \\ \underline{-2 \quad 8 \quad -22} \\ 1 \quad -4 \quad 11 \quad -17 \end{array}$$

$$\frac{x^3 - 2x^2 + 3x + 5}{x + 2} = x^2 - 4x + 11 + \frac{-17}{x + 2}$$

$$a + b + c + d = 1 - 4 + 11 - 17 = -9$$

$$\begin{array}{r} 30. \quad h \overline{) 1 \quad 3-h \quad -2h \quad -2h^2 \quad h^3} \\ \underline{h \quad 3h \quad h^2 \quad -h^3} \\ 1 \quad 3 \quad h \quad -h^2 \quad 0 \end{array}$$

$x^3 + 3x^2 + hx - h^2$ is the quotient and 0 is the remainder.

$$\begin{array}{r} 31. \quad -y \overline{) 1 \quad 3y \quad -3y^2 \quad -y^3 \quad 4y^4} \\ \underline{-y \quad -2y^2 \quad 5y^3 \quad 4y^4} \\ 1 \quad 2y \quad -5y^2 \quad 4y^3 \quad 0 \end{array}$$

Yes, $x+y$ is a factor of

$$x^4 + 3x^3y - 3x^2y^2 - xy^3 + 4y^4.$$

32. Answers will vary.

Section R.7

1. lowest terms

2. Least Common Multiple

3. d

4. a

$$\begin{aligned} 5. \text{ True; } \frac{\frac{1}{3} + \frac{1}{5}}{\frac{1}{5} - \frac{1}{x}} &= \frac{\frac{x+3}{3x}}{\frac{x-5}{5x}} \\ &= \frac{x+3}{3x} \cdot \frac{5x}{x-5} = \frac{5(x+3)}{3(x-5)} \end{aligned}$$

6. False;

$$2x^3 + 6x^2 = 2x^2(x+3)$$

$$6x^4 + 4x^3 = 2x^3(3x+2)$$

$$LCM = 2x^3(x+3)(3x+2)$$

$$7. \quad \frac{3x+9}{x^2-9} = \frac{3(x+3)}{(x-3)(x+3)} = \frac{3}{x-3}$$

$$8. \quad \frac{4x^2+8x}{12x+24} = \frac{4x(x+2)}{12(x+2)} = \frac{x}{3}$$

$$9. \quad \frac{x^2-2x}{3x-6} = \frac{x(x-2)}{3(x-2)} = \frac{x}{3}$$

$$10. \quad \frac{15x^2+24x}{3x^2} = \frac{3x(5x+8)}{3x^2} = \frac{5x+8}{x}$$

$$11. \quad \frac{24x^2}{12x^2-6x} = \frac{24x^2}{6x(2x-1)} = \frac{4x}{2x-1}$$

$$12. \quad \frac{x^2+4x+4}{x^2-4} = \frac{(x+2)(x+2)}{(x-2)(x+2)} = \frac{x+2}{x-2}$$

Section R.7: Rational Expressions

$$\begin{aligned} 13. \quad \frac{y^2 - 49}{3y^2 - 18y - 21} &= \frac{(y+7)(y-7)}{3(y^2 - 6y - 7)} \\ &= \frac{(y+7)(y-7)}{3(y-7)(y+1)} \\ &= \frac{y+7}{3(y+1)} \end{aligned}$$

$$14. \quad \frac{3y^2 - y - 2}{3y^2 + 5y + 2} = \frac{(3y+2)(y-1)}{(3y+2)(y+1)} = \frac{y-1}{y+1}$$

$$15. \quad \frac{x^2 + 4x - 12}{x^2 - 4x + 4} = \frac{(x+6)(x-2)}{(x-2)(x-2)} = \frac{x+6}{x-2}$$

$$16. \quad \frac{x - x^2}{x^2 + x - 2} = \frac{-x(x-1)}{(x+2)(x-1)} = \frac{-x}{x+2} = -\frac{x}{x+2}$$

$$\begin{aligned} 17. \quad \frac{x^2 + x - 20}{4 - x} &= \frac{(x+5)(x-4)}{2 - x} \\ &= \frac{(x+5)(x-4)}{-1(x-4)} \\ &= -(x+5) \end{aligned}$$

$$18. \quad \frac{2x^2 + 5x - 3}{1 - 2x} = \frac{(2x-1)(x+3)}{-1(2x-1)} = -(x+3) = -x-3$$

$$\begin{aligned} 19. \quad \frac{3x+6}{5x^2} \cdot \frac{x}{x^2-4} &= \frac{3(x+2)}{5x^2} \cdot \frac{x}{(x-2)(x+2)} \\ &= \frac{3}{5x(x-2)} \end{aligned}$$

$$20. \quad \frac{3}{2x} \cdot \frac{x^2}{6x+10} = \frac{3}{2} \cdot \frac{x}{2(3x+5)} = \frac{3x}{4(3x+5)}$$

$$\begin{aligned} 21. \quad \frac{4x^2}{x^2-16} \cdot \frac{x^3-64}{2x} &= \frac{4x^2}{(x-4)(x+4)} \cdot \frac{(x-4)(x^2+4x+16)}{2x} \\ &= \frac{2x \cdot 2x(x-4)(x^2+4x+16)}{2x(x-4)(x+4)} \\ &= \frac{2x(x^2+4x+16)}{x+4} \end{aligned}$$

$$\begin{aligned} 22. \quad \frac{12}{x^2+x} \cdot \frac{x^3+1}{4x-2} &= \frac{12}{x(x+1)} \cdot \frac{(x+1)(x^2-x+1)}{2(2x-1)} \\ &= \frac{2 \cdot 6(x+1)(x^2-x+1)}{2x(x+1)(2x-1)} \\ &= \frac{6(x^2-x+1)}{x(2x-1)} \end{aligned}$$

$$\begin{aligned} 23. \quad \frac{4x-8}{-3x} \cdot \frac{12}{12-6x} &= \frac{4(x-2)}{-3x} \cdot \frac{12}{6(2-x)} \\ &= \frac{4(x-2)}{-3x} \cdot \frac{2}{(-1)(x-2)} \\ &= \frac{8}{3x} \end{aligned}$$

$$24. \quad \frac{6x-27}{5x} \cdot \frac{2}{4x-18} = \frac{3(2x-9)}{5x} \cdot \frac{2}{2(2x-9)} = \frac{3}{5x}$$

$$\begin{aligned} 25. \quad \frac{x^2-4x-12}{x^2+2x-48} \cdot \frac{x^2+4x-32}{x^2+10x+16} &= \frac{(x-6)(x+2)}{(x+8)(x-6)} \cdot \frac{(x+8)(x-4)}{(x+8)(x+2)} \\ &= \frac{x-4}{x+8} \end{aligned}$$

$$\begin{aligned} 26. \quad \frac{x^2+x-6}{x^2+4x-5} \cdot \frac{x^2-25}{x^2+2x-15} &= \frac{(x-2)(x+3)}{(x+5)(x-1)} \cdot \frac{(x+5)(x-5)}{(x+5)(x-3)} \\ &= \frac{(x-2)(x+3)(x-5)}{(x+5)(x-1)(x-3)} \end{aligned}$$

$$\begin{aligned} 27. \quad \frac{6x}{\frac{x^2-4}{3x-9}} &= \frac{6x}{x^2-4} \cdot \frac{2x+4}{3x-9} \\ &= \frac{6x}{(x-2)(x+2)} \cdot \frac{2(x+2)}{3(x-3)} \\ &= \frac{4x}{(x-2)(x-3)} \end{aligned}$$

Chapter R: Review

$$\begin{aligned} 28. \frac{\frac{12x}{5x+20}}{\frac{4x^2}{x^2-16}} &= \frac{12x}{5x+20} \cdot \frac{x^2-16}{4x^2} \\ &= \frac{12x}{5(x+4)} \cdot \frac{(x+4)(x-4)}{4x^2} \\ &= \frac{3(x-4)}{5x} \end{aligned}$$

$$\begin{aligned} 29. \frac{\frac{8x}{x^2-1}}{\frac{10x}{x+1}} &= \frac{8x}{x^2-1} \cdot \frac{x+1}{10x} \\ &= \frac{8x}{(x-1)(x+1)} \cdot \frac{x+1}{10x} \\ &= \frac{4}{5(x-1)} \end{aligned}$$

$$\begin{aligned} 30. \frac{\frac{x-2}{4x}}{\frac{x^2-4x+4}{12x}} &= \frac{x-2}{4x} \cdot \frac{12x}{x^2-4x+4} \\ &= \frac{x-2}{4x} \cdot \frac{12x}{(x-2)(x-2)} \\ &= \frac{3}{x-2} \end{aligned}$$

$$\begin{aligned} 31. \frac{\frac{4-x}{4+x}}{\frac{4x}{x^2-16}} &= \frac{4-x}{4+x} \cdot \frac{x^2-16}{4x} \\ &= \frac{4-x}{4+x} \cdot \frac{(x+4)(x-4)}{4x} \\ &= \frac{(4-x)(x-4)}{4x} \\ &= -\frac{(x-4)^2}{4x} \end{aligned}$$

$$\begin{aligned} 32. \frac{\frac{3+x}{3-x}}{\frac{x^2-9}{9x^3}} &= \frac{3+x}{3-x} \cdot \frac{9x^3}{x^2-9} \\ &= \frac{3+x}{3-x} \cdot \frac{9x^3}{(x+3)(x-3)} \\ &= \frac{9x^3}{(3-x)(x-3)} \\ &= \frac{9x^3}{-(x-3)^2} \\ &= -\frac{9x^3}{(x-3)^2} \end{aligned}$$

$$\begin{aligned} 33. \frac{\frac{x^2+7x+12}{x^2-7x+12}}{\frac{x^2+x-12}{x^2-x-12}} &= \frac{x^2+7x+12}{x^2-7x+12} \cdot \frac{x^2-x-12}{x^2+x-12} \\ &= \frac{(x+3)(x+4)}{(x-3)(x-4)} \cdot \frac{(x-4)(x+3)}{(x+4)(x-3)} \\ &= \frac{(x+3)^2}{(x-3)^2} \end{aligned}$$

$$\begin{aligned} 34. \frac{\frac{x^2+7x+6}{x^2+x-6}}{\frac{x^2+5x-6}{x^2+5x+6}} &= \frac{x^2+7x+6}{x^2+x-6} \cdot \frac{x^2+5x+6}{x^2+5x-6} \\ &= \frac{(x+6)(x+1)}{(x+3)(x-2)} \cdot \frac{(x+2)(x+3)}{(x+6)(x-1)} \\ &= \frac{(x+1)(x+2)}{(x-2)(x-1)} \end{aligned}$$

$$\begin{aligned} 35. \frac{\frac{5x^2-7x-6}{2x^2+3x-5}}{\frac{15x^2+14x+3}{2x^2+13x+20}} &= \frac{5x^2-7x-6}{2x^2+3x-5} \cdot \frac{2x^2+13x+20}{15x^2+14x+3} \\ &= \frac{(5x+3)(x-2)}{(x-1)(2x+5)} \cdot \frac{(2x+5)(x+4)}{(5x+3)(3x+1)} \\ &= \frac{(x-2)(x+4)}{(x-1)(3x+1)} \end{aligned}$$

Section R.7: Rational Expressions

$$\begin{aligned} 36. \frac{\frac{9x^2+3x-2}{12x^2+5x-2}}{\frac{9x^2-6x+1}{8x^2-10x-3}} &= \frac{9x^2+3x-2}{12x^2+5x-2} \cdot \frac{8x^2-10x-3}{9x^2-6x+1} \\ &= \frac{(3x+2)(3x-1)}{(3x+2)(4x-1)} \cdot \frac{(4x+1)(2x-3)}{(3x-1)(3x-1)} \\ &= \frac{(4x+1)(2x-3)}{(4x-1)(3x-1)} \end{aligned}$$

$$37. \frac{x}{2} + \frac{5}{2} = \frac{x+5}{2}$$

$$38. \frac{3}{x} - \frac{6}{x} = \frac{3-6}{x} = \frac{-3}{x} = -\frac{3}{x}$$

$$39. \frac{x^2}{2x-3} - \frac{4}{2x-3} = \frac{x^2-4}{2x-3} = \frac{(x+2)(x-2)}{2x-3}$$

$$40. \frac{3x^2}{2x-1} - \frac{9}{2x-1} = \frac{3x^2-9}{2x-1} = \frac{3(x^2-3)}{2x-1}$$

$$41. \frac{x+5}{x-4} + \frac{3x-2}{x-4} = \frac{x+5+3x-2}{x-4} = \frac{4x+3}{x-4}$$

$$42. \frac{2x-5}{3x+2} + \frac{x+4}{3x+2} = \frac{2x-5+x+4}{3x+2} = \frac{3x-1}{3x+2}$$

$$\begin{aligned} 43. \frac{3x+5}{2x-1} - \frac{2x-4}{2x-1} &= \frac{(3x+5)-(2x-4)}{2x-1} \\ &= \frac{3x+5-2x+4}{2x-1} \\ &= \frac{x+9}{2x-1} \end{aligned}$$

$$\begin{aligned} 44. \frac{5x-4}{3x+4} - \frac{x+1}{3x+4} &= \frac{(5x-4)-(x+1)}{3x+4} \\ &= \frac{5x-4-x-1}{3x+4} \\ &= \frac{4x-5}{3x+4} \end{aligned}$$

$$45. \frac{4}{x-2} + \frac{x}{2-x} = \frac{4}{x-2} - \frac{x}{x-2} = \frac{4-x}{x-2}$$

$$46. \frac{6}{x-1} - \frac{x}{1-x} = \frac{6}{x-1} + \frac{x}{x-1} = \frac{x+6}{x-1}$$

$$\begin{aligned} 47. \frac{7}{x-3} - \frac{3}{x+1} &= \frac{7(x+1)}{(x-3)(x+1)} - \frac{3(x-3)}{(x+1)(x-3)} \\ &= \frac{7x+7-3x+9}{(x+1)(x-3)} \\ &= \frac{4x+16}{(x+1)(x-3)} \\ &= \frac{4(x+4)}{(x+1)(x-3)} \end{aligned}$$

$$\begin{aligned} 48. \frac{2}{x+5} - \frac{5}{x-5} &= \frac{2(x-5)}{(x+5)(x-5)} - \frac{5(x+5)}{(x+5)(x-5)} \\ &= \frac{2x-10-5x-25}{(x+5)(x-5)} \\ &= \frac{-3x-35}{(x+5)(x-5)} \\ &= -\frac{3x+35}{(x+5)(x-5)} \end{aligned}$$

$$\begin{aligned} 49. \frac{x}{x+1} + \frac{2x-3}{x-1} &= \frac{x(x-1)}{(x+1)(x-1)} + \frac{(2x-3)(x+1)}{(x-1)(x+1)} \\ &= \frac{x^2-x+2x^2-x-3}{(x-1)(x+1)} \\ &= \frac{3x^2-2x-3}{(x-1)(x+1)} \end{aligned}$$

$$\begin{aligned} 50. \frac{3x}{x-4} + \frac{2x}{x+3} &= \frac{3x(x+3)}{(x-4)(x+3)} + \frac{2x(x-4)}{(x-4)(x+3)} \\ &= \frac{3x^2+9x+2x^2-8x}{(x-4)(x+3)} \\ &= \frac{5x^2+x}{(x-4)(x+3)} \\ &= \frac{x(5x+1)}{(x-4)(x+3)} \end{aligned}$$

$$\begin{aligned} 51. \frac{x-3}{x+2} - \frac{x+4}{x-2} &= \frac{(x-3)(x-2)}{(x+2)(x-2)} - \frac{(x+4)(x+2)}{(x-2)(x+2)} \\ &= \frac{x^2-5x+6-(x^2+6x+8)}{(x+2)(x-2)} \\ &= \frac{x^2-5x+6-x^2-6x-8}{(x+2)(x-2)} \\ &= \frac{-11x-2}{(x+2)(x-2)} \quad \text{or} \quad \frac{-(11x+2)}{(x+2)(x-2)} \end{aligned}$$

Chapter R: Review

$$\begin{aligned}
 52. \quad \frac{2x-3}{x-1} - \frac{2x+1}{x+1} &= \frac{(2x-3)(x+1)}{(x-1)(x+1)} - \frac{(2x+1)(x-1)}{(x+1)(x-1)} \\
 &= \frac{2x^2 - x - 3 - (2x^2 - x - 1)}{(x+1)(x-1)} \\
 &= \frac{2x^2 - x - 3 - 2x^2 + x + 1}{(x+1)(x-1)} \\
 &= \frac{-2}{(x+1)(x-1)} \\
 &= -\frac{2}{(x+1)(x-1)}
 \end{aligned}$$

$$\begin{aligned}
 53. \quad \frac{x}{x^2-4} + \frac{1}{x} &= \frac{x^2 + x^2 - 4}{x(x^2 - 4)} \\
 &= \frac{2x^2 - 4}{x(x^2 - 4)} \\
 &= \frac{2(x^2 - 2)}{x(x-2)(x+2)}
 \end{aligned}$$

$$\begin{aligned}
 54. \quad \frac{x-1}{x^3} + \frac{x}{x^2+1} &= \frac{(x-1)(x^2+1) + x^4}{x^3(x^2+1)} \\
 &= \frac{x^3 - x^2 + x - 1 + x^4}{x^3(x^2+1)} \\
 &= \frac{x^4 + x^3 - x^2 + x - 1}{x^3(x^2+1)}
 \end{aligned}$$

$$\begin{aligned}
 55. \quad x^2 - 4 &= (x+2)(x-2) \\
 x^2 - x - 2 &= (x+1)(x-2) \\
 \text{Therefore, LCM} &= (x+2)(x-2)(x+1).
 \end{aligned}$$

$$\begin{aligned}
 56. \quad x^2 - x - 12 &= (x+3)(x-4) \\
 x^2 - 8x + 16 &= (x-4)(x-4) \\
 \text{Therefore, LCM} &= (x+3)(x-4)^2.
 \end{aligned}$$

$$\begin{aligned}
 57. \quad x^3 - x &= x(x^2 - 1) = x(x+1)(x-1) \\
 x^2 - x &= x(x-1) \\
 \text{Therefore, LCM} &= x(x+1)(x-1).
 \end{aligned}$$

$$\begin{aligned}
 58. \quad 3x^2 - 27 &= 3(x^2 - 9) = 3(x+3)(x-3) \\
 2x^2 - x - 15 &= (2x+5)(x-3) \\
 \text{Therefore, LCM} &= 3(2x+5)(x-3)(x+3).
 \end{aligned}$$

$$\begin{aligned}
 59. \quad 4x^3 - 4x^2 + x &= x(4x^2 - 4x + 1) \\
 &= x(2x-1)(2x-1) \\
 2x^3 - x^2 &= x^2(2x-1) \\
 x^3 & \\
 \text{Therefore, LCM} &= x^3(2x-1)^2.
 \end{aligned}$$

$$\begin{aligned}
 60. \quad x-3 & \\
 x^2 + 3x &= x(x+3) \\
 x^3 - 9x &= x(x^2 - 9) = x(x+3)(x-3) \\
 \text{Therefore, LCM} &= x(x+3)(x-3).
 \end{aligned}$$

$$\begin{aligned}
 61. \quad x^3 - x &= x(x^2 - 1) = x(x+1)(x-1) \\
 x^3 - 2x^2 + x &= x(x^2 - 2x + 1) = x(x-1)^2 \\
 x^3 - 1 &= (x-1)(x^2 + x + 1) \\
 \text{Therefore, LCM} &= x(x+1)(x-1)^2(x^2 + x + 1).
 \end{aligned}$$

$$\begin{aligned}
 62. \quad x^2 + 4x + 4 &= (x+2)^2 \\
 x^3 + 2x^2 &= x^2(x+2) \\
 (x+2)^3 & \\
 \text{Therefore, LCM} &= x^2(x+2)^3.
 \end{aligned}$$

$$\begin{aligned}
 63. \quad \frac{x}{x^2-7x+6} - \frac{x}{x^2-2x-24} & \\
 &= \frac{x}{(x-6)(x-1)} - \frac{x}{(x-6)(x+4)} \\
 &= \frac{x(x+4)}{(x-6)(x-1)(x+4)} - \frac{x(x-1)}{(x-6)(x+4)(x-1)} \\
 &= \frac{x^2 + 4x - x^2 + x}{(x-6)(x+4)(x-1)} = \frac{5x}{(x-6)(x+4)(x-1)}
 \end{aligned}$$

$$\begin{aligned}
 64. \quad \frac{x}{x-3} - \frac{x+1}{x^2+5x-24} & \\
 &= \frac{x}{(x-3)} - \frac{x+1}{(x-3)(x+8)} \\
 &= \frac{x(x+8)}{(x-3)(x+8)} - \frac{x+1}{(x-3)(x+8)} \\
 &= \frac{x^2 + 8x - x - 1}{(x-3)(x+8)} = \frac{x^2 + 7x - 1}{(x-3)(x+8)}
 \end{aligned}$$

Section R.7: Rational Expressions

$$\begin{aligned}
 65. \quad & \frac{4x}{x^2-4} - \frac{2}{x^2+x-6} \\
 &= \frac{4x}{(x-2)(x+2)} - \frac{2}{(x+3)(x-2)} \\
 &= \frac{4x(x+3)}{(x-2)(x+2)(x+3)} - \frac{2(x+2)}{(x+3)(x-2)(x+2)} \\
 &= \frac{4x^2+12x-2x-4}{(x-2)(x+2)(x+3)} \\
 &= \frac{4x^2+10x-4}{(x-2)(x+2)(x+3)} \\
 &= \frac{2(2x^2+5x-2)}{(x-2)(x+2)(x+3)}
 \end{aligned}$$

$$\begin{aligned}
 66. \quad & \frac{3x}{x-1} - \frac{x-4}{x^2-2x+1} = \frac{3x}{(x-1)} - \frac{x-4}{(x-1)^2} \\
 &= \frac{3x(x-1)}{(x-1)(x-1)} - \frac{x-4}{(x-1)^2} \\
 &= \frac{3x^2-3x-x+4}{(x-1)^2} \\
 &= \frac{3x^2-4x+4}{(x-1)^2}
 \end{aligned}$$

$$\begin{aligned}
 67. \quad & \frac{3}{(x-1)^2(x+1)} + \frac{2}{(x-1)(x+1)^2} \\
 &= \frac{3(x+1)+2(x-1)}{(x-1)^2(x+1)^2} \\
 &= \frac{3x+3+2x-2}{(x-1)^2(x+1)^2} \\
 &= \frac{5x+1}{(x-1)^2(x+1)^2}
 \end{aligned}$$

$$\begin{aligned}
 68. \quad & \frac{2}{(x+2)^2(x-1)} - \frac{6}{(x+2)(x-1)^2} \\
 &= \frac{2(x-1)-6(x+2)}{(x+2)^2(x-1)^2} \\
 &= \frac{2x-2-6x-12}{(x+2)^2(x-1)^2} \\
 &= \frac{-4x-14}{(x+2)^2(x-1)^2} \\
 &= \frac{-2(2x+7)}{(x+2)^2(x-1)^2}
 \end{aligned}$$

$$\begin{aligned}
 69. \quad & \frac{x+4}{x^2-x-2} - \frac{2x+3}{x^2+2x-8} \\
 &= \frac{x+4}{(x-2)(x+1)} - \frac{2x+3}{(x+4)(x-2)} \\
 &= \frac{(x+4)(x+4)}{(x-2)(x+1)(x+4)} - \frac{(2x+3)(x+1)}{(x+4)(x-2)(x+1)} \\
 &= \frac{x^2+8x+16-(2x^2+5x+3)}{(x-2)(x+1)(x+4)} \\
 &= \frac{-x^2+3x+13}{(x-2)(x+1)(x+4)}
 \end{aligned}$$

$$\begin{aligned}
 70. \quad & \frac{2x-3}{x^2+8x+7} - \frac{x-2}{(x+1)^2} \\
 &= \frac{2x-3}{(x+1)(x+7)} - \frac{x-2}{(x+1)^2} \\
 &= \frac{(2x-3)(x+1)}{(x+1)(x+7)(x+1)} - \frac{(x-2)(x+7)}{(x+1)^2(x+7)} \\
 &= \frac{2x^2-x-3-(x^2+5x-14)}{(x+1)^2(x+7)} \\
 &= \frac{x^2-6x+11}{(x+1)^2(x+7)}
 \end{aligned}$$

$$\begin{aligned}
 71. \quad & \frac{1}{x} - \frac{2}{x^2+x} + \frac{3}{x^3-x^2} \\
 &= \frac{1}{x} - \frac{2}{x(x+1)} + \frac{3}{x^2(x-1)} \\
 &= \frac{x(x+1)(x-1)-2x(x-1)+3(x+1)}{x^2(x+1)(x-1)} \\
 &= \frac{x(x^2-1)-2x^2+2x+3x+3}{x^2(x+1)(x-1)} \\
 &= \frac{x^3-x-2x^2+5x+3}{x^2(x+1)(x-1)} \\
 &= \frac{x^3-2x^2+4x+3}{x^2(x+1)(x-1)}
 \end{aligned}$$

Chapter R: Review

$$\begin{aligned}
 72. \quad & \frac{x}{(x-1)^2} + \frac{2}{x} - \frac{x+1}{x^3-x^2} \\
 &= \frac{x}{(x-1)^2} + \frac{2}{x} - \frac{x+1}{x^2(x-1)} \\
 &= \frac{x^3+2x(x-1)^2-(x+1)(x-1)}{x^2(x-1)^2} \\
 &= \frac{x^3+2x(x^2-2x+1)-(x^2-1)}{x^2(x-1)^2} \\
 &= \frac{x^3+2x^3-4x^2+2x-x^2+1}{x^2(x-1)^2} \\
 &= \frac{3x^3-5x^2+2x+1}{x^2(x-1)^2}
 \end{aligned}$$

$$\begin{aligned}
 73. \quad & \frac{1}{h} \left(\frac{1}{x+h} - \frac{1}{x} \right) = \frac{1}{h} \left(\frac{1 \cdot x}{(x+h)x} - \frac{1(x+h)}{x(x+h)} \right) \\
 &= \frac{1}{h} \left(\frac{x-x-h}{x(x+h)} \right) \\
 &= \frac{-h}{hx(x+h)} \\
 &= \frac{-1}{x(x+h)}
 \end{aligned}$$

$$\begin{aligned}
 74. \quad & \frac{1}{h} \left(\frac{1}{(x+h)^2} - \frac{1}{x^2} \right) \\
 &= \frac{1}{h} \left(\frac{1 \cdot x^2}{(x+h)^2 x^2} - \frac{1(x+h)^2}{x^2(x+h)^2} \right) \\
 &= \frac{1}{h} \left(\frac{x^2 - (x^2 + 2xh + h^2)}{x^2(x+h)^2} \right) \\
 &= \frac{-2xh - h^2}{hx^2(x+h)^2} \\
 &= \frac{h(-2x-h)}{hx^2(x+h)^2} \\
 &= \frac{-2x-h}{x^2(x+h)^2} \\
 &= -\frac{2x+h}{x^2(x+h)^2}
 \end{aligned}$$

$$\begin{aligned}
 75. \quad & \frac{1+\frac{1}{x}}{1-\frac{1}{x}} = \frac{\left(\frac{x}{x} + \frac{1}{x}\right)}{\left(\frac{x}{x} - \frac{1}{x}\right)} = \frac{\left(\frac{x+1}{x}\right)}{\left(\frac{x-1}{x}\right)} = \frac{x+1}{x} \cdot \frac{x}{x-1} = \frac{x+1}{x-1}
 \end{aligned}$$

$$\begin{aligned}
 76. \quad & \frac{4+\frac{1}{x^2}}{3-\frac{1}{x^2}} = \frac{\left(\frac{4x^2}{x^2} + \frac{1}{x^2}\right)}{\left(\frac{3x^2}{x^2} - \frac{1}{x^2}\right)} = \frac{\left(\frac{4x^2+1}{x^2}\right)}{\left(\frac{3x^2-1}{x^2}\right)} \\
 &= \frac{4x^2+1}{x^2} \cdot \frac{x^2}{3x^2-1} \\
 &= \frac{4x^2+1}{3x^2-1}
 \end{aligned}$$

$$\begin{aligned}
 77. \quad & \frac{2-\frac{x+1}{x}}{3+\frac{x-1}{x+1}} = \frac{\frac{2x}{x} - \frac{x+1}{x}}{\frac{3(x+1)}{x+1} + \frac{x-1}{x+1}} = \frac{\frac{2x-x-1}{x}}{\frac{3x+3+x-1}{x+1}} \\
 &= \frac{\frac{x-1}{x}}{\frac{4x+2}{x+1}} = \frac{x-1}{x} \cdot \frac{x+1}{2(2x+1)} \\
 &= \frac{(x-1)(x+1)}{2x(2x+1)}
 \end{aligned}$$

$$\begin{aligned}
 78. \quad & \frac{1-\frac{x}{x+1}}{2-\frac{x-1}{x}} = \frac{\left(\frac{x+1}{x+1} - \frac{x}{x+1}\right)}{\left(\frac{2x}{x} - \frac{x-1}{x}\right)} = \frac{\left(\frac{1}{x+1}\right)}{\left(\frac{x+1}{x}\right)} \\
 &= \frac{1}{x+1} \cdot \frac{x}{x+1} \\
 &= \frac{x}{(x+1)^2}
 \end{aligned}$$

$$\begin{aligned}
 79. \quad & \frac{\frac{x+4}{x-2} - \frac{x-3}{x+1}}{x+1} \\
 &= \frac{\left(\frac{(x+4)(x+1)}{(x-2)(x+1)} - \frac{(x-3)(x-2)}{(x+1)(x-2)}\right)}{x+1} \\
 &= \frac{\left(\frac{x^2+5x+4-(x^2-5x+6)}{(x-2)(x+1)}\right)}{x+1} \\
 &= \frac{10x-2}{(x-2)(x+1)} \cdot \frac{1}{x+1} \\
 &= \frac{2(5x-1)}{(x-2)(x+1)^2}
 \end{aligned}$$

Section R.7: Rational Expressions

$$\begin{aligned}
 80. \quad & \frac{\frac{x-2}{x+1} - \frac{x}{x-2}}{x+3} \\
 &= \frac{\left(\frac{(x-2)(x-2)}{(x+1)(x-2)} - \frac{x(x+1)}{(x-2)(x+1)} \right)}{x+3} \\
 &= \frac{\left(\frac{x^2 - 4x + 4 - (x^2 + x)}{(x-2)(x+1)} \right)}{x+3} \\
 &= \frac{-5x + 4}{(x-2)(x+1)} \cdot \frac{1}{x+3} \\
 &= \frac{-5x + 4}{(x-2)(x+1)(x+3)} \\
 &= \frac{-(5x-4)}{(x-2)(x+1)(x+3)}
 \end{aligned}$$

$$\begin{aligned}
 81. \quad & \frac{\frac{x-2}{x+2} + \frac{x-1}{x+1}}{\frac{x}{x+1} - \frac{2x-3}{x}} \\
 &= \frac{\left(\frac{(x-2)(x+1)}{(x+2)(x+1)} + \frac{(x-1)(x+2)}{(x+1)(x+2)} \right)}{\left(\frac{x^2}{(x+1)(x)} - \frac{(2x-3)(x+1)}{x(x+1)} \right)} \\
 &= \frac{\left(\frac{x^2 - x - 2 + x^2 + x - 2}{(x+2)(x+1)} \right)}{\left(\frac{x^2 - (2x^2 - x - 3)}{x(x+1)} \right)} \\
 &= \frac{\left(\frac{2x^2 - 4}{(x+2)(x+1)} \right)}{\left(\frac{-x^2 + x + 3}{x(x+1)} \right)} \\
 &= \frac{2(x^2 - 2)}{(x+2)(x+1)} \cdot \frac{x(x+1)}{-(x^2 - x - 3)} \\
 &= \frac{2x(x^2 - 2)}{-(x+2)(x^2 - x - 3)} \\
 &= \frac{-2x(x^2 - 2)}{(x+2)(x^2 - x - 3)}
 \end{aligned}$$

$$\begin{aligned}
 82. \quad & \frac{\frac{2x+5}{x} - \frac{x}{x-3}}{\frac{x^2}{x-3} - \frac{(x+1)^2}{x+3}} \\
 &= \frac{\left(\frac{(2x+5)(x-3)}{x(x-3)} - \frac{x(x)}{x(x-3)} \right)}{\left(\frac{x^2(x+3)}{(x-3)(x+3)} - \frac{(x-3)(x+1)^2}{(x-3)(x+3)} \right)} \\
 &= \frac{\left(\frac{2x^2 - x - 15 - x^2}{x(x-3)} \right)}{\left(\frac{x^3 + 3x^2 - (x^3 - x^2 - 5x - 3)}{(x-3)(x+3)} \right)} \\
 &= \frac{\left(\frac{x^2 - x - 15}{x(x-3)} \right)}{\left(\frac{4x^2 + 5x + 3}{(x-3)(x+3)} \right)} \\
 &= \frac{x^2 - x - 15}{x(x-3)} \cdot \frac{(x-3)(x+3)}{4x^2 + 5x + 3} \\
 &= \frac{(x^2 - x - 15)(x+3)}{x(4x^2 + 5x + 3)}
 \end{aligned}$$

$$\begin{aligned}
 83. \quad & 1 - \frac{1}{1 - \frac{1}{x}} = 1 - \frac{1}{\frac{x-1}{x}} \\
 &= 1 - \frac{x}{x-1} \\
 &= \frac{x-1-x}{x-1} \\
 &= \frac{-1}{x-1}
 \end{aligned}$$

$$\begin{aligned}
 84. \quad & 1 - \frac{1}{1 - \frac{1}{1-x}} = 1 - \frac{1}{\frac{1-x-1}{1-x}} = 1 - \frac{1}{\frac{-x}{1-x}} \\
 &= 1 - \frac{1-x}{-x} = 1 + \frac{1-x}{x} \\
 &= \frac{x+1-x}{x} \\
 &= \frac{1}{x}
 \end{aligned}$$

Chapter R: Review

$$\begin{aligned}
 85. \quad \frac{2(x-1)^{-1} + 3}{3(x-1)^{-1} + 2} &= \frac{\frac{2}{x-1} + 3}{\frac{3}{x-1} + 2} = \frac{\frac{2}{x-1} + \frac{3(x-1)}{x-1}}{\frac{3}{x-1} + \frac{2(x-1)}{x-1}} \\
 &= \frac{\frac{2+3(x-1)}{x-1}}{\frac{3+2(x-1)}{x-1}} \\
 &= \frac{2+3(x-1)}{3+2(x-1)} = \frac{2+3x-3}{3+2x-2} \\
 &= \frac{3x-1}{2x+1}
 \end{aligned}$$

$$\begin{aligned}
 86. \quad \frac{4(x+2)^{-1} - 3}{3(x+2)^{-1} - 1} &= \frac{\frac{4}{x+2} - 3}{\frac{3}{x+2} - 1} = \frac{\frac{4}{x+2} - \frac{3(x+2)}{x+2}}{\frac{3}{x+2} - \frac{1(x+2)}{x+2}} \\
 &= \frac{\frac{4-3(x+2)}{x+2}}{\frac{3-(x+2)}{x+2}} \\
 &= \frac{4-3(x+2)}{3-(x+2)} = \frac{4-3x-6}{3-x-2} \\
 &= \frac{-3x-2}{-x+1} = \frac{3x+2}{x-1}
 \end{aligned}$$

$$\begin{aligned}
 87. \quad \frac{(2x+3) \cdot 3 - (3x-5) \cdot 2}{(3x-5)^2} &= \frac{6x+9-6x+10}{(3x-5)^2} \\
 &= \frac{19}{(3x-5)^2}
 \end{aligned}$$

$$\begin{aligned}
 88. \quad \frac{(4x+1) \cdot 5 - (5x-2) \cdot 4}{(5x-2)^2} &= \frac{20x+5-20x+8}{(5x-2)^2} \\
 &= \frac{13}{(5x-2)^2}
 \end{aligned}$$

$$\begin{aligned}
 89. \quad \frac{x \cdot 2x - (x^2+1) \cdot 1}{(x^2+1)^2} &= \frac{2x^2 - x^2 - 1}{(x^2+1)^2} \\
 &= \frac{x^2 - 1}{(x^2+1)^2} \\
 &= \frac{(x-1)(x+1)}{(x^2+1)^2}
 \end{aligned}$$

$$\begin{aligned}
 90. \quad \frac{x \cdot 2x - (x^2-4) \cdot 1}{(x^2-4)^2} &= \frac{2x^2 - x^2 + 4}{(x^2-4)^2} = \frac{x^2 + 4}{(x^2-4)^2} \\
 &= \frac{x^2 + 4}{(x+2)^2(x-2)^2}
 \end{aligned}$$

$$\begin{aligned}
 91. \quad \frac{(3x+1) \cdot 2x - x^2 \cdot 3}{(3x+1)^2} &= \frac{6x^2 + 2x - 3x^2}{(3x+1)^2} \\
 &= \frac{3x^2 + 2x}{(3x+1)^2} \\
 &= \frac{x(3x+2)}{(3x+1)^2}
 \end{aligned}$$

$$\begin{aligned}
 92. \quad \frac{(2x-5) \cdot 3x^2 - x^3 \cdot 2}{(2x-5)^2} &= \frac{6x^3 - 15x^2 - 2x^3}{(2x-5)^2} \\
 &= \frac{4x^3 - 15x^2}{(2x-5)^2} \\
 &= \frac{x^2(4x-15)}{(2x-5)^2}
 \end{aligned}$$

$$\begin{aligned}
 93. \quad \frac{(x^2+1) \cdot 3 - (3x+4) \cdot 2x}{(x^2+1)^2} &= \frac{3x^2 + 3 - 6x^2 - 8x}{(x^2+1)^2} \\
 &= \frac{-3x^2 - 8x + 3}{(x^2+1)^2} \\
 &= \frac{-(3x^2 + 8x - 3)}{(x^2+1)^2} \\
 &= -\frac{(3x-1)(x+3)}{(x^2+1)^2}
 \end{aligned}$$

Section R.7: Rational Expressions

$$\begin{aligned}
 94. \quad \frac{(x^2+9) \cdot 2 - (2x-5) \cdot 2x}{(x^2+9)^2} &= \frac{2x^2+18-4x^2+10x}{(x^2+9)^2} \\
 &= \frac{-2x^2+10x+18}{(x^2+9)^2} \\
 &= \frac{-2(x^2-5x-9)}{(x^2+9)^2}
 \end{aligned}$$

$$\begin{aligned}
 95. \quad \frac{1}{f} &= (n-1) \left(\frac{1}{R_1} + \frac{1}{R_2} \right) \\
 \frac{1}{f} &= (n-1) \left(\frac{R_2+R_1}{R_1 \cdot R_2} \right) \\
 \frac{R_1 \cdot R_2}{f} &= (n-1)(R_2+R_1) \\
 \frac{f}{R_1 \cdot R_2} &= \frac{1}{(n-1)(R_2+R_1)} \\
 f &= \frac{R_1 \cdot R_2}{(n-1)(R_2+R_1)} \\
 f &= \frac{0.1(0.2)}{(1.5-1)(0.2+0.1)} \\
 &= \frac{0.02}{0.5(0.3)} = \frac{0.02}{0.15} = \frac{2}{15} \text{ meters}
 \end{aligned}$$

$$\begin{aligned}
 96. \quad \frac{1}{R} &= \frac{1}{R_1} + \frac{1}{R_2} + \frac{1}{R_3} = \frac{R_2 R_3 + R_1 R_3 + R_1 R_2}{R_1 R_2 R_3} \\
 R &= \frac{R_1 R_2 R_3}{R_2 R_3 + R_1 R_3 + R_1 R_2} \\
 &= \frac{5 \cdot 4 \cdot 10}{4 \cdot 10 + 5 \cdot 10 + 5 \cdot 4} \\
 &= \frac{200}{110} = \frac{20}{11} \text{ ohms}
 \end{aligned}$$

$$\begin{aligned}
 97. \quad \frac{x^2+3xk-3x-9k}{x^2+2x-15} &= \frac{x(x+3k)-3(x+3k)}{(x-3)(x+5)} \\
 \frac{(x-3)(x+3k)}{(x-3)(x+5)} &= \frac{(x+3k)}{(x+5)} \\
 &= \frac{x+12}{x+5} \\
 x+3k &= x+12 \\
 3k &= 12 \\
 k &= 4
 \end{aligned}$$

$$\begin{aligned}
 98. \quad \frac{A}{x+2} + \frac{7}{x-3} &= \frac{12x-1}{x^2-x+6} \\
 A(x-3) + 7(x+2) &= 12x-1 \\
 Ax-3A+7x+14 &= 12x-1 \\
 x(A+7)-3A+14 &= 12x-1 \\
 &\text{so} \\
 A+7 &= 12 \\
 A &= 5
 \end{aligned}$$

$$\begin{aligned}
 99. \quad 1 + \frac{1}{x} &= \frac{x+1}{x} \Rightarrow a=1, b=1, c=0 \\
 1 + \frac{1}{1+\frac{1}{x}} &= 1 + \frac{1}{\left(\frac{x+1}{x}\right)} = 1 + \frac{x}{x+1} \\
 &= \frac{x+1+x}{x+1} = \frac{2x+1}{x+1} \\
 \Rightarrow a &= 2, b=1, c=1
 \end{aligned}$$

$$\begin{aligned}
 1 + \frac{1}{1+\frac{1}{1+\frac{1}{x}}} &= 1 + \frac{1}{\left(\frac{2x+1}{x+1}\right)} = 1 + \frac{x+1}{2x+1} \\
 &= \frac{2x+1+x+1}{2x+1} = \frac{3x+2}{2x+1} \\
 \Rightarrow a &= 3, b=2, c=1
 \end{aligned}$$

$$\begin{aligned}
 1 + \frac{1}{1+\frac{1}{1+\frac{1}{1+\frac{1}{x}}}}} &= 1 + \frac{1}{\left(\frac{3x+2}{2x+1}\right)} = 1 + \frac{2x+1}{3x+2} \\
 &= \frac{3x+2+2x+1}{3x+2} = \frac{5x+3}{3x+2} \\
 \Rightarrow a &= 5, b=3, c=2
 \end{aligned}$$

If we continue this process, the values of a , b and c produce the following sequences:

$a: 1, 2, 3, 5, 8, 13, 21, \dots$

$b: 1, 1, 2, 3, 5, 8, 13, 21, \dots$

$c: 0, 1, 1, 2, 3, 5, 8, 13, 21, \dots$

In each case we have a *Fibonacci sequence*, where the next value in the list is obtained from the sum of the previous 2 values in the list.

Chapter R: Review

100. Answers will vary.

101. Answers will vary.

Section R.8

1. 9; -9

2. 4; $|-4| = 4$

3. index

4. cube root

5. b

6. d

7. c

8. c

9. true

10. False; $\sqrt[4]{(-3)^4} = |-3| = 3$

11. $\sqrt[3]{27} = \sqrt[3]{3^3} = 3$

12. $\sqrt[4]{16} = \sqrt[4]{2^4} = 2$

13. $\sqrt[3]{-8} = \sqrt[3]{(-2)^3} = -2$

14. $\sqrt[3]{-1} = \sqrt[3]{(-1)^3} = -1$

15. $\sqrt{8} = \sqrt{4 \cdot 2} = 2\sqrt{2}$

16. $\sqrt{75} = \sqrt{25 \cdot 3} = 5\sqrt{3}$

17. $\sqrt{700} = \sqrt{100 \cdot 7} = 10\sqrt{7}$

18. $\sqrt{45x^3} = \sqrt{9 \cdot 5 \cdot x^2 \cdot x} = 3x\sqrt{5x}$

19. $\sqrt[3]{32} = \sqrt[3]{8 \cdot 4} = 2\sqrt[3]{4}$

20. $\sqrt[3]{54} = \sqrt[3]{27 \cdot 2} = 3\sqrt[3]{2}$

21. $\sqrt[3]{-8x^4} = \sqrt[3]{-8x^3 \cdot x} = -2x\sqrt[3]{x}$

22. $\sqrt[3]{192x^5} = \sqrt[3]{64 \cdot 3x^3 \cdot x^2} = 4x\sqrt[3]{3x^2}$

23. $\sqrt[4]{243} = \sqrt[4]{81 \cdot 3} = 3\sqrt[4]{3}$

24. $\sqrt[4]{48x^5} = \sqrt[4]{16x^4 \cdot 3x} = 2x\sqrt[4]{3x}$

25. $\sqrt[4]{x^{12}y^8} = \sqrt[4]{(x^3)^4(y^2)^4} = x^3y^2$

26. $\sqrt[5]{x^{10}y^5} = \sqrt[5]{(x^2)^5y^5} = x^2y$

27. $\sqrt[4]{\frac{x^9y^7}{xy^3}} = \sqrt[4]{x^8y^4} = x^2y$

28. $\sqrt[3]{\frac{3xy^2}{81x^4y^2}} = \sqrt[3]{\frac{1}{27x^3}} = \frac{\sqrt[3]{1}}{\sqrt[3]{27x^3}} = \frac{1}{3x}$

29. $\sqrt{64x} = 8\sqrt{x}$

30. $\sqrt{9x^5} = 3\sqrt{x^4 \cdot x} = 3x^2\sqrt{x}$

31. $\sqrt[4]{162x^9y^{12}} = \sqrt[4]{2(3)^4x(x^2)^4(y^3)^4}$
 $= 3x^2y^3\sqrt[4]{2x}$

32. $\sqrt[3]{-40x^{14}y^{10}} = \sqrt[3]{5(-2)^3x^2(x^4)^3y(y^3)^3}$
 $= -2x^4y^3\sqrt[3]{5x^2y}$

33. $\sqrt{15x^2}\sqrt{5x} = \sqrt{75x^2 \cdot x} = \sqrt{25 \cdot 3 \cdot x^2 \cdot x} = 5x\sqrt{3x}$

34. $\sqrt{5x}\sqrt{20x^3} = \sqrt{100x^4} = 10x^2$

35. $(\sqrt{5}\sqrt[3]{9})^2 = (\sqrt{5})^2(\sqrt[3]{9})^2$
 $= 5 \cdot \sqrt[3]{9^2} = 5\sqrt[3]{81} = 5 \cdot 3\sqrt[3]{3} = 15\sqrt[3]{3}$

36. $(\sqrt[3]{3}\sqrt{10})^4 = (\sqrt[3]{3})^4(\sqrt{10})^4$
 $= \sqrt[3]{3^4} \cdot 10^2 = 3\sqrt[3]{3} \cdot 100 = 300\sqrt[3]{3}$

37. $(3\sqrt{6})(2\sqrt{2}) = 6\sqrt{12} = 6\sqrt{4 \cdot 3} = 12\sqrt{3}$

Section R.8: *n*th Roots; Rational Exponents

$$38. (5\sqrt{8})(-3\sqrt{3}) = -15\sqrt{24} = -30\sqrt{6}$$

$$39. 3\sqrt{2} + 4\sqrt{2} = (3+4)\sqrt{2} = 7\sqrt{2}$$

$$40. 6\sqrt{5} - 4\sqrt{5} = (6-4)\sqrt{5} = 2\sqrt{5}$$

$$\begin{aligned} 41. -\sqrt{48} + 5\sqrt{12} &= -\sqrt{16 \cdot 3} + 5\sqrt{4 \cdot 3} \\ &= -4\sqrt{3} + 5 \cdot 2\sqrt{3} \\ &= (-4+10)\sqrt{3} \\ &= 6\sqrt{3} \end{aligned}$$

$$\begin{aligned} 42. 2\sqrt{12} - 3\sqrt{27} &= 2\sqrt{4 \cdot 3} - 3\sqrt{9 \cdot 3} \\ &= 4\sqrt{3} - 9\sqrt{3} \\ &= (4-9)\sqrt{3} \\ &= -5\sqrt{3} \end{aligned}$$

$$\begin{aligned} 43. (\sqrt{3}+3)(\sqrt{3}-1) &= (\sqrt{3})^2 + 3\sqrt{3} - \sqrt{3} - 3 \\ &= 3 + 2\sqrt{3} - 3 \\ &= 2\sqrt{3} \end{aligned}$$

$$\begin{aligned} 44. (\sqrt{5}-2)(\sqrt{5}+3) &= (\sqrt{5})^2 - 2\sqrt{5} + 3\sqrt{5} - 6 \\ &= 5 + \sqrt{5} - 6 \\ &= \sqrt{5} - 1 \end{aligned}$$

$$\begin{aligned} 45. 5\sqrt[3]{2} - 2\sqrt[3]{54} &= 5\sqrt[3]{2} - 2 \cdot 3\sqrt[3]{2} \\ &= 5\sqrt[3]{2} - 6\sqrt[3]{2} \\ &= (5-6)\sqrt[3]{2} \\ &= -\sqrt[3]{2} \end{aligned}$$

$$\begin{aligned} 46. 9\sqrt[3]{24} - \sqrt[3]{81} &= 9 \cdot 2\sqrt[3]{3} - 3\sqrt[3]{3} \\ &= 18\sqrt[3]{3} - 3\sqrt[3]{3} \\ &= (18-3)\sqrt[3]{3} \\ &= 15\sqrt[3]{3} \end{aligned}$$

$$\begin{aligned} 47. (\sqrt{x}-1)^2 &= (\sqrt{x})^2 - 2\sqrt{x} + 1 \\ &= x - 2\sqrt{x} + 1 \end{aligned}$$

$$\begin{aligned} 48. (\sqrt{x} + \sqrt{5})^2 &= (\sqrt{x})^2 + 2(\sqrt{x})(\sqrt{5}) + (\sqrt{5})^2 \\ &= x + 2\sqrt{5x} + 5 \end{aligned}$$

$$\begin{aligned} 49. \sqrt[3]{16x^4} - \sqrt[3]{2x} &= \sqrt[3]{8x^3 \cdot 2x} - \sqrt[3]{2x} \\ &= 2x\sqrt[3]{2x} - \sqrt[3]{2x} \\ &= (2x-1)\sqrt[3]{2x} \end{aligned}$$

$$\begin{aligned} 50. \sqrt[4]{32x} + \sqrt[4]{2x^5} &= \sqrt[4]{16 \cdot 2x} + \sqrt[4]{x^4 \cdot 2x} \\ &= 2\sqrt[4]{2x} + x\sqrt[4]{2x} \\ &= (2+x)\sqrt[4]{2x} \text{ or } (x+2)\sqrt[4]{2x} \end{aligned}$$

$$\begin{aligned} 51. \sqrt{8x^3} - 3\sqrt{50x} &= \sqrt{4x^2 \cdot 2x} - 3\sqrt{25 \cdot 2x} \\ &= 2x\sqrt{2x} - 15\sqrt{2x} \\ &= (2x-15)\sqrt{2x} \end{aligned}$$

$$\begin{aligned} 52. 3x\sqrt{9y} + 4\sqrt{25y} &= 9x\sqrt{y} + 20\sqrt{y} \\ &= (9x+20)\sqrt{y} \end{aligned}$$

$$\begin{aligned} 53. \sqrt[3]{16x^4y} - 3x\sqrt[3]{2xy} + 5\sqrt[3]{-2xy^4} \\ &= \sqrt[3]{8x^3 \cdot 2xy} - 3x\sqrt[3]{2xy} + 5\sqrt[3]{-y^3 \cdot 2xy} \\ &= 2x\sqrt[3]{2xy} - 3x\sqrt[3]{2xy} - 5y\sqrt[3]{2xy} \\ &= (2x-3x-5y)\sqrt[3]{2xy} \\ &= (-x-5y)\sqrt[3]{2xy} \text{ or } -(x+5y)\sqrt[3]{2xy} \end{aligned}$$

$$\begin{aligned} 54. 8xy - \sqrt{25x^2y^2} + \sqrt[3]{8x^3y^3} &= 8xy - 5xy + 2xy \\ &= (8-5+2)xy \\ &= 5xy \end{aligned}$$

$$55. \frac{1}{\sqrt{2}} = \frac{1}{\sqrt{2}} \cdot \frac{\sqrt{2}}{\sqrt{2}} = \frac{\sqrt{2}}{2}$$

$$56. \frac{2}{\sqrt{3}} = \frac{2}{\sqrt{3}} \cdot \frac{\sqrt{3}}{\sqrt{3}} = \frac{2\sqrt{3}}{3}$$

$$57. \frac{-\sqrt{3}}{\sqrt{5}} = \frac{-\sqrt{3}}{\sqrt{5}} \cdot \frac{\sqrt{5}}{\sqrt{5}} = \frac{-\sqrt{15}}{5}$$

$$58. \frac{-\sqrt{3}}{\sqrt{8}} = \frac{-\sqrt{3}}{2\sqrt{2}} = \frac{-\sqrt{3}}{2\sqrt{2}} \cdot \frac{\sqrt{2}}{\sqrt{2}} = \frac{-\sqrt{6}}{2 \cdot 2} = \frac{-\sqrt{6}}{4}$$

Chapter R: Review

$$\begin{aligned} 59. \quad \frac{\sqrt{3}}{5-\sqrt{2}} &= \frac{\sqrt{3}}{5-\sqrt{2}} \cdot \frac{5+\sqrt{2}}{5+\sqrt{2}} \\ &= \frac{\sqrt{3}(5+\sqrt{2})}{25-2} \\ &= \frac{\sqrt{3}(5+\sqrt{2})}{23} \quad \text{or} \quad \frac{5\sqrt{3}+\sqrt{6}}{23} \end{aligned}$$

$$\begin{aligned} 60. \quad \frac{\sqrt{2}}{\sqrt{7}+2} &= \frac{\sqrt{2}}{\sqrt{7}+2} \cdot \frac{\sqrt{7}-2}{\sqrt{7}-2} \\ &= \frac{\sqrt{2}(\sqrt{7}-2)}{7-4} \\ &= \frac{\sqrt{2}(\sqrt{7}-2)}{3} \quad \text{or} \quad \frac{\sqrt{14}-2\sqrt{2}}{3} \end{aligned}$$

$$\begin{aligned} 61. \quad \frac{2-\sqrt{5}}{2+3\sqrt{5}} &= \frac{2-\sqrt{5}}{2+3\sqrt{5}} \cdot \frac{2-3\sqrt{5}}{2-3\sqrt{5}} \\ &= \frac{4-2\sqrt{5}-6\sqrt{5}+15}{4-45} \\ &= \frac{19-8\sqrt{5}}{-41} = \frac{8\sqrt{5}-19}{41} \end{aligned}$$

$$\begin{aligned} 62. \quad \frac{\sqrt{3}-1}{2\sqrt{3}+3} &= \frac{\sqrt{3}-1}{2\sqrt{3}+3} \cdot \frac{2\sqrt{3}-3}{2\sqrt{3}-3} \\ &= \frac{6-2\sqrt{3}-3\sqrt{3}+3}{12-9} = \frac{9-5\sqrt{3}}{3} \end{aligned}$$

$$\begin{aligned} 63. \quad \frac{5}{\sqrt{2}-1} &= \frac{5}{\sqrt{2}-1} \cdot \frac{\sqrt{2}+1}{\sqrt{2}+1} \\ &= \frac{5\sqrt{2}+5}{2-1} = 5\sqrt{2}+5 \end{aligned}$$

$$\begin{aligned} 64. \quad \frac{-3}{\sqrt{5}+4} &= \frac{-3}{\sqrt{5}+4} \cdot \frac{\sqrt{5}-4}{\sqrt{5}-4} \\ &= \frac{-3\sqrt{5}+12}{5-16} = \frac{-3\sqrt{5}+12}{-11} \\ &= \frac{3\sqrt{5}-12}{11} \end{aligned}$$

$$65. \quad \frac{5}{\sqrt[3]{2}} = \frac{5}{\sqrt[3]{2}} \cdot \frac{\sqrt[3]{4}}{\sqrt[3]{4}} = \frac{5\sqrt[3]{4}}{2}$$

$$66. \quad \frac{-2}{\sqrt[3]{9}} = \frac{-2}{\sqrt[3]{9}} \cdot \frac{\sqrt[3]{3}}{\sqrt[3]{3}} = \frac{-2\sqrt[3]{3}}{3}$$

$$\begin{aligned} 67. \quad \frac{\sqrt{x+h}-\sqrt{x}}{\sqrt{x+h}+\sqrt{x}} &= \frac{\sqrt{x+h}-\sqrt{x}}{\sqrt{x+h}+\sqrt{x}} \cdot \frac{\sqrt{x+h}-\sqrt{x}}{\sqrt{x+h}-\sqrt{x}} \\ &= \frac{(x+h)-2\sqrt{x(x+h)}+x}{(x+h)-x} \\ &= \frac{x+h-2\sqrt{x^2+xh}+x}{x+h-x} \\ &= \frac{2x+h-2\sqrt{x^2+xh}}{h} \end{aligned}$$

$$\begin{aligned} 68. \quad \frac{\sqrt{x+h}+\sqrt{x-h}}{\sqrt{x+h}-\sqrt{x-h}} &= \frac{\sqrt{x+h}+\sqrt{x-h}}{\sqrt{x+h}-\sqrt{x-h}} \cdot \frac{\sqrt{x+h}+\sqrt{x-h}}{\sqrt{x+h}+\sqrt{x-h}} \\ &= \frac{(x+h)+2\sqrt{(x-h)(x+h)}+(x-h)}{(x+h)-(x-h)} \\ &= \frac{x+h+2\sqrt{x^2-h^2}+x-h}{x+h-x+h} \\ &= \frac{2x+2\sqrt{x^2-h^2}}{2h} \\ &= \frac{x+\sqrt{x^2-h^2}}{h} \end{aligned}$$

$$\begin{aligned} 69. \quad \frac{\sqrt{11}+1}{2} &= \frac{\sqrt{11}+1}{2} \cdot \frac{\sqrt{11}-1}{\sqrt{11}-1} \\ &= \frac{11-1}{2(\sqrt{11}-1)} = \frac{10}{2(\sqrt{11}-1)} \\ &= \frac{5}{\sqrt{11}-1} \end{aligned}$$

$$\begin{aligned} 70. \quad \frac{5-\sqrt{43}}{3} &= \frac{5-\sqrt{43}}{3} \cdot \frac{5+\sqrt{43}}{5+\sqrt{43}} = \frac{25-43}{3(5+\sqrt{43})} \\ &= \frac{-18}{3(5+\sqrt{43})} = \frac{-6}{5+\sqrt{43}} = -\frac{6}{5+\sqrt{43}} \end{aligned}$$

$$\begin{aligned} 71. \quad \frac{\sqrt{6}-\sqrt{15}}{\sqrt{15}} &= \frac{\sqrt{6}-\sqrt{15}}{\sqrt{15}} \cdot \frac{\sqrt{6}+\sqrt{15}}{\sqrt{6}+\sqrt{15}} \\ &= \frac{6-15}{\sqrt{90}-15} = \frac{-9}{3\sqrt{10}-15} \\ &= \frac{-9}{3(\sqrt{10}-5)} = -\frac{3}{\sqrt{10}-5} \end{aligned}$$

Section R.8: *n*th Roots; Rational Exponents

$$\begin{aligned} 72. \quad \frac{\sqrt{5}+\sqrt{3}}{\sqrt{5}} &= \frac{\sqrt{5}+\sqrt{3}}{\sqrt{5}} \cdot \frac{\sqrt{5}-\sqrt{3}}{\sqrt{5}-\sqrt{3}} \\ &= \frac{5-3}{5-\sqrt{15}} = \frac{2}{5-\sqrt{15}} \end{aligned}$$

$$\begin{aligned} 73. \quad \frac{\sqrt{x}-\sqrt{c}}{x-c} &= \frac{\sqrt{x}-\sqrt{c}}{x-c} \cdot \frac{\sqrt{x}+\sqrt{c}}{\sqrt{x}+\sqrt{c}} \\ &= \frac{x-c}{(x-c)(\sqrt{x}+\sqrt{c})} = \frac{1}{\sqrt{x}+\sqrt{c}} \end{aligned}$$

$$\begin{aligned} 74. \quad \frac{\sqrt{x}-2}{x-4} &= \frac{\sqrt{x}-2}{x-4} \cdot \frac{\sqrt{x}+2}{\sqrt{x}+2} \\ &= \frac{x-4}{(x-4)(\sqrt{x}+2)} = \frac{1}{\sqrt{x}+2} \end{aligned}$$

$$\begin{aligned} 75. \quad \frac{\sqrt{x-7}-1}{x-8} &= \frac{\sqrt{x-7}-1}{x-8} \cdot \frac{\sqrt{x-7}+1}{\sqrt{x-7}+1} \\ &= \frac{x-7-1}{(x-8)(\sqrt{x-7}+1)} \\ &= \frac{x-8}{(x-8)(\sqrt{x-7}+1)} \\ &= \frac{1}{\sqrt{x-7}+1} \end{aligned}$$

$$\begin{aligned} 76. \quad \frac{4-\sqrt{x-9}}{x-25} &= \frac{4-\sqrt{x-9}}{x-25} \cdot \frac{4+\sqrt{x-9}}{4+\sqrt{x-9}} \\ &= \frac{16-(x-9)}{(x-25)(4+\sqrt{x-9})} \\ &= \frac{25-x}{(x-25)(4+\sqrt{x-9})} \\ &= -\frac{x-25}{(x-25)(4+\sqrt{x-9})} \\ &= -\frac{1}{4+\sqrt{x-9}} \end{aligned}$$

$$77. \quad 8^{2/3} = (\sqrt[3]{8})^2 = 2^2 = 4$$

$$78. \quad 4^{3/2} = (\sqrt{4})^3 = 2^3 = 8$$

$$79. \quad (-64)^{1/3} = \sqrt[3]{-64} = -4$$

$$80. \quad 16^{3/4} = (\sqrt[4]{16})^3 = 2^3 = 8$$

$$81. \quad 100^{3/2} = (\sqrt{100})^3 = 10^3 = 1000$$

$$82. \quad 25^{3/2} = (\sqrt{25})^3 = 5^3 = 125$$

$$83. \quad 4^{-3/2} = \frac{1}{4^{3/2}} = \frac{1}{(\sqrt{4})^3} = \frac{1}{2^3} = \frac{1}{8}$$

$$84. \quad 16^{-3/2} = \frac{1}{16^{3/2}} = \frac{1}{(\sqrt{16})^3} = \frac{1}{4^3} = \frac{1}{64}$$

$$\begin{aligned} 85. \quad \left(\frac{9}{8}\right)^{3/2} &= \left(\sqrt{\frac{9}{8}}\right)^3 = \left(\frac{3}{2\sqrt{2}}\right)^3 = \frac{3^3}{2^3(\sqrt{2})^3} \\ &= \frac{27}{8 \cdot 2\sqrt{2}} = \frac{27}{16\sqrt{2}} = \frac{27}{16\sqrt{2}} \cdot \frac{\sqrt{2}}{\sqrt{2}} \\ &= \frac{27\sqrt{2}}{32} \end{aligned}$$

$$86. \quad \left(\frac{27}{8}\right)^{2/3} = \left(\sqrt[3]{\frac{27}{8}}\right)^2 = \left(\frac{3}{2}\right)^2 = \frac{9}{4}$$

$$\begin{aligned} 87. \quad \left(\frac{8}{9}\right)^{-3/2} &= \left(\frac{9}{8}\right)^{3/2} = \left(\sqrt{\frac{9}{8}}\right)^3 = \left(\frac{3}{2\sqrt{2}}\right)^3 \\ &= \frac{3^3}{2^3(\sqrt{2})^3} = \frac{27}{8 \cdot 2\sqrt{2}} = \frac{27}{16\sqrt{2}} \\ &= \frac{27}{16\sqrt{2}} \cdot \frac{\sqrt{2}}{\sqrt{2}} = \frac{27\sqrt{2}}{32} \end{aligned}$$

$$88. \quad \left(\frac{8}{27}\right)^{-2/3} = \left(\frac{27}{8}\right)^{2/3} = \left(\sqrt[3]{\frac{27}{8}}\right)^2 = \left(\frac{3}{2}\right)^2 = \frac{9}{4}$$

$$89. \quad (-1000)^{-1/3} = \left(-\frac{1}{1000}\right)^{1/3} = \sqrt[3]{-\frac{1}{1000}} = -\frac{1}{10}$$

$$90. \quad -25^{-1/2} = -\left(\frac{1}{25}\right)^{1/2} = -\sqrt{\frac{1}{25}} = -\frac{1}{5}$$

Chapter R: Review

$$91. \left(-\frac{64}{125}\right)^{-2/3} = \left(-\frac{125}{64}\right)^{2/3} = \left(\sqrt[3]{-\frac{125}{64}}\right)^2 \\ = \left(-\frac{5}{4}\right)^2 = \frac{25}{16}$$

$$92. -81^{-3/4} = -\left(\frac{1}{81}\right)^{3/4} = -\left(\sqrt[4]{\frac{1}{81}}\right)^3 \\ = -\left(\frac{1}{3}\right)^3 = -\frac{1}{27}$$

$$93. x^{3/4} x^{1/3} x^{-1/2} = x^{3/4+1/3-1/2} = x^{7/12}$$

$$94. x^{2/3} x^{1/2} x^{-1/4} = x^{2/3+1/2-1/4} = x^{11/12}$$

$$95. (x^3 y^6)^{1/3} = (x^3)^{1/3} (y^6)^{1/3} = xy^2$$

$$96. (x^4 y^8)^{3/4} = (x^4)^{3/4} (y^8)^{3/4} = x^3 y^6$$

$$97. \frac{(x^2 y)^{1/3} (xy^2)^{2/3}}{x^{2/3} y^{2/3}} = \frac{(x^2)^{1/3} (y)^{1/3} (x)^{2/3} (y^2)^{2/3}}{x^{2/3} y^{2/3}} \\ = \frac{x^{2/3} y^{1/3} x^{2/3} y^{4/3}}{x^{2/3} y^{2/3}} \\ = x^{2/3+2/3-2/3} y^{1/3+4/3-2/3} \\ = x^{2/3} y^1 = x^{2/3} y$$

$$98. \frac{(xy)^{1/4} (x^2 y^2)^{1/2}}{(x^2 y)^{3/4}} = \frac{x^{1/4} y^{1/4} (x^2)^{1/2} (y^2)^{1/2}}{(x^2)^{3/4} y^{3/4}} \\ = \frac{x^{1/4} y^{1/4} xy}{x^{3/2} y^{3/4}} \\ = x^{1/4+1-3/2} y^{1/4+1-3/4} \\ = x^{-1/4} y^{1/2} = \frac{y^{1/2}}{x^{1/4}}$$

$$99. \frac{(16x^2 y^{-1/3})^{3/4}}{(xy^2)^{1/4}} = \frac{16^{3/4} (x^2)^{3/4} (y^{-1/3})^{3/4}}{x^{1/4} (y^2)^{1/4}} \\ = \frac{(\sqrt[4]{16})^3 x^{3/2} y^{-1/4}}{x^{1/4} y^{1/2}} \\ = 2^3 x^{3/2-1/4} y^{-1/4-1/2} \\ = 8x^{5/4} y^{-3/4} \\ = \frac{8x^{5/4}}{y^{3/4}}$$

$$100. \frac{(4x^{-1} y^{1/3})^{3/2}}{(xy)^{3/2}} = \frac{4^{3/2} (x^{-1})^{3/2} (y^{1/3})^{3/2}}{x^{3/2} y^{3/2}} \\ = \frac{(\sqrt{4})^3 x^{-3/2} y^{1/2}}{x^{3/2} y^{3/2}} \\ = 2^3 x^{-3/2-3/2} y^{1/2-3/2} \\ = 8x^{-3} y^{-1} \\ = \frac{8}{x^3 y}$$

$$101. \frac{x}{(1+x)^{1/2}} + 2(1+x)^{1/2} = \frac{x+2(1+x)^{1/2} (1+x)^{1/2}}{(1+x)^{1/2}} \\ = \frac{x+2(1+x)}{(1+x)^{1/2}} \\ = \frac{x+2+2x}{(1+x)^{1/2}} \\ = \frac{3x+2}{(1+x)^{1/2}}$$

$$102. \frac{1+x}{2x^{1/2}} + x^{1/2} = \frac{1+x+x^{1/2} \cdot 2x^{1/2}}{2x^{1/2}} \\ = \frac{1+x+2x}{2x^{1/2}} = \frac{3x+1}{2x^{1/2}}$$

Section R.8: *n*th Roots; Rational Exponents

$$\begin{aligned}
 103. \quad & 2x(x^2+1)^{1/2} + x^2 \cdot \frac{1}{2}(x^2+1)^{-1/2} \cdot 2x \\
 &= 2x(x^2+1)^{1/2} + \frac{x^3}{(x^2+1)^{1/2}} \\
 &= \frac{2x(x^2+1)^{1/2} \cdot (x^2+1)^{1/2} + x^3}{(x^2+1)^{1/2}} \\
 &= \frac{2x(x^2+1)^{1/2+1/2} + x^3}{(x^2+1)^{1/2}} = \frac{2x(x^2+1)^1 + x^3}{(x^2+1)^{1/2}} \\
 &= \frac{2x^3 + 2x + x^3}{(x^2+1)^{1/2}} = \frac{3x^3 + 2x}{(x^2+1)^{1/2}} \\
 &= \frac{x(3x^2 + 2)}{(x^2+1)^{1/2}}
 \end{aligned}$$

$$\begin{aligned}
 104. \quad & (x+1)^{1/3} + x \cdot \frac{1}{3}(x+1)^{-2/3}, x \neq -1 \\
 &= (x+1)^{1/3} + \frac{x}{3(x+1)^{2/3}} \\
 &= \frac{3(x+1)^{2/3}(x+1)^{1/3} + x}{3(x+1)^{2/3}} \\
 &= \frac{3(x+1)^{2/3+1/3} + x}{3(x+1)^{2/3}} = \frac{3(x+1)^1 + x}{3(x+1)^{2/3}} \\
 &= \frac{3x+3+x}{3(x+1)^{2/3}} = \frac{4x+3}{3(x+1)^{2/3}}
 \end{aligned}$$

$$\begin{aligned}
 105. \quad & \sqrt{4x+3} \cdot \frac{1}{2\sqrt{x-5}} + \sqrt{x-5} \cdot \frac{1}{5\sqrt{4x+3}}, x > 5 \\
 &= \frac{\sqrt{4x+3}}{2\sqrt{x-5}} + \frac{\sqrt{x-5}}{5\sqrt{4x+3}} \\
 &= \frac{\sqrt{4x+3} \cdot 5 \cdot \sqrt{4x+3} + \sqrt{x-5} \cdot 2 \cdot \sqrt{x-5}}{10\sqrt{x-5}\sqrt{4x+3}} \\
 &= \frac{5(4x+3) + 2(x-5)}{10\sqrt{(x-5)(4x+3)}} \\
 &= \frac{20x+15+2x-10}{10\sqrt{(x-5)(4x+3)}} \\
 &= \frac{22x+5}{10\sqrt{(x-5)(4x+3)}}
 \end{aligned}$$

$$\begin{aligned}
 106. \quad & \frac{\sqrt[3]{8x+1}}{3\sqrt[3]{(x-2)^2}} + \frac{\sqrt[3]{x-2}}{24\sqrt[3]{(8x+1)^2}}, x \neq 2, x \neq -\frac{1}{8} \\
 &= \frac{8\sqrt[3]{8x+1} \cdot \sqrt[3]{(8x+1)^2} + \sqrt[3]{x-2} \cdot \sqrt[3]{(x-2)^2}}{24\sqrt[3]{(x-2)^2} \cdot \sqrt[3]{(8x+1)^2}} \\
 &= \frac{8\sqrt[3]{(8x+1)^3} + \sqrt[3]{(x-2)^3}}{24\sqrt[3]{(x-2)^2} \cdot \sqrt[3]{(8x+1)^2}} \\
 &= \frac{8(8x+1) + x-2}{24\sqrt[3]{(x-2)^2(8x+1)^2}} \\
 &= \frac{64x+8+x-2}{24\sqrt[3]{(x-2)^2(8x+1)^2}} \\
 &= \frac{65x+6}{24\sqrt[3]{(x-2)^2(8x+1)^2}}
 \end{aligned}$$

$$\begin{aligned}
 107. \quad & \frac{\left(\sqrt{1+x} - x \cdot \frac{1}{2\sqrt{1+x}}\right)}{1+x} = \frac{\left(\sqrt{1+x} - \frac{x}{2\sqrt{1+x}}\right)}{1+x} \\
 &= \frac{\left(\frac{2\sqrt{1+x}\sqrt{1+x} - x}{2\sqrt{1+x}}\right)}{1+x} \\
 &= \frac{2(1+x) - x}{2(1+x)^{1/2}} \cdot \frac{1}{1+x} \\
 &= \frac{2+x}{2(1+x)^{3/2}}
 \end{aligned}$$

$$\begin{aligned}
 108. \quad & \frac{\left(\sqrt{x^2+1} - x \cdot \frac{2x}{2\sqrt{x^2+1}}\right)}{x^2+1} \\
 &= \frac{\left(\sqrt{x^2+1} - \frac{x^2}{\sqrt{x^2+1}}\right)}{x^2+1} \\
 &= \frac{\left(\sqrt{x^2+1} \cdot \frac{\sqrt{x^2+1}}{\sqrt{x^2+1}} - \frac{x^2}{\sqrt{x^2+1}}\right)}{x^2+1} \\
 &= \frac{\left(\frac{x^2+1-x^2}{\sqrt{x^2+1}}\right)}{x^2+1} = \frac{1}{\sqrt{x^2+1}} \cdot \frac{1}{x^2+1} \\
 &= \frac{1}{(x^2+1)^{3/2}}
 \end{aligned}$$

Chapter R: Review

$$\begin{aligned}
 109. \quad & \frac{(x+4)^{1/2} - 2x(x+4)^{-1/2}}{x+4} \\
 &= \frac{\left((x+4)^{1/2} - \frac{2x}{(x+4)^{1/2}} \right)}{x+4} \\
 &= \frac{\left((x+4)^{1/2} \cdot \frac{(x+4)^{1/2}}{(x+4)^{1/2}} - \frac{2x}{(x+4)^{1/2}} \right)}{x+4} \\
 &= \frac{\left(\frac{x+4-2x}{(x+4)^{1/2}} \right)}{x+4} \\
 &= \frac{-x+4}{(x+4)^{1/2}} \cdot \frac{1}{x+4} \\
 &= \frac{-x+4}{(x+4)^{3/2}} \\
 &= \frac{4-x}{(x+4)^{3/2}}
 \end{aligned}$$

$$\begin{aligned}
 110. \quad & \frac{(9-x^2)^{1/2} + x^2(9-x^2)^{-1/2}}{9-x^2}, -3 < x < 3 \\
 &= \frac{\left((9-x^2)^{1/2} + \frac{x^2}{(9-x^2)^{1/2}} \right)}{9-x^2} \\
 &= \frac{\left(\frac{(9-x^2)^{1/2} \cdot (9-x^2)^{1/2} + x^2}{(9-x^2)^{1/2}} \right)}{9-x^2} \\
 &= \frac{(9-x^2)^{1/2} \cdot (9-x^2)^{1/2} + x^2}{(9-x^2)^{1/2}} \cdot \frac{1}{9-x^2} \\
 &= \frac{9-x^2+x^2}{(9-x^2)^{1/2}} \cdot \frac{1}{9-x^2} \\
 &= \frac{9}{(9-x^2)^{3/2}}
 \end{aligned}$$

$$\begin{aligned}
 111. \quad & \frac{\frac{x^2}{(x^2-1)^{1/2}} - (x^2-1)^{1/2}}{x^2}, x < -1 \text{ or } x > 1 \\
 &= \frac{\left(\frac{x^2 - (x^2-1)^{1/2} \cdot (x^2-1)^{1/2}}{(x^2-1)^{1/2}} \right)}{x^2} \\
 &= \frac{x^2 - (x^2-1)^{1/2} \cdot (x^2-1)^{1/2}}{(x^2-1)^{1/2}} \cdot \frac{1}{x^2} \\
 &= \frac{x^2 - (x^2-1)}{(x^2-1)^{1/2}} \cdot \frac{1}{x^2} \\
 &= \frac{x^2 - x^2 + 1}{(x^2-1)^{1/2}} \cdot \frac{1}{x^2} \\
 &= \frac{1}{x^2(x^2-1)^{1/2}}
 \end{aligned}$$

$$\begin{aligned}
 112. \quad & \frac{(x^2+4)^{1/2} - x^2(x^2+4)^{-1/2}}{x^2+4} \\
 &= \frac{\left((x^2+4)^{1/2} - \frac{x^2}{(x^2+4)^{1/2}} \right)}{x^2+4} \\
 &= \frac{\left(\frac{(x^2+4)^{1/2} \cdot (x^2+4)^{1/2} - x^2}{(x^2+4)^{1/2}} \right)}{x^2+4} \\
 &= \frac{(x^2+4)^{1/2} \cdot (x^2+4)^{1/2} - x^2}{(x^2+4)^{1/2}} \cdot \frac{1}{x^2+4} \\
 &= \frac{x^2+4-x^2}{(x^2+4)^{1/2}} \cdot \frac{1}{x^2+4} = \frac{4}{(x^2+4)^{3/2}}
 \end{aligned}$$

Section R.8: nth Roots; Rational Exponents

$$\begin{aligned}
 113. \quad & \frac{\frac{1+x^2}{2\sqrt{x}} - 2x\sqrt{x}}{(1+x^2)^2}, x > 0 \\
 & = \frac{\left(\frac{1+x^2 - (2\sqrt{x})(2x\sqrt{x})}{2\sqrt{x}} \right)}{(1+x^2)^2} \\
 & = \frac{1+x^2 - (2\sqrt{x})(2x\sqrt{x})}{2\sqrt{x}} \cdot \frac{1}{(1+x^2)^2} \\
 & = \frac{1+x^2 - 4x^2}{2\sqrt{x}} \cdot \frac{1}{(1+x^2)^2} = \frac{1-3x^2}{2\sqrt{x}(1+x^2)^2}
 \end{aligned}$$

$$\begin{aligned}
 114. \quad & \frac{2x(1-x^2)^{1/3} + \frac{2}{3}x^3(1-x^2)^{-2/3}}{(1-x^2)^{2/3}}, x \neq -1, x \neq 1 \\
 & = \frac{\left(2x(1-x^2)^{1/3} + \frac{2x^3}{3(1-x^2)^{2/3}} \right)}{(1-x^2)^{2/3}} \\
 & = \frac{\left(\frac{2x(1-x^2)^{1/3} \cdot 3(1-x^2)^{2/3} + 2x^3}{3(1-x^2)^{2/3}} \right)}{(1-x^2)^{2/3}} \\
 & = \frac{6x(1-x^2)^{1/3+2/3} + 2x^3}{3(1-x^2)^{2/3}} \cdot \frac{1}{(1-x^2)^{2/3}} \\
 & = \frac{6x(1-x^2) + 2x^3}{3(1-x^2)^{2/3+2/3}} = \frac{6x - 6x^3 + 2x^3}{3(1-x^2)^{4/3}} \\
 & = \frac{6x - 4x^3}{3(1-x^2)^{4/3}} = \frac{2x(3-2x^2)}{3(1-x^2)^{4/3}}
 \end{aligned}$$

$$\begin{aligned}
 115. \quad & (x+1)^{3/2} + x \cdot \frac{3}{2}(x+1)^{1/2} \\
 & = (x+1)^{1/2} \left(x+1 + \frac{3}{2}x \right) \\
 & = (x+1)^{1/2} \left(\frac{5}{2}x + 1 \right) \\
 & = \frac{1}{2}(x+1)^{1/2} (5x+2)
 \end{aligned}$$

$$\begin{aligned}
 116. \quad & (x^2+4)^{4/3} + x \cdot \frac{4}{3}(x^2+4)^{1/3} \cdot 2x \\
 & = (x^2+4)^{1/3} \left(x^2+4 + \frac{8}{3}x^2 \right) \\
 & = (x^2+4)^{1/3} \left(\frac{11}{3}x^2 + 4 \right) \\
 & = \frac{1}{3}(x^2+4)^{1/3} (11x^2+12)
 \end{aligned}$$

$$\begin{aligned}
 117. \quad & 6x^{1/2}(x^2+x) - 8x^{3/2} - 8x^{1/2} \\
 & = 2x^{1/2}(3(x^2+x) - 4x - 4) \\
 & = 2x^{1/2}(3x^2 - x - 4) \\
 & = 2x^{1/2}(3x-4)(x+1)
 \end{aligned}$$

$$\begin{aligned}
 118. \quad & 6x^{1/2}(2x+3) + x^{3/2} \cdot 8 \\
 & = 2x^{1/2}(3(2x+3) + 4x) \\
 & = 2x^{1/2}(10x+9)
 \end{aligned}$$

$$\begin{aligned}
 119. \quad & 3(x^2+4)^{4/3} + x \cdot 4(x^2+4)^{1/3} \cdot 2x \\
 & = (x^2+4)^{1/3} [3(x^2+4) + 8x^2] \\
 & = (x^2+4)^{1/3} [3x^2 + 12 + 8x^2] \\
 & = (x^2+4)^{1/3} (11x^2 + 12)
 \end{aligned}$$

$$\begin{aligned}
 120. \quad & 2x(3x+4)^{4/3} + x^2 \cdot 4(3x+4)^{1/3} \\
 & = 2x(3x+4)^{1/3} [(3x+4) + 2x] \\
 & = 2x(3x+4)^{1/3} (5x+4)
 \end{aligned}$$

$$\begin{aligned}
 121. \quad & 4(3x+5)^{1/3}(2x+3)^{3/2} + 3(3x+5)^{4/3}(2x+3)^{1/2} \\
 & = (3x+5)^{1/3}(2x+3)^{1/2} [4(2x+3) + 3(3x+5)] \\
 & = (3x+5)^{1/3}(2x+3)^{1/2} (8x+12+9x+15) \\
 & = (3x+5)^{1/3}(2x+3)^{1/2} (17x+27)
 \end{aligned}$$


where $x \geq -\frac{3}{2}$.


Chapter R: Review


$$\begin{aligned}
 122. \quad & 6(6x+1)^{1/3}(4x-3)^{3/2} + 6(6x+1)^{4/3}(4x-3)^{1/2} \\
 &= 6(6x+1)^{1/3}(4x-3)^{1/2}[(4x-3) + (6x+1)] \\
 &= 6(6x+1)^{1/3}(4x-3)^{1/2}(10x-2) \\
 &= 6(6x+1)^{1/3}(4x-3)^{1/2}(2)(5x-1) \\
 &= 12(6x+1)^{1/3}(4x-3)^{1/2}(5x-1) \\
 &\text{where } x \geq \frac{3}{4}.
 \end{aligned}$$


$$\begin{aligned}
 123. \quad & 3x^{-1/2} + \frac{3}{2}x^{1/2}, x > 0 \\
 &= \frac{3}{x^{1/2}} + \frac{3}{2}x^{1/2} \\
 &= \frac{3 \cdot 2 + 3x^{1/2} \cdot x^{1/2}}{2x^{1/2}} = \frac{6 + 3x}{2x^{1/2}} = \frac{3(x+2)}{2x^{1/2}}
 \end{aligned}$$


$$\begin{aligned}
 124. \quad & 8x^{1/3} - 4x^{-2/3}, x \neq 0 \\
 &= 8x^{1/3} - \frac{4}{x^{2/3}} \\
 &= \frac{8x^{1/3} \cdot x^{2/3} - 4}{x^{2/3}} = \frac{8x - 4}{x^{2/3}} = \frac{4(2x-1)}{x^{2/3}}
 \end{aligned}$$


$$125. \quad \sqrt{2} \approx 1.41$$



$$126. \quad \sqrt{7} \approx 2.65$$



$$127. \quad \sqrt[3]{4} \approx 1.59$$


$$128. \quad \sqrt[3]{-5} \approx -1.71$$


$$129. \quad \frac{2+\sqrt{3}}{3-\sqrt{5}} \approx 4.89$$


$$130. \quad \frac{\sqrt{5}-2}{\sqrt{2}+4} \approx 0.04$$


$$131. \quad \frac{3\sqrt[3]{5}-\sqrt{2}}{\sqrt{3}} \approx 2.15$$


$$132. \quad \frac{2\sqrt{3}-\sqrt[3]{4}}{\sqrt{2}} \approx 1.33$$


$$133. \quad \text{a. } V = 40(12)^2 \sqrt{\frac{96}{12}} - 0.608 \approx 15,660.4 \text{ gallons}$$

$$\text{b. } V = 40(1)^2 \sqrt{\frac{96}{1}} - 0.608 \approx 390.7 \text{ gallons}$$

$$134. \quad \text{a. } v = \sqrt{64 \cdot 4 + 0^2} = \sqrt{256} = 16 \text{ feet per second}$$

$$\text{b. } v = \sqrt{64 \cdot 16 + 0^2} = \sqrt{1024} = 32 \text{ feet per second}$$

$$\text{c. } v = \sqrt{64 \cdot 2 + 4^2} = \sqrt{144} = 12 \text{ feet per second}$$

$$135. \quad T = 2\pi \sqrt{\frac{64}{32}} = 2\pi \sqrt{2} \approx 8.89 \text{ seconds}$$

Section R.8: nth Roots; Rational Exponents

$$\begin{aligned} 136. \quad T &= 2\pi\sqrt{\frac{16}{32}} = 2\pi\sqrt{\frac{1}{2}} = \frac{2\pi}{\sqrt{2}} \\ &= \pi\sqrt{2} \approx 4.44 \text{ seconds} \end{aligned}$$

$$\begin{array}{r} 137. \quad \sqrt{3} \overline{) 1 \quad -4 \quad -3 \quad 13} \\ \underline{ \sqrt{3} \quad 3-4\sqrt{3} \quad -12} \\ 1 \quad \sqrt{3}-4 \quad -4\sqrt{3} \quad 1 \end{array}$$

The quotient is $x^2 + (\sqrt{3} - 4)x - 4\sqrt{3}$

The remainder is 1

$$\begin{array}{r} 138. \quad 1+\sqrt{2} \overline{) 1 \quad -9 \quad 13 \quad 7} \\ \underline{ 1+\sqrt{2} \quad -6-7\sqrt{2} \quad -7} \\ 1 \quad \sqrt{3}-8 \quad -7-7\sqrt{2} \quad 0 \end{array}$$

Yes, $1 + \sqrt{2}$ is a factor of $x^3 - 9x^2 + 13x + 7$.

139. Answers may vary. One possibility follows: If

$$a = -5, \text{ then } \sqrt{a^2} = \sqrt{(-5)^2} = \sqrt{25} = 5 \neq a.$$

Since we use the principal square root, which is always non-negative,

$$\sqrt{a^2} = \begin{cases} a & \text{if } a \geq 0 \\ -a & \text{if } a < 0 \end{cases}$$

which is the definition of $|a|$, so $\sqrt{a^2} = |a|$.