


Solutions for Introduction to Quantum Mechanics 2nd Edition by Griffiths

[CLICK HERE TO ACCESS COMPLETE Solutions](#)



Solutions

Chapter 2

The Time-Independent Schrödinger Equation

Problem 2.1

(a)

$$\Psi(x, t) = \psi(x)e^{-i(E_0 + i\Gamma)t/\hbar} = \psi(x)e^{\Gamma t/\hbar}e^{-iE_0 t/\hbar} \implies |\Psi|^2 = |\psi|^2 e^{2\Gamma t/\hbar}.$$

$$\int_{-\infty}^{\infty} |\Psi(x, t)|^2 dx = e^{2\Gamma t/\hbar} \int_{-\infty}^{\infty} |\psi|^2 dx.$$

The second term is independent of t , so if the product is to be 1 for all time, the first term ($e^{2\Gamma t/\hbar}$) must also be constant, and hence $\Gamma = 0$. QED

(b) If ψ satisfies Eq. 2.5, $-\frac{\hbar^2}{2m} \frac{d^2\psi}{dx^2} + V\psi = E\psi$, then (taking the complex conjugate and noting that V and E are real): $-\frac{\hbar^2}{2m} \frac{d^2\psi^*}{dx^2} + V\psi^* = E\psi^*$, so ψ^* also satisfies Eq. 2.5. Now, if ψ_1 and ψ_2 satisfy Eq. 2.5, so too does any linear combination of them ($\psi_3 \equiv c_1\psi_1 + c_2\psi_2$):

$$\begin{aligned} -\frac{\hbar^2}{2m} \frac{d^2\psi_3}{dx^2} + V\psi_3 &= -\frac{\hbar^2}{2m} \left(c_1 \frac{d^2\psi_1}{dx^2} + c_2 \frac{d^2\psi_2}{dx^2} \right) + V(c_1\psi_1 + c_2\psi_2) \\ &= c_1 \left[-\frac{\hbar^2}{2m} \frac{d^2\psi_1}{dx^2} + V\psi_1 \right] + c_2 \left[-\frac{\hbar^2}{2m} \frac{d^2\psi_2}{dx^2} + V\psi_2 \right] \\ &= c_1(E\psi_1) + c_2(E\psi_2) = E(c_1\psi_1 + c_2\psi_2) = E\psi_3. \end{aligned}$$

Thus, $(\psi + \psi^*)$ and $i(\psi - \psi^*)$ – both of which are *real* – satisfy Eq. 2.5. *Conclusion:* From any complex solution, we can always construct two *real* solutions (of course, if ψ is already real, the second one will be zero). In particular, since $\psi = \frac{1}{2}[(\psi + \psi^*) - i(i(\psi - \psi^*))]$, ψ can be expressed as a linear combination of two real solutions. QED


(c) If $\psi(x)$ satisfies Eq. 2.5, then, changing variables $x \rightarrow -x$ and noting that $d^2/d(-x)^2 = d^2/dx^2$,

$$-\frac{\hbar^2}{2m} \frac{d^2\psi(-x)}{dx^2} + V(-x)\psi(-x) = E\psi(-x);$$

so if $V(-x) = V(x)$ then $\psi(-x)$ also satisfies Eq. 2.5. It follows that $\psi_+(x) \equiv \psi(x) + \psi(-x)$ (which is *even*: $\psi_+(-x) = \psi_+(x)$) and $\psi_-(x) \equiv \psi(x) - \psi(-x)$ (which is *odd*: $\psi_-(-x) = -\psi_-(x)$) both satisfy Eq. 2.5. But $\psi(x) = \frac{1}{2}(\psi_+(x) + \psi_-(x))$, so any solution can be expressed as a linear combination of even and odd solutions. QED

Problem 2.2

Given $\frac{d^2\psi}{dx^2} = \frac{2m}{\hbar^2}[V(x) - E]\psi$, if $E < V_{\min}$, then ψ'' and ψ always have the same sign: If ψ is positive(negative), then ψ'' is also positive(negative). This means that ψ always curves away from the axis (see Figure). However, it has got to go to zero as $x \rightarrow -\infty$ (else it would not be normalizable). At some point it's got to *depart* from zero (if it *doesn't*, it's going to be identically zero *everywhere*), in (say) the positive direction. At this point its slope is positive, and *increasing*, so ψ gets bigger and bigger as x increases. It can't ever "turn over" and head back toward the axis, because that would require a negative second derivative—it always has to bend away from the axis. By the same token, if it starts out heading negative, it just runs more and more negative. In neither case is there any way for it to come back to zero, as it must (at $x \rightarrow \infty$) in order to be normalizable. QED



Problem 2.3

Equation 2.20 says $\frac{d^2\psi}{dx^2} = -\frac{2mE}{\hbar^2}\psi$; Eq. 2.23 says $\psi(0) = \psi(a) = 0$. If $E = 0$, $d^2\psi/dx^2 = 0$, so $\psi(x) = A + Bx$; $\psi(0) = A = 0 \Rightarrow \psi = Bx$; $\psi(a) = Ba = 0 \Rightarrow B = 0$, so $\psi = 0$. If $E < 0$, $d^2\psi/dx^2 = \kappa^2\psi$, with $\kappa \equiv \sqrt{-2mE}/\hbar$ real, so $\psi(x) = Ae^{\kappa x} + Be^{-\kappa x}$. This time $\psi(0) = A + B = 0 \Rightarrow B = -A$, so $\psi = A(e^{\kappa x} - e^{-\kappa x})$, while $\psi(a) = A(e^{\kappa a} - e^{-\kappa a}) = 0 \Rightarrow$ either $A = 0$, so $\psi = 0$, or else $e^{\kappa a} = e^{-\kappa a}$, so $e^{2\kappa a} = 1$, so $2\kappa a = \ln(1) = 0$, so $\kappa = 0$, and again $\psi = 0$. In all cases, then, the boundary conditions force $\psi = 0$, which is unacceptable (non-normalizable).

Problem 2.4

$$\begin{aligned} \langle x \rangle &= \int x |\psi|^2 dx = \frac{2}{a} \int_0^a x \sin^2 \left(\frac{n\pi}{a} x \right) dx. \quad \text{Let } y \equiv \frac{n\pi}{a} x, \text{ so } dx = \frac{a}{n\pi} dy; \quad y : 0 \rightarrow n\pi. \\ &= \frac{2}{a} \left(\frac{a}{n\pi} \right)^2 \int_0^{n\pi} y \sin^2 y dy = \frac{2a}{n^2\pi^2} \left[\frac{y^2}{4} - \frac{y \sin 2y}{4} - \frac{\cos 2y}{8} \right] \Big|_0^{n\pi} \\ &= \frac{2a}{n^2\pi^2} \left[\frac{n^2\pi^2}{4} - \frac{\cos 2n\pi}{8} + \frac{1}{8} \right] = \boxed{\frac{a}{2}} \quad (\text{Independent of } n.) \end{aligned}$$

$$\begin{aligned} \langle x^2 \rangle &= \frac{2}{a} \int_0^a x^2 \sin^2 \left(\frac{n\pi}{a} x \right) dx = \frac{2}{a} \left(\frac{a}{n\pi} \right)^3 \int_0^{n\pi} y^2 \sin^2 y dy \\ &= \frac{2a^2}{(n\pi)^3} \left[\frac{y^3}{6} - \left(\frac{y^2}{4} - \frac{1}{8} \right) \sin 2y - \frac{y \cos 2y}{4} \right] \Big|_0^{n\pi} \\ &= \frac{2a^2}{(n\pi)^3} \left[\frac{(n\pi)^3}{6} - \frac{n\pi \cos(2n\pi)}{4} \right] = \boxed{a^2 \left[\frac{1}{3} - \frac{1}{2(n\pi)^2} \right]}. \end{aligned}$$

$$\langle p \rangle = m \frac{d\langle x \rangle}{dt} = \boxed{0}. \quad (\text{Note : Eq. 1.33 is much faster than Eq. 1.35.})$$

$$\begin{aligned} \langle p^2 \rangle &= \int \psi_n^* \left(\frac{\hbar}{i} \frac{d}{dx} \right)^2 \psi_n dx = -\hbar^2 \int \psi_n^* \left(\frac{d^2 \psi_n}{dx^2} \right) dx \\ &= (-\hbar^2) \left(-\frac{2mE_n}{\hbar^2} \right) \int \psi_n^* \psi_n dx = 2mE_n = \boxed{\left(\frac{n\pi\hbar}{a} \right)^2}. \end{aligned}$$

$$\sigma_x^2 = \langle x^2 \rangle - \langle x \rangle^2 = a^2 \left(\frac{1}{3} - \frac{1}{2(n\pi)^2} - \frac{1}{4} \right) = \frac{a^2}{4} \left(\frac{1}{3} - \frac{2}{(n\pi)^2} \right); \quad \boxed{\sigma_x = \frac{a}{2} \sqrt{\frac{1}{3} - \frac{2}{(n\pi)^2}}}.$$

$$\sigma_p^2 = \langle p^2 \rangle - \langle p \rangle^2 = \left(\frac{n\pi\hbar}{a} \right)^2; \quad \boxed{\sigma_p = \frac{n\pi\hbar}{a}}. \quad \therefore \sigma_x \sigma_p = \boxed{\frac{\hbar}{2} \sqrt{\frac{(n\pi)^2}{3} - 2}}.$$

The product $\sigma_x \sigma_p$ is smallest for $n = 1$; in that case, $\sigma_x \sigma_p = \frac{\hbar}{2} \sqrt{\frac{\pi^2}{3} - 2} = (1.136)\hbar/2 > \hbar/2. \quad \checkmark$

Problem 2.5

(a)

$$|\Psi|^2 = \Psi^* \Psi = |A|^2 (\psi_1^* + \psi_2^*) (\psi_1 + \psi_2) = |A|^2 [\psi_1^* \psi_1 + \psi_1^* \psi_2 + \psi_2^* \psi_1 + \psi_2^* \psi_2].$$

$$1 = \int |\Psi|^2 dx = |A|^2 \int [|\psi_1|^2 + \psi_1^* \psi_2 + \psi_2^* \psi_1 + |\psi_2|^2] dx = 2|A|^2 \Rightarrow \boxed{A = 1/\sqrt{2}}.$$

(b)

$$\Psi(x, t) = \frac{1}{\sqrt{2}} \left[\psi_1 e^{-iE_1 t/\hbar} + \psi_2 e^{-iE_2 t/\hbar} \right] \quad (\text{but } \frac{E_n}{\hbar} = n^2 \omega)$$

$$= \frac{1}{\sqrt{2}} \sqrt{\frac{2}{a}} \left[\sin\left(\frac{\pi}{a}x\right) e^{-i\omega t} + \sin\left(\frac{2\pi}{a}x\right) e^{-i4\omega t} \right] = \boxed{\frac{1}{\sqrt{a}} e^{-i\omega t} \left[\sin\left(\frac{\pi}{a}x\right) + \sin\left(\frac{2\pi}{a}x\right) e^{-3i\omega t} \right]}.$$

$$\begin{aligned} |\Psi(x, t)|^2 &= \frac{1}{a} \left[\sin^2\left(\frac{\pi}{a}x\right) + \sin\left(\frac{\pi}{a}x\right) \sin\left(\frac{2\pi}{a}x\right) (e^{-3i\omega t} + e^{3i\omega t}) + \sin^2\left(\frac{2\pi}{a}x\right) \right] \\ &= \boxed{\frac{1}{a} \left[\sin^2\left(\frac{\pi}{a}x\right) + \sin^2\left(\frac{2\pi}{a}x\right) + 2 \sin\left(\frac{\pi}{a}x\right) \sin\left(\frac{2\pi}{a}x\right) \cos(3\omega t) \right]}. \end{aligned}$$

(c)

$$\begin{aligned} \langle x \rangle &= \int x |\Psi(x, t)|^2 dx \\ &= \frac{1}{a} \int_0^a x \left[\sin^2\left(\frac{\pi}{a}x\right) + \sin^2\left(\frac{2\pi}{a}x\right) + 2 \sin\left(\frac{\pi}{a}x\right) \sin\left(\frac{2\pi}{a}x\right) \cos(3\omega t) \right] dx \end{aligned}$$

$$\int_0^a x \sin^2\left(\frac{\pi}{a}x\right) dx = \left[\frac{x^2}{4} - \frac{x \sin\left(\frac{2\pi}{a}x\right)}{4\pi/a} - \frac{\cos\left(\frac{2\pi}{a}x\right)}{8(\pi/a)^2} \right] \Big|_0^a = \frac{a^2}{4} = \int_0^a x \sin^2\left(\frac{2\pi}{a}x\right) dx.$$

$$\begin{aligned} \int_0^a x \sin\left(\frac{\pi}{a}x\right) \sin\left(\frac{2\pi}{a}x\right) dx &= \frac{1}{2} \int_0^a x \left[\cos\left(\frac{\pi}{a}x\right) - \cos\left(\frac{3\pi}{a}x\right) \right] dx \\ &= \frac{1}{2} \left[\frac{a^2}{\pi^2} \cos\left(\frac{\pi}{a}x\right) + \frac{ax}{\pi} \sin\left(\frac{\pi}{a}x\right) - \frac{a^2}{9\pi^2} \cos\left(\frac{3\pi}{a}x\right) - \frac{ax}{3\pi} \sin\left(\frac{3\pi}{a}x\right) \right]_0^a \\ &= \frac{1}{2} \left[\frac{a^2}{\pi^2} (\cos(\pi) - \cos(0)) - \frac{a^2}{9\pi^2} (\cos(3\pi) - \cos(0)) \right] = -\frac{a^2}{\pi^2} \left(1 - \frac{1}{9} \right) = -\frac{8a^2}{9\pi^2}. \end{aligned}$$

$$\therefore \langle x \rangle = \frac{1}{a} \left[\frac{a^2}{4} + \frac{a^2}{4} - \frac{16a^2}{9\pi^2} \cos(3\omega t) \right] = \boxed{\frac{a}{2} \left[1 - \frac{32}{9\pi^2} \cos(3\omega t) \right]}.$$

Amplitude: $\boxed{\frac{32}{9\pi^2} \left(\frac{a}{2} \right) = 0.3603(a/2)}$; angular frequency: $\boxed{3\omega = \frac{3\pi^2\hbar}{2ma^2}}.$

(d)

$$\langle p \rangle = m \frac{d\langle x \rangle}{dt} = m \left(\frac{a}{2} \right) \left(-\frac{32}{9\pi^2} \right) (-3\omega) \sin(3\omega t) = \boxed{\frac{8\hbar}{3a} \sin(3\omega t)}.$$

(e) You could get either $\boxed{E_1 = \pi^2\hbar^2/2ma^2}$ or $\boxed{E_2 = 2\pi^2\hbar^2/ma^2}$, with equal probability $\boxed{P_1 = P_2 = 1/2}$.

So $\langle H \rangle = \boxed{\frac{1}{2}(E_1 + E_2) = \frac{5\pi^2\hbar^2}{4ma^2}}$; it's the *average* of E_1 and E_2 .

Problem 2.6

From Problem 2.5, we see that

$$\Psi(x, t) = \boxed{\frac{1}{\sqrt{a}} e^{-i\omega t} \left[\sin\left(\frac{\pi}{a}x\right) + \sin\left(\frac{2\pi}{a}x\right) e^{-3i\omega t} e^{i\phi} \right]};$$


$$|\Psi(x, t)|^2 = \boxed{\frac{1}{a} \left[\sin^2\left(\frac{\pi}{a}x\right) + \sin^2\left(\frac{2\pi}{a}x\right) + 2 \sin\left(\frac{\pi}{a}x\right) \sin\left(\frac{2\pi}{a}x\right) \cos(3\omega t - \phi) \right]};$$

and hence $\boxed{\langle x \rangle = \frac{a}{2} \left[1 - \frac{32}{9\pi^2} \cos(3\omega t - \phi) \right]}$. This amounts physically to starting the clock at a different time (i.e., shifting the $t = 0$ point).

If $\phi = \frac{\pi}{2}$, so $\Psi(x, 0) = A[\psi_1(x) + i\psi_2(x)]$, then $\cos(3\omega t - \phi) = \sin(3\omega t)$; $\langle x \rangle$ starts at $\frac{a}{2}$.

If $\phi = \pi$, so $\Psi(x, 0) = A[\psi_1(x) - \psi_2(x)]$, then $\cos(3\omega t - \phi) = -\cos(3\omega t)$; $\langle x \rangle$ starts at $\frac{a}{2} \left(1 + \frac{32}{9\pi^2} \right)$.

Problem 2.7



(a)

$$\begin{aligned}
 1 &= A^2 \int_0^{a/2} x^2 dx + A^2 \int_{a/2}^a (a-x)^2 dx = A^2 \left[\frac{x^3}{3} \Big|_0^{a/2} - \frac{(a-x)^3}{3} \Big|_{a/2}^a \right] \\
 &= \frac{A^2}{3} \left(\frac{a^3}{8} + \frac{a^3}{8} \right) = \frac{A^2 a^3}{12} \Rightarrow \boxed{A = \frac{2\sqrt{3}}{\sqrt{a^3}}}.
 \end{aligned}$$

(b)

$$\begin{aligned}
 c_n &= \sqrt{\frac{2}{a}} \frac{2\sqrt{3}}{a\sqrt{a}} \left[\int_0^{a/2} x \sin\left(\frac{n\pi}{a}x\right) dx + \int_{a/2}^a (a-x) \sin\left(\frac{n\pi}{a}x\right) dx \right] \\
 &= \frac{2\sqrt{6}}{a^2} \left\{ \left[\left(\frac{a}{n\pi}\right)^2 \sin\left(\frac{n\pi}{a}x\right) - \frac{xa}{n\pi} \cos\left(\frac{n\pi}{a}x\right) \right] \Big|_0^{a/2} \right. \\
 &\quad \left. + a \left[-\frac{a}{n\pi} \cos\left(\frac{n\pi}{a}x\right) \right] \Big|_{a/2}^a - \left[\left(\frac{a}{n\pi}\right)^2 \sin\left(\frac{n\pi}{a}x\right) - \left(\frac{ax}{n\pi}\right) \cos\left(\frac{n\pi}{a}x\right) \right] \Big|_{a/2}^a \right\} \\
 &= \frac{2\sqrt{6}}{a^2} \left[\left(\frac{a}{n\pi}\right)^2 \sin\left(\frac{n\pi}{2}\right) - \frac{a^2}{2n\pi} \cos\left(\frac{n\pi}{2}\right) - \frac{a^2}{n\pi} \cos n\pi + \frac{a^2}{n\pi} \cos\left(\frac{n\pi}{2}\right) \right. \\
 &\quad \left. + \left(\frac{a}{n\pi}\right)^2 \sin\left(\frac{n\pi}{2}\right) + \frac{a^2}{n\pi} \cos n\pi - \frac{a^2}{2n\pi} \cos\left(\frac{n\pi}{2}\right) \right] \\
 &= \frac{2\sqrt{6}}{a^2} \frac{2}{(n\pi)^2} \sin\left(\frac{n\pi}{2}\right) = \frac{4\sqrt{6}}{(n\pi)^2} \sin\left(\frac{n\pi}{2}\right) = \begin{cases} 0, & n \text{ even,} \\ (-1)^{(n-1)/2} \frac{4\sqrt{6}}{(n\pi)^2}, & n \text{ odd.} \end{cases}
 \end{aligned}$$

$$\text{So } \Psi(x,t) = \frac{4\sqrt{6}}{\pi^2} \sqrt{\frac{2}{a}} \sum_{n=1,3,5,\dots} (-1)^{(n-1)/2} \frac{1}{n^2} \sin\left(\frac{n\pi}{a}x\right) e^{-iE_n t/\hbar}, \text{ where } E_n = \frac{n^2 \pi^2 \hbar^2}{2ma^2}.$$

(c)

$$P_1 = |c_1|^2 = \frac{16 \cdot 6}{\pi^4} = \boxed{0.9855}.$$

(d)

$$\langle H \rangle = \sum |c_n|^2 E_n = \frac{96}{\pi^4} \frac{\pi^2 \hbar^2}{2ma^2} \underbrace{\left(\frac{1}{1} + \frac{1}{3^2} + \frac{1}{5^2} + \frac{1}{7^2} + \dots \right)}_{\pi^2/8} = \frac{48\hbar^2}{\pi^2 ma^2} \frac{\pi^2}{8} = \boxed{\frac{6\hbar^2}{ma^2}}.$$

Problem 2.8

(a)

$$\Psi(x, 0) = \begin{cases} A, & 0 < x < a/2; \\ 0, & \text{otherwise.} \end{cases} \quad 1 = A^2 \int_0^{a/2} dx = A^2(a/2) \Rightarrow A = \sqrt{\frac{2}{a}}.$$

(b) From Eq. 2.37,

$$c_1 = A \sqrt{\frac{2}{a}} \int_0^{a/2} \sin\left(\frac{\pi}{a}x\right) dx = \frac{2}{a} \left[-\frac{a}{\pi} \cos\left(\frac{\pi}{a}x\right) \right] \Big|_0^{a/2} = -\frac{2}{\pi} \left[\cos\left(\frac{\pi}{2}\right) - \cos 0 \right] = \frac{2}{\pi}.$$

$$P_1 = |c_1|^2 = \boxed{(2/\pi)^2 = 0.4053.}$$

Problem 2.9

$$\hat{H}\Psi(x, 0) = -\frac{\hbar^2}{2m} \frac{\partial^2}{\partial x^2} [Ax(a-x)] = -A \frac{\hbar^2}{2m} \frac{\partial}{\partial x} (a-2x) = A \frac{\hbar^2}{m}.$$

$$\begin{aligned} \int \Psi(x, 0)^* \hat{H}\Psi(x, 0) dx &= A^2 \frac{\hbar^2}{m} \int_0^a x(a-x) dx = A^2 \frac{\hbar^2}{m} \left(a \frac{x^2}{2} - \frac{x^3}{3} \right) \Big|_0^a \\ &= A^2 \frac{\hbar^2}{m} \left(\frac{a^3}{2} - \frac{a^3}{3} \right) = \frac{30}{a^5} \frac{\hbar^2}{m} \frac{a^3}{6} = \boxed{\frac{5\hbar^2}{ma^2}} \end{aligned}$$

(same as Example 2.3).

Problem 2.10


(a) Using Eqs. 2.47 and 2.59,

$$\begin{aligned} a_+ \psi_0 &= \frac{1}{\sqrt{2\hbar m\omega}} \left(-\hbar \frac{d}{dx} + m\omega x \right) \left(\frac{m\omega}{\pi\hbar} \right)^{1/4} e^{-\frac{m\omega}{2\hbar} x^2} \\ &= \frac{1}{\sqrt{2\hbar m\omega}} \left(\frac{m\omega}{\pi\hbar} \right)^{1/4} \left[-\hbar \left(-\frac{m\omega}{2\hbar} \right) 2x + m\omega x \right] e^{-\frac{m\omega}{2\hbar} x^2} = \frac{1}{\sqrt{2\hbar m\omega}} \left(\frac{m\omega}{\pi\hbar} \right)^{1/4} 2m\omega x e^{-\frac{m\omega}{2\hbar} x^2}. \\ (a_+)^2 \psi_0 &= \frac{1}{2\hbar m\omega} \left(\frac{m\omega}{\pi\hbar} \right)^{1/4} 2m\omega \left(-\hbar \frac{d}{dx} + m\omega x \right) x e^{-\frac{m\omega}{2\hbar} x^2} \\ &= \frac{1}{\hbar} \left(\frac{m\omega}{\pi\hbar} \right)^{1/4} \left[-\hbar \left(1 - x \frac{m\omega}{2\hbar} 2x \right) + m\omega x^2 \right] e^{-\frac{m\omega}{2\hbar} x^2} = \left(\frac{m\omega}{\pi\hbar} \right)^{1/4} \left(\frac{2m\omega}{\hbar} x^2 - 1 \right) e^{-\frac{m\omega}{2\hbar} x^2}. \end{aligned}$$

Therefore, from Eq. 2.67,

$$\psi_2 = \frac{1}{\sqrt{2}} (a_+)^2 \psi_0 = \boxed{\frac{1}{\sqrt{2}} \left(\frac{m\omega}{\pi\hbar} \right)^{1/4} \left(\frac{2m\omega}{\hbar} x^2 - 1 \right) e^{-\frac{m\omega}{2\hbar} x^2}.$$

(b)



(c) Since ψ_0 and ψ_2 are even, whereas ψ_1 is odd, $\int \psi_0^* \psi_1 dx$ and $\int \psi_2^* \psi_1 dx$ vanish automatically. The only one we need to check is $\int \psi_2^* \psi_0 dx$:

$$\begin{aligned} \int \psi_2^* \psi_0 dx &= \frac{1}{\sqrt{2}} \sqrt{\frac{m\omega}{\pi\hbar}} \int_{-\infty}^{\infty} \left(\frac{2m\omega}{\hbar} x^2 - 1 \right) e^{-\frac{m\omega}{\hbar} x^2} dx \\ &= -\sqrt{\frac{m\omega}{2\pi\hbar}} \left(\int_{-\infty}^{\infty} e^{-\frac{m\omega}{\hbar} x^2} dx - \frac{2m\omega}{\hbar} \int_{-\infty}^{\infty} x^2 e^{-\frac{m\omega}{\hbar} x^2} dx \right) \\ &= -\sqrt{\frac{m\omega}{2\pi\hbar}} \left(\sqrt{\frac{\pi\hbar}{m\omega}} - \frac{2m\omega}{\hbar} \frac{\hbar}{2m\omega} \sqrt{\frac{\pi\hbar}{m\omega}} \right) = 0. \quad \checkmark \end{aligned}$$

Problem 2.11

(a) Note that ψ_0 is even, and ψ_1 is odd. In either case $|\psi|^2$ is even, so $\langle x \rangle = \int x |\psi|^2 dx = \boxed{0}$. Therefore $\langle p \rangle = m d\langle x \rangle / dt = \boxed{0}$. (These results hold for *any* stationary state of the harmonic oscillator.)

From Eqs. 2.59 and 2.62, $\psi_0 = \alpha e^{-\xi^2/2}$, $\psi_1 = \sqrt{2}\alpha\xi e^{-\xi^2/2}$. So

$n = 0$:

$$\langle x^2 \rangle = \alpha^2 \int_{-\infty}^{\infty} x^2 e^{-\xi^2} dx = \alpha^2 \left(\frac{\hbar}{m\omega} \right)^{3/2} \int_{-\infty}^{\infty} \xi^2 e^{-\xi^2} d\xi = \frac{1}{\sqrt{\pi}} \left(\frac{\hbar}{m\omega} \right) \frac{\sqrt{\pi}}{2} = \boxed{\frac{\hbar}{2m\omega}}.$$

$$\begin{aligned} \langle p^2 \rangle &= \int \psi_0 \left(\frac{\hbar}{i} \frac{d}{dx} \right)^2 \psi_0 dx = -\hbar^2 \alpha^2 \sqrt{\frac{m\omega}{\hbar}} \int_{-\infty}^{\infty} e^{-\xi^2/2} \left(\frac{d^2}{d\xi^2} e^{-\xi^2/2} \right) d\xi \\ &= -\frac{m\hbar\omega}{\sqrt{\pi}} \int_{-\infty}^{\infty} (\xi^2 - 1) e^{-\xi^2} d\xi = -\frac{m\hbar\omega}{\sqrt{\pi}} \left(\frac{\sqrt{\pi}}{2} - \sqrt{\pi} \right) = \boxed{\frac{m\hbar\omega}{2}}. \end{aligned}$$

$n = 1$:

$$\langle x^2 \rangle = 2\alpha^2 \int_{-\infty}^{\infty} x^2 \xi^2 e^{-\xi^2} dx = 2\alpha^2 \left(\frac{\hbar}{m\omega} \right)^{3/2} \int_{-\infty}^{\infty} \xi^4 e^{-\xi^2} d\xi = \frac{2\hbar}{\sqrt{\pi}m\omega} \frac{3\sqrt{\pi}}{4} = \boxed{\frac{3\hbar}{2m\omega}}.$$

$$\begin{aligned} \langle p^2 \rangle &= -\hbar^2 2\alpha^2 \sqrt{\frac{m\omega}{\hbar}} \int_{-\infty}^{\infty} \xi e^{-\xi^2/2} \left[\frac{d^2}{d\xi^2} (\xi e^{-\xi^2/2}) \right] d\xi \\ &= -\frac{2m\omega\hbar}{\sqrt{\pi}} \int_{-\infty}^{\infty} (\xi^4 - 3\xi^2) e^{-\xi^2} d\xi = -\frac{2m\omega\hbar}{\sqrt{\pi}} \left(\frac{3}{4}\sqrt{\pi} - 3\frac{\sqrt{\pi}}{2} \right) = \boxed{\frac{3m\hbar\omega}{2}}. \end{aligned}$$

(b) $\underline{n = 0}$:

$$\sigma_x = \sqrt{\langle x^2 \rangle - \langle x \rangle^2} = \sqrt{\frac{\hbar}{2m\omega}}; \quad \sigma_p = \sqrt{\langle p^2 \rangle - \langle p \rangle^2} = \sqrt{\frac{m\hbar\omega}{2}};$$

$$\sigma_x \sigma_p = \sqrt{\frac{\hbar}{2m\omega}} \sqrt{\frac{m\omega\hbar}{2}} = \frac{\hbar}{2}. \quad (\text{Right at the uncertainty limit.}) \checkmark$$

$\underline{n = 1}$:

$$\sigma_x = \sqrt{\frac{3\hbar}{2m\omega}}; \quad \sigma_p = \sqrt{\frac{3m\hbar\omega}{2}}; \quad \sigma_x \sigma_p = 3\frac{\hbar}{2} > \frac{\hbar}{2}. \checkmark$$

(c)

$$\langle T \rangle = \frac{1}{2m} \langle p^2 \rangle = \begin{cases} \frac{1}{4}\hbar\omega & (n=0) \\ \frac{3}{4}\hbar\omega & (n=1) \end{cases}; \quad \langle V \rangle = \frac{1}{2}m\omega^2 \langle x^2 \rangle = \begin{cases} \frac{1}{4}\hbar\omega & (n=0) \\ \frac{3}{4}\hbar\omega & (n=1) \end{cases}.$$

$$\langle T \rangle + \langle V \rangle = \langle H \rangle = \begin{cases} \frac{1}{2}\hbar\omega & (n=0) = E_0 \\ \frac{3}{2}\hbar\omega & (n=1) = E_1 \end{cases}, \text{ as expected.}$$

Problem 2.12

From Eq. 2.69,

$$x = \sqrt{\frac{\hbar}{2m\omega}}(a_+ + a_-), \quad p = i\sqrt{\frac{\hbar m\omega}{2}}(a_+ - a_-),$$

so

$$\langle x \rangle = \sqrt{\frac{\hbar}{2m\omega}} \int \psi_n^*(a_+ + a_-)\psi_n dx.$$

But (Eq. 2.66)

$$a_+\psi_n = \sqrt{n+1}\psi_{n+1}, \quad a_-\psi_n = \sqrt{n}\psi_{n-1}.$$

So

$$\langle x \rangle = \sqrt{\frac{\hbar}{2m\omega}} \left[\sqrt{n+1} \int \psi_n^* \psi_{n+1} dx + \sqrt{n} \int \psi_n^* \psi_{n-1} dx \right] = \boxed{0} \text{ (by orthogonality).}$$

$$\langle p \rangle = m \frac{d\langle x \rangle}{dt} = \boxed{0}. \quad x^2 = \frac{\hbar}{2m\omega}(a_+ + a_-)^2 = \frac{\hbar}{2m\omega}(a_+^2 + a_+a_- + a_-a_+ + a_-^2).$$

$$\langle x^2 \rangle = \frac{\hbar}{2m\omega} \int \psi_n^*(a_+^2 + a_+a_- + a_-a_+ + a_-^2)\psi_n. \quad \text{But}$$

$$\begin{cases} a_+^2 \psi_n &= a_+ (\sqrt{n+1}\psi_{n+1}) = \sqrt{n+1}\sqrt{n+2}\psi_{n+2} = \sqrt{(n+1)(n+2)}\psi_{n+2}. \\ a_+a_- \psi_n &= a_+ (\sqrt{n}\psi_{n-1}) = \sqrt{n}\sqrt{n}\psi_n = n\psi_n. \\ a_-a_+ \psi_n &= a_- (\sqrt{n+1}\psi_{n+1}) = \sqrt{n+1}\sqrt{n+1}\psi_n = (n+1)\psi_n. \\ a_-^2 \psi_n &= a_- (\sqrt{n}\psi_{n-1}) = \sqrt{n}\sqrt{n-1}\psi_{n-2} = \sqrt{(n-1)n}\psi_{n-2}. \end{cases}$$

So

$$\langle x^2 \rangle = \frac{\hbar}{2m\omega} \left[0 + n \int |\psi_n|^2 dx + (n+1) \int |\psi_n|^2 dx + 0 \right] = \frac{\hbar}{2m\omega}(2n+1) = \boxed{\left(n + \frac{1}{2}\right) \frac{\hbar}{m\omega}}.$$

$$p^2 = -\frac{\hbar m \omega}{2} (a_+ - a_-)^2 = -\frac{\hbar m \omega}{2} (a_+^2 - a_+ a_- - a_- a_+ + a_-^2) \Rightarrow$$

$$\langle p^2 \rangle = -\frac{\hbar m \omega}{2} [0 - n - (n+1) + 0] = \frac{\hbar m \omega}{2} (2n+1) = \boxed{\left(n + \frac{1}{2}\right) m \hbar \omega}.$$

$$\langle T \rangle = \langle p^2 / 2m \rangle = \boxed{\frac{1}{2} \left(n + \frac{1}{2}\right) \hbar \omega}.$$

$$\sigma_x = \sqrt{\langle x^2 \rangle - \langle x \rangle^2} = \sqrt{n + \frac{1}{2}} \sqrt{\frac{\hbar}{m \omega}}; \quad \sigma_p = \sqrt{\langle p^2 \rangle - \langle p \rangle^2} = \sqrt{n + \frac{1}{2}} \sqrt{m \hbar \omega}; \quad \sigma_x \sigma_p = \left(n + \frac{1}{2}\right) \hbar \geq \frac{\hbar}{2}. \quad \checkmark$$

Problem 2.13

(a)

$$\begin{aligned} 1 &= \int |\Psi(x, 0)|^2 dx = |A|^2 \int (9|\psi_0|^2 + 12\psi_0^* \psi_1 + 12\psi_1^* \psi_0 + 16|\psi_1|^2) dx \\ &= |A|^2 (9 + 0 + 0 + 16) = 25|A|^2 \Rightarrow \boxed{A = 1/5}. \end{aligned}$$

(b)

$$\Psi(x, t) = \frac{1}{5} \left[3\psi_0(x) e^{-iE_0 t/\hbar} + 4\psi_1(x) e^{-iE_1 t/\hbar} \right] = \boxed{\frac{1}{5} \left[3\psi_0(x) e^{-i\omega t/2} + 4\psi_1(x) e^{-3i\omega t/2} \right]}.$$

(Here ψ_0 and ψ_1 are given by Eqs. 2.59 and 2.62; E_0 and E_1 by Eq. 2.61.)

$$\begin{aligned} |\Psi(x, t)|^2 &= \frac{1}{25} \left[9\psi_0^2 + 12\psi_0\psi_1 e^{i\omega t/2} e^{-3i\omega t/2} + 12\psi_0\psi_1 e^{-i\omega t/2} e^{3i\omega t/2} + 16\psi_1^2 \right] \\ &= \boxed{\frac{1}{25} \left[9\psi_0^2 + 16\psi_1^2 + 24\psi_0\psi_1 \cos(\omega t) \right]}. \end{aligned}$$

(With ψ_2 in place of ψ_1 the frequency would be $(E_2 - E_0)/\hbar = [(5/2)\hbar\omega - (1/2)\hbar\omega]/\hbar = 2\omega$.)

(c)

$$\langle x \rangle = \frac{1}{25} \left[9 \int x\psi_0^2 dx + 16 \int x\psi_1^2 dx + 24 \cos(\omega t) \int x\psi_0\psi_1 dx \right].$$

But $\int x\psi_0^2 dx = \int x\psi_1^2 dx = 0$ (see Problem 2.11 or 2.12), while

$$\begin{aligned} \int x\psi_0\psi_1 dx &= \sqrt{\frac{m\omega}{\pi\hbar}} \sqrt{\frac{2m\omega}{\hbar}} \int x e^{-\frac{m\omega}{2\hbar}x^2} x e^{-\frac{m\omega}{2\hbar}x^2} dx = \sqrt{\frac{2}{\pi}} \left(\frac{m\omega}{\hbar}\right) \int_{-\infty}^{\infty} x^2 e^{-\frac{m\omega}{\hbar}x^2} dx \\ &= \sqrt{\frac{2}{\pi}} \left(\frac{m\omega}{\hbar}\right) 2\sqrt{\pi} 2 \left(\frac{1}{2} \sqrt{\frac{\hbar}{m\omega}}\right)^3 = \sqrt{\frac{\hbar}{2m\omega}}. \end{aligned}$$

So

$$\langle x \rangle = \boxed{\frac{24}{25} \sqrt{\frac{\hbar}{2m\omega}} \cos(\omega t)}; \quad \langle p \rangle = m \frac{d}{dt} \langle x \rangle = \boxed{-\frac{24}{25} \sqrt{\frac{m\omega\hbar}{2}} \sin(\omega t)}.$$

Ehrenfest's theorem says $d\langle p \rangle/dt = -\langle \partial V/\partial x \rangle$. Here

$$\frac{d\langle p \rangle}{dt} = -\frac{24}{25} \sqrt{\frac{m\omega\hbar}{2}} \omega \cos(\omega t), \quad V = \frac{1}{2} m\omega^2 x^2 \Rightarrow \frac{\partial V}{\partial x} = m\omega^2 x,$$

so

$$-\langle \frac{\partial V}{\partial x} \rangle = -m\omega^2 \langle x \rangle = -m\omega^2 \frac{24}{25} \sqrt{\frac{\hbar}{2m\omega}} \cos(\omega t) = -\frac{24}{25} \sqrt{\frac{\hbar m\omega}{2}} \omega \cos(\omega t),$$

so Ehrenfest's theorem holds.

(d) You could get $E_0 = \frac{1}{2}\hbar\omega$, with probability $|c_0|^2 = 9/25$, or $E_1 = \frac{3}{2}\hbar\omega$, with probability $|c_1|^2 = 16/25$.

Problem 2.14

The new allowed energies are $E'_n = (n + \frac{1}{2})\hbar\omega' = 2(n + \frac{1}{2})\hbar\omega = \hbar\omega, 3\hbar\omega, 5\hbar\omega, \dots$. So the probability of getting $\frac{1}{2}\hbar\omega$ is **zero**. The probability of getting $\hbar\omega$ (the new ground state energy) is $P_0 = |c_0|^2$, where $c_0 = \int \Psi(x, 0)\psi'_0 dx$, with

$$\Psi(x, 0) = \psi_0(x) = \left(\frac{m\omega}{\pi\hbar}\right)^{1/4} e^{-\frac{m\omega}{2\hbar}x^2}, \quad \psi_0(x)' = \left(\frac{m2\omega}{\pi\hbar}\right)^{1/4} e^{-\frac{m2\omega}{2\hbar}x^2}.$$

So

$$c_0 = 2^{1/4} \sqrt{\frac{m\omega}{\pi\hbar}} \int_{-\infty}^{\infty} e^{-\frac{3m\omega}{2\hbar}x^2} dx = 2^{1/4} \sqrt{\frac{m\omega}{\pi\hbar}} 2\sqrt{\pi} \left(\frac{1}{2} \sqrt{\frac{2\hbar}{3m\omega}}\right) = 2^{1/4} \sqrt{\frac{2}{3}}.$$

Therefore

$$P_0 = \frac{2}{3}\sqrt{2} = 0.9428.$$

Problem 2.15

$$\psi_0 = \left(\frac{m\omega}{\pi\hbar}\right)^{1/4} e^{-\xi^2/2}, \text{ so } P = 2\sqrt{\frac{m\omega}{\pi\hbar}} \int_{x_0}^{\infty} e^{-\xi^2} dx = 2\sqrt{\frac{m\omega}{\pi\hbar}} \sqrt{\frac{\hbar}{m\omega}} \int_{\xi_0}^{\infty} e^{-\xi^2} d\xi.$$

Classically allowed region extends out to: $\frac{1}{2}m\omega^2 x_0^2 = E_0 = \frac{1}{2}\hbar\omega$, or $x_0 = \sqrt{\frac{\hbar}{m\omega}}$, so $\xi_0 = 1$.

$$P = \frac{2}{\sqrt{\pi}} \int_1^{\infty} e^{-\xi^2} d\xi = 2(1 - F(\sqrt{2})) \text{ (in notation of CRC Table)} = \boxed{0.157}.$$

Problem 2.16

$n=5$: $j=1 \Rightarrow a_3 = \frac{-2(5-1)}{(1+1)(1+2)}a_1 = -\frac{4}{3}a_1$; $j=3 \Rightarrow a_5 = \frac{-2(5-3)}{(3+1)(3+2)}a_3 = -\frac{1}{5}a_3 = \frac{4}{15}a_1$; $j=5 \Rightarrow a_7 = 0$. So $H_5(\xi) = a_1\xi - \frac{4}{3}a_1\xi^3 + \frac{4}{15}a_1\xi^5 = \frac{a_1}{15}(15\xi - 20\xi^3 + 4\xi^5)$. By convention the coefficient of ξ^5 is 2^5 , so $a_1 = 15 \cdot 8$, and $H_5(\xi) = 120\xi - 160\xi^3 + 32\xi^5$ (which agrees with Table 2.1).

$n=6$: $j=0 \Rightarrow a_2 = \frac{-2(6-0)}{(0+1)(0+2)}a_0 = -6a_0$; $j=2 \Rightarrow a_4 = \frac{-2(6-2)}{(2+1)(2+2)}a_2 = -\frac{2}{3}a_2 = 4a_0$; $j=4 \Rightarrow a_6 = \frac{-2(6-4)}{(4+1)(4+2)}a_4 = -\frac{2}{15}a_4 = -\frac{8}{15}a_0$; $j=6 \Rightarrow a_8 = 0$. So $H_6(\xi) = a_0 - 6a_0\xi^2 + 4a_0\xi^4 - \frac{8}{15}a_0\xi^6$. The coefficient of ξ^6 is 2^6 , so $2^6 = -\frac{8}{15}a_0 \Rightarrow a_0 = -15 \cdot 8 = -120$. $H_6(\xi) = -120 + 720\xi^2 - 480\xi^4 + 64\xi^6$.

Problem 2.17

(a)

$$\begin{aligned}\frac{d}{d\xi}(e^{-\xi^2}) &= -2\xi e^{-\xi^2}; \quad \left(\frac{d}{d\xi}\right)^2 e^{-\xi^2} = \frac{d}{d\xi}(-2\xi e^{-\xi^2}) = (-2 + 4\xi^2)e^{-\xi^2}; \\ \left(\frac{d}{d\xi}\right)^3 e^{-\xi^2} &= \frac{d}{d\xi} \left[(-2 + 4\xi^2)e^{-\xi^2} \right] = \left[8\xi + (-2 + 4\xi^2)(-2\xi) \right] e^{-\xi^2} = (12\xi - 8\xi^3)e^{-\xi^2}; \\ \left(\frac{d}{d\xi}\right)^4 e^{-\xi^2} &= \frac{d}{d\xi} \left[(12\xi - 8\xi^3)e^{-\xi^2} \right] = \left[12 - 24\xi^2 + (12\xi - 8\xi^3)(-2\xi) \right] e^{-\xi^2} = (12 - 48\xi^2 + 16\xi^4)e^{-\xi^2}. \\ H_3(\xi) &= -e^{\xi^2} \left(\frac{d}{d\xi}\right)^3 e^{-\xi^2} = \boxed{-12\xi + 8\xi^3}; \quad H_4(\xi) = e^{\xi^2} \left(\frac{d}{d\xi}\right)^4 e^{-\xi^2} = \boxed{12 - 48\xi^2 + 16\xi^4}.\end{aligned}$$

(b)

$$\begin{aligned}H_5 &= 2\xi H_4 - 8H_3 = 2\xi(12 - 48\xi^2 + 16\xi^4) - 8(-12\xi + 8\xi^3) = \boxed{120\xi - 160\xi^3 + 32\xi^5}. \\ H_6 &= 2\xi H_5 - 10H_4 = 2\xi(120\xi - 160\xi^3 + 32\xi^5) - 10(12 - 48\xi^2 + 16\xi^4) = \boxed{-120 + 720\xi^2 - 480\xi^4 + 64\xi^6}.\end{aligned}$$

(c)

$$\begin{aligned}\frac{dH_5}{d\xi} &= 120 - 480\xi^2 + 160\xi^4 = 10(12 - 48\xi^2 + 16\xi^4) = (2)(5)H_4. \quad \checkmark \\ \frac{dH_6}{d\xi} &= 1440\xi - 1920\xi^3 + 384\xi^5 = 12(120\xi - 160\xi^3 + 32\xi^5) = (2)(6)H_5. \quad \checkmark\end{aligned}$$

(d)

$$\begin{aligned}\frac{d}{dz}(e^{-z^2+2z\xi}) &= (-2z + 2\xi)e^{-z^2+2z\xi}; \quad \text{setting } z = 0, \quad \boxed{H_1(\xi) = 2\xi}. \\ \left(\frac{d}{dz}\right)^2 (e^{-z^2+2z\xi}) &= \frac{d}{dz} \left[(-2z + 2\xi)e^{-z^2+2z\xi} \right] \\ &= \left[-2 + (-2z + 2\xi)^2 \right] e^{-z^2+2z\xi}; \quad \text{setting } z = 0, \quad \boxed{H_2(\xi) = -2 + 4\xi^2}. \\ \left(\frac{d}{dz}\right)^3 (e^{-z^2+2z\xi}) &= \frac{d}{dz} \left\{ \left[-2 + (-2z + 2\xi)^2 \right] e^{-z^2+2z\xi} \right\} \\ &= \left\{ 2(-2z + 2\xi)(-2) + \left[-2 + (-2z + 2\xi)^2 \right] (-2z + 2\xi) \right\} e^{-z^2+2z\xi}; \\ \text{setting } z = 0, \quad H_3(\xi) &= -8\xi + (-2 + 4\xi^2)(2\xi) = \boxed{-12\xi + 8\xi^3}.\end{aligned}$$

Problem 2.18

$$\begin{aligned} Ae^{ikx} + Be^{-ikx} &= A(\cos kx + i \sin kx) + B(\cos kx - i \sin kx) = (A + B) \cos kx + i(A - B) \sin kx \\ &= C \cos kx + D \sin kx, \text{ with } \boxed{C = A + B; D = i(A - B)}. \end{aligned}$$

$$\begin{aligned} C \cos kx + D \sin kx &= C \left(\frac{e^{ikx} + e^{-ikx}}{2} \right) + D \left(\frac{e^{ikx} - e^{-ikx}}{2i} \right) = \frac{1}{2}(C - iD)e^{ikx} + \frac{1}{2}(C + iD)e^{-ikx} \\ &= Ae^{ikx} + Be^{-ikx}, \text{ with } \boxed{A = \frac{1}{2}(C - iD); B = \frac{1}{2}(C + iD)}. \end{aligned}$$

Problem 2.19

Equation 2.94 says $\Psi = Ae^{i(kx - \frac{\hbar k^2}{2m}t)}$, so

$$\begin{aligned} J &= \frac{i\hbar}{2m} \left(\Psi \frac{\partial \Psi^*}{\partial x} - \Psi^* \frac{\partial \Psi}{\partial x} \right) = \frac{i\hbar}{2m} |A|^2 \left[e^{i(kx - \frac{\hbar k^2}{2m}t)} (-ik) e^{-i(kx - \frac{\hbar k^2}{2m}t)} - e^{-i(kx - \frac{\hbar k^2}{2m}t)} (ik) e^{i(kx - \frac{\hbar k^2}{2m}t)} \right] \\ &= \frac{i\hbar}{2m} |A|^2 (-2ik) = \boxed{\frac{\hbar k}{m} |A|^2}. \end{aligned}$$

It flows in the positive (x) direction (as you would expect).

Problem 2.20

(a)

$$\begin{aligned} f(x) &= b_0 + \sum_{n=1}^{\infty} \frac{a_n}{2i} (e^{in\pi x/a} - e^{-in\pi x/a}) + \sum_{n=1}^{\infty} \frac{b_n}{2} (e^{in\pi x/a} + e^{-in\pi x/a}) \\ &= b_0 + \sum_{n=1}^{\infty} \left(\frac{a_n}{2i} + \frac{b_n}{2} \right) e^{in\pi x/a} + \sum_{n=1}^{\infty} \left(-\frac{a_n}{2i} + \frac{b_n}{2} \right) e^{-in\pi x/a}. \end{aligned}$$

Let

$$\boxed{c_0 \equiv b_0; c_n = \frac{1}{2}(-ia_n + b_n), \text{ for } n = 1, 2, 3, \dots; c_n \equiv \frac{1}{2}(ia_{-n} + b_{-n}), \text{ for } n = -1, -2, -3, \dots}$$

$$\text{Then } f(x) = \sum_{n=-\infty}^{\infty} c_n e^{in\pi x/a}. \quad \text{QED}$$

(b)

$$\begin{aligned} \int_{-a}^a f(x) e^{-im\pi x/a} dx &= \sum_{n=-\infty}^{\infty} c_n \int_{-a}^a e^{i(n-m)\pi x/a} dx. \quad \text{But for } n \neq m, \\ \int_{-a}^a e^{i(n-m)\pi x/a} dx &= \frac{e^{i(n-m)\pi x/a}}{i(n-m)\pi/a} \Big|_{-a}^a = \frac{e^{i(n-m)\pi} - e^{-i(n-m)\pi}}{i(n-m)\pi/a} = \frac{(-1)^{n-m} - (-1)^{n-m}}{i(n-m)\pi/a} = 0, \end{aligned}$$

whereas for $n = m$,

$$\int_{-a}^a e^{i(n-m)\pi x/a} dx = \int_{-a}^a dx = 2a.$$

So all terms except $n = m$ are zero, and

$$\int_{-a}^a f(x) e^{-im\pi x/a} dx = 2ac_m, \text{ so } c_n = \frac{1}{2a} \int_{-a}^a f(x) e^{-in\pi x/a} dx. \quad \text{QED}$$

(c)

$$f(x) = \sum_{n=-\infty}^{\infty} \sqrt{\frac{\pi}{2}} \frac{1}{a} F(k) e^{ikx} = \frac{1}{\sqrt{2\pi}} \sum F(k) e^{ikx} \Delta k,$$

where $\Delta k \equiv \frac{\pi}{a}$ is the increment in k from n to $(n+1)$.

$$F(k) = \sqrt{\frac{2}{\pi}} \frac{1}{2a} \int_{-a}^a f(x) e^{-ikx} dx = \frac{1}{\sqrt{2\pi}} \int_{-a}^a f(x) e^{-ikx} dx.$$

(d) As $a \rightarrow \infty$, k becomes a continuous variable,

$$f(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} F(k) e^{ikx} dk; \quad F(k) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} f(x) e^{-ikx} dx.$$

Problem 2.21

(a)

$$1 = \int_{-\infty}^{\infty} |\Psi(x, 0)|^2 dx = 2|A|^2 \int_0^{\infty} e^{-2ax} dx = 2|A|^2 \left. \frac{e^{-2ax}}{-2a} \right|_0^{\infty} = \frac{|A|^2}{a} \Rightarrow A = \boxed{\sqrt{a}}.$$

(b)

$$\phi(k) = \frac{A}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{-a|x|} e^{-ikx} dx = \frac{A}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{-a|x|} (\cos kx - i \sin kx) dx.$$

The cosine integrand is even, and the sine is odd, so the latter vanishes and

$$\begin{aligned} \phi(k) &= 2 \frac{A}{\sqrt{2\pi}} \int_0^{\infty} e^{-ax} \cos kx dx = \frac{A}{\sqrt{2\pi}} \int_0^{\infty} e^{-ax} (e^{ikx} + e^{-ikx}) dx \\ &= \frac{A}{\sqrt{2\pi}} \int_0^{\infty} (e^{(ik-a)x} + e^{-(ik+a)x}) dx = \frac{A}{\sqrt{2\pi}} \left[\frac{e^{(ik-a)x}}{ik-a} + \frac{e^{-(ik+a)x}}{-(ik+a)} \right] \Big|_0^{\infty} \\ &= \frac{A}{\sqrt{2\pi}} \left(\frac{-1}{ik-a} + \frac{1}{ik+a} \right) = \frac{A}{\sqrt{2\pi}} \frac{-ik-a+ik-a}{-k^2-a^2} = \boxed{\sqrt{\frac{a}{2\pi}} \frac{2a}{k^2+a^2}}. \end{aligned}$$

(c)

$$\Psi(x, t) = \frac{1}{\sqrt{2\pi}} 2\sqrt{\frac{a^3}{2\pi}} \int_{-\infty}^{\infty} \frac{1}{k^2+a^2} e^{i(kx - \frac{\hbar k^2}{2m}t)} dk = \boxed{\frac{a^{3/2}}{\pi} \int_{-\infty}^{\infty} \frac{1}{k^2+a^2} e^{i(kx - \frac{\hbar k^2}{2m}t)} dk}.$$

- (d) For *large* a , $\Psi(x, 0)$ is a sharp narrow spike whereas $\phi(k) \cong \sqrt{2/\pi a}$ is broad and flat; position is well-defined but momentum is ill-defined. For *small* a , $\Psi(x, 0)$ is a broad and flat whereas $\phi(k) \cong (\sqrt{2a^3/\pi})/k^2$ is a sharp narrow spike; position is ill-defined but momentum is well-defined.

Problem 2.22

(a)

$$1 = |A|^2 \int_{-\infty}^{\infty} e^{-2ax^2} dx = |A|^2 \sqrt{\frac{\pi}{2a}}; \quad \boxed{A = \left(\frac{2a}{\pi}\right)^{1/4}}.$$

(b)

$$\int_{-\infty}^{\infty} e^{-(ax^2+bx)} dx = \int_{-\infty}^{\infty} e^{-y^2+(b^2/4a)} \frac{1}{\sqrt{a}} dy = \frac{1}{\sqrt{a}} e^{b^2/4a} \int_{-\infty}^{\infty} e^{-y^2} dy = \sqrt{\frac{\pi}{a}} e^{b^2/4a}.$$

$$\phi(k) = \frac{1}{\sqrt{2\pi}} A \int_{-\infty}^{\infty} e^{-ax^2} e^{-ikx} dx = \frac{1}{\sqrt{2\pi}} \left(\frac{2a}{\pi}\right)^{1/4} \sqrt{\frac{\pi}{a}} e^{-k^2/4a} = \frac{1}{(2\pi a)^{1/4}} e^{-k^2/4a}.$$


$$\begin{aligned} \Psi(x, t) &= \frac{1}{\sqrt{2\pi}} \frac{1}{(2\pi a)^{1/4}} \int_{-\infty}^{\infty} \underbrace{e^{-k^2/4a} e^{i(kx - \hbar k^2 t/2m)}}_{e^{-[(\frac{1}{4a} + i\hbar t/2m)k^2 - ixk]}} dk \\ &= \frac{1}{\sqrt{2\pi}(2\pi a)^{1/4}} \frac{\sqrt{\pi}}{\sqrt{\frac{1}{4a} + i\hbar t/2m}} e^{-x^2/4(\frac{1}{4a} + i\hbar t/2m)} = \boxed{\left(\frac{2a}{\pi}\right)^{1/4} \frac{e^{-ax^2/(1+2i\hbar at/m)}}{\sqrt{1+2i\hbar at/m}}}. \end{aligned}$$

(c)

Let $\theta \equiv 2\hbar at/m$. Then $|\Psi|^2 = \sqrt{\frac{2a}{\pi}} \frac{e^{-ax^2/(1+i\theta)} e^{-ax^2/(1-i\theta)}}{\sqrt{(1+i\theta)(1-i\theta)}}$. The exponent is

$$-\frac{ax^2}{(1+i\theta)} - \frac{ax^2}{(1-i\theta)} = -ax^2 \frac{(1-i\theta+1+i\theta)}{(1+i\theta)(1-i\theta)} = \frac{-2ax^2}{1+\theta^2}; \quad |\Psi|^2 = \sqrt{\frac{2a}{\pi}} \frac{e^{-2ax^2/(1+\theta^2)}}{\sqrt{1+\theta^2}}.$$

Or, with $w \equiv \sqrt{\frac{a}{1+\theta^2}}$, $|\Psi|^2 = \sqrt{\frac{2}{\pi}} w e^{-2w^2 x^2}$. As t increases, the graph of $|\Psi|^2$ flattens out and broadens.



(d)

$$\langle x \rangle = \int_{-\infty}^{\infty} x |\Psi|^2 dx = \boxed{0} \text{ (odd integrand); } \langle p \rangle = m \frac{d\langle x \rangle}{dt} = \boxed{0}.$$

$$\langle x^2 \rangle = \sqrt{\frac{2}{\pi}} w \int_{-\infty}^{\infty} x^2 e^{-2w^2 x^2} dx = \sqrt{\frac{2}{\pi}} w \frac{1}{4w^2} \sqrt{\frac{\pi}{2w^2}} = \boxed{\frac{1}{4w^2}}. \quad \langle p^2 \rangle = -\hbar^2 \int_{-\infty}^{\infty} \Psi^* \frac{\partial^2 \Psi}{\partial x^2} dx.$$

$$\text{Write } \Psi = B e^{-bx^2}, \text{ where } B \equiv \left(\frac{2a}{\pi}\right)^{1/4} \frac{1}{\sqrt{1+i\theta}} \text{ and } b \equiv \frac{a}{1+i\theta}.$$

$$\frac{\partial^2 \Psi}{\partial x^2} = B \frac{\partial}{\partial x} (-2bx e^{-bx^2}) = -2bB(1 - 2bx^2)e^{-bx^2}.$$

$$\Psi^* \frac{\partial^2 \Psi}{\partial x^2} = -2b|B|^2(1 - 2bx^2)e^{-(b+b^*)x^2}; \quad b + b^* = \frac{a}{1+i\theta} + \frac{a}{1-i\theta} = \frac{2a}{1+\theta^2} = 2w^2.$$

$$|B|^2 = \sqrt{\frac{2a}{\pi}} \frac{1}{\sqrt{1+\theta^2}} = \sqrt{\frac{2}{\pi}} w. \quad \text{So } \Psi^* \frac{\partial^2 \Psi}{\partial x^2} = -2b \sqrt{\frac{2}{\pi}} w (1 - 2bx^2) e^{-2w^2 x^2}.$$

$$\begin{aligned} \langle p^2 \rangle &= 2b\hbar^2 \sqrt{\frac{2}{\pi}} w \int_{-\infty}^{\infty} (1 - 2bx^2) e^{-2w^2 x^2} dx \\ &= 2b\hbar^2 \sqrt{\frac{2}{\pi}} w \left(\sqrt{\frac{\pi}{2w^2}} - 2b \frac{1}{4w^2} \sqrt{\frac{\pi}{2w^2}} \right) = 2b\hbar^2 \left(1 - \frac{b}{2w^2} \right). \end{aligned}$$

$$\text{But } 1 - \frac{b}{2w^2} = 1 - \left(\frac{a}{1+i\theta} \right) \left(\frac{1+\theta^2}{2a} \right) = 1 - \frac{(1-i\theta)}{2} = \frac{1+i\theta}{2} = \frac{a}{2b}, \text{ so}$$

$$\langle p^2 \rangle = 2b\hbar^2 \frac{a}{2b} = \boxed{\hbar^2 a}. \quad \boxed{\sigma_x = \frac{1}{2w}}; \quad \boxed{\sigma_p = \hbar \sqrt{a}}.$$

(e)

$$\sigma_x \sigma_p = \frac{1}{2w} \hbar \sqrt{a} = \frac{\hbar}{2} \sqrt{1+\theta^2} = \frac{\hbar}{2} \sqrt{1+(2\hbar a t/m)^2} \geq \frac{\hbar}{2}. \quad \checkmark$$

Closest at $\boxed{t=0}$, at which time it is right at the uncertainty limit.

Problem 2.23

(a)

$$(-2)^3 - 3(-2)^2 + 2(-2) - 1 = -8 - 12 - 4 - 1 = \boxed{-25}.$$

(b)

$$\cos(3\pi) + 2 = -1 + 2 = \boxed{1}.$$

(c)

$$\boxed{0} \text{ (} x=2 \text{ is outside the domain of integration).}$$

Problem 2.24

(a) Let $y \equiv cx$, so $dx = \frac{1}{c}dy$. $\begin{cases} \text{If } c > 0, y : -\infty \rightarrow \infty. \\ \text{If } c < 0, y : \infty \rightarrow -\infty. \end{cases}$

$$\int_{-\infty}^{\infty} f(x)\delta(cx)dx = \begin{cases} \frac{1}{c} \int_{-\infty}^{\infty} f(y/c)\delta(y)dy = \frac{1}{c}f(0) & (c > 0); \text{ or} \\ \frac{1}{c} \int_{\infty}^{-\infty} f(y/c)\delta(y)dy = -\frac{1}{c} \int_{-\infty}^{\infty} f(y/c)\delta(y)dy = -\frac{1}{c}f(0) & (c < 0). \end{cases}$$

In either case, $\int_{-\infty}^{\infty} f(x)\delta(cx)dx = \frac{1}{|c|}f(0) = \int_{-\infty}^{\infty} f(x)\frac{1}{|c|}\delta(x)dx$. So $\delta(cx) = \frac{1}{|c|}\delta(x)$. ✓

(b)

$$\begin{aligned} \int_{-\infty}^{\infty} f(x)\frac{d\theta}{dx}dx &= f\theta \Big|_{-\infty}^{\infty} - \int_{-\infty}^{\infty} \frac{df}{dx}\theta dx \quad (\text{integration by parts}) \\ &= f(\infty) - \int_0^{\infty} \frac{df}{dx}dx = f(\infty) - f(\infty) + f(0) = f(0) = \int_{-\infty}^{\infty} f(x)\delta(x)dx. \end{aligned}$$

So $d\theta/dx = \delta(x)$. ✓ [Makes sense: The θ function is constant (so derivative is zero) except at $x = 0$, where the derivative is infinite.]

Problem 2.25

$$\psi(x) = \frac{\sqrt{m\alpha}}{\hbar} e^{-m\alpha|x|/\hbar^2} = \frac{\sqrt{m\alpha}}{\hbar} \begin{cases} e^{-m\alpha x/\hbar^2}, & (x \geq 0), \\ e^{m\alpha x/\hbar^2}, & (x \leq 0). \end{cases}$$

$$\langle x \rangle = 0 \text{ (odd integrand).}$$

$$\langle x^2 \rangle = \int_{-\infty}^{\infty} x^2 |\psi|^2 dx = 2 \frac{m\alpha}{\hbar^2} \int_0^{\infty} x^2 e^{-2m\alpha x/\hbar^2} dx = \frac{2m\alpha}{\hbar^2} 2 \left(\frac{\hbar^2}{2m\alpha} \right)^3 = \frac{\hbar^4}{2m^2\alpha^2}; \quad \sigma_x = \frac{\hbar^2}{\sqrt{2}m\alpha}.$$

$$\frac{d\psi}{dx} = \frac{\sqrt{m\alpha}}{\hbar} \begin{cases} -\frac{m\alpha}{\hbar^2} e^{-m\alpha x/\hbar^2}, & (x \geq 0) \\ \frac{m\alpha}{\hbar^2} e^{m\alpha x/\hbar^2}, & (x \leq 0) \end{cases} = \left(\frac{\sqrt{m\alpha}}{\hbar} \right)^3 \left[-\theta(x)e^{-m\alpha x/\hbar^2} + \theta(-x)e^{m\alpha x/\hbar^2} \right].$$

$$\begin{aligned} \frac{d^2\psi}{dx^2} &= \left(\frac{\sqrt{m\alpha}}{\hbar} \right)^3 \left[-\delta(x)e^{-m\alpha x/\hbar^2} + \frac{m\alpha}{\hbar^2}\theta(x)e^{-m\alpha x/\hbar^2} - \delta(-x)e^{m\alpha x/\hbar^2} + \frac{m\alpha}{\hbar^2}\theta(-x)e^{m\alpha x/\hbar^2} \right] \\ &= \left(\frac{\sqrt{m\alpha}}{\hbar} \right)^3 \left[-2\delta(x) + \frac{m\alpha}{\hbar^2}e^{-m\alpha|x|/\hbar^2} \right]. \end{aligned}$$

In the last step I used the fact that $\delta(-x) = \delta(x)$ (Eq. 2.142), $f(x)\delta(x) = f(0)\delta(x)$ (Eq. 2.112), and $\theta(-x) + \theta(x) = 1$ (Eq. 2.143). Since $d\psi/dx$ is an odd function, $\langle p \rangle = 0$.

$$\begin{aligned} \langle p^2 \rangle &= -\hbar^2 \int_{-\infty}^{\infty} \psi \frac{d^2\psi}{dx^2} dx = -\hbar^2 \frac{\sqrt{m\alpha}}{\hbar} \left(\frac{\sqrt{m\alpha}}{\hbar} \right)^3 \int_{-\infty}^{\infty} e^{-m\alpha|x|/\hbar^2} \left[-2\delta(x) + \frac{m\alpha}{\hbar^2}e^{-m\alpha|x|/\hbar^2} \right] dx \\ &= \left(\frac{m\alpha}{\hbar} \right)^2 \left[2 - 2 \frac{m\alpha}{\hbar^2} \int_0^{\infty} e^{-2m\alpha x/\hbar^2} dx \right] = 2 \left(\frac{m\alpha}{\hbar} \right)^2 \left[1 - \frac{m\alpha}{\hbar^2} \frac{\hbar^2}{2m\alpha} \right] = \left(\frac{m\alpha}{\hbar} \right)^2. \end{aligned}$$

Evidently

$$\sigma_p = \frac{m\alpha}{\hbar}, \quad \text{so} \quad \sigma_x \sigma_p = \frac{\hbar^2}{\sqrt{2}m\alpha} \frac{m\alpha}{\hbar} = \sqrt{2} \frac{\hbar}{2} > \frac{\hbar}{2}. \quad \checkmark$$


Problem 2.26

Put $f(x) = \delta(x)$ into Eq. 2.102: $F(k) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \delta(x) e^{-ikx} dx = \boxed{\frac{1}{\sqrt{2\pi}}}$.

$\therefore f(x) = \delta(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi}} e^{ikx} dk = \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{ikx} dk$. QED

Problem 2.27

(a)



(b) From Problem 2.1(c) the solutions are even or odd. Look first for *even solutions*:

$$\psi(x) = \begin{cases} Ae^{-\kappa x} & (x > a), \\ B(e^{\kappa x} + e^{-\kappa x}) & (-a < x < a), \\ Ae^{\kappa x} & (x < -a). \end{cases}$$

Continuity at a : $Ae^{-\kappa a} = B(e^{\kappa a} + e^{-\kappa a})$, or $A = B(e^{2\kappa a} + 1)$.


Discontinuous derivative at a , $\Delta \frac{d\psi}{dx} = -\frac{2m\alpha}{\hbar^2} \psi(a)$:

$$-\kappa Ae^{-\kappa a} - B(\kappa e^{\kappa a} - \kappa e^{-\kappa a}) = -\frac{2m\alpha}{\hbar^2} Ae^{-\kappa a} \Rightarrow A + B(e^{2\kappa a} - 1) = \frac{2m\alpha}{\hbar^2 \kappa} A; \text{ or}$$

$$B(e^{2\kappa a} - 1) = A \left(\frac{2m\alpha}{\hbar^2 \kappa} - 1 \right) = B(e^{2\kappa a} + 1) \left(\frac{2m\alpha}{\hbar^2 \kappa} - 1 \right) \Rightarrow e^{2\kappa a} - 1 = e^{2\kappa a} \left(\frac{2m\alpha}{\hbar^2 \kappa} - 1 \right) + \frac{2m\alpha}{\hbar^2 \kappa} - 1.$$

$$1 = \frac{2m\alpha}{\hbar^2 \kappa} - 1 + \frac{2m\alpha}{\hbar^2 \kappa} e^{-2\kappa a}; \quad \frac{\hbar^2 \kappa}{m\alpha} = 1 + e^{-2\kappa a}, \text{ or } \boxed{e^{-2\kappa a} = \frac{\hbar^2 \kappa}{m\alpha} - 1}.$$

This is a transcendental equation for κ (and hence for E). I'll solve it graphically: Let $z \equiv 2\kappa a$, $c \equiv \frac{\hbar^2}{2am\alpha}$, so $e^{-z} = cz - 1$. Plot both sides and look for intersections:



From the graph, noting that c and z are both positive, we see that there is one (and only one) solution (for even ψ). If $\alpha = \frac{\hbar^2}{2ma}$, so $c = 1$, the calculator gives $z = 1.278$, so $\kappa^2 = -\frac{2mE}{\hbar^2} = \frac{z^2}{(2a)^2} \Rightarrow E = -\frac{(1.278)^2}{8} \left(\frac{\hbar^2}{ma^2} \right) = -0.204 \left(\frac{\hbar^2}{ma^2} \right)$.

Now look for *odd solutions*:


$$\psi(x) = \begin{cases} Ae^{-\kappa x} & (x > a), \\ B(e^{\kappa x} - e^{-\kappa x}) & (-a < x < a), \\ -Ae^{\kappa x} & (x < -a). \end{cases}$$

Continuity at a : $Ae^{-\kappa a} = B(e^{\kappa a} - e^{-\kappa a})$, or $A = B(e^{2\kappa a} - 1)$.

Discontinuity in ψ' : $-\kappa Ae^{-\kappa a} - B(\kappa e^{\kappa a} + \kappa e^{-\kappa a}) = -\frac{2m\alpha}{\hbar^2} Ae^{-\kappa a} \Rightarrow B(e^{2\kappa a} + 1) = A \left(\frac{2m\alpha}{\hbar^2 \kappa} - 1 \right)$,


$$e^{2\kappa a} + 1 = (e^{2\kappa a} - 1) \left(\frac{2m\alpha}{\hbar^2 \kappa} - 1 \right) = e^{2\kappa a} \left(\frac{2m\alpha}{\hbar^2 \kappa} - 1 \right) - \frac{2m\alpha}{\hbar^2 \kappa} + 1,$$

$$1 = \frac{2m\alpha}{\hbar^2 \kappa} - 1 - \frac{2m\alpha}{\hbar^2 \kappa} e^{-2\kappa a}; \quad \frac{\hbar^2 \kappa}{m\alpha} = 1 - e^{-2\kappa a}, \quad \boxed{e^{-2\kappa a} = 1 - \frac{\hbar^2 \kappa}{m\alpha}}, \text{ or } e^{-z} = 1 - cz.$$



This time there may or may not be a solution. Both graphs have their y -intercepts at 1, but if c is too large (α too small), there may be no intersection (solid line), whereas if c is smaller (dashed line) there will be. (Note that $z = 0 \Rightarrow \kappa = 0$ is *not* a solution, since ψ is then non-normalizable.) The slope of e^{-z} (at $z = 0$) is -1 ; the slope of $(1 - cz)$ is $-c$. So there is an *odd* solution $\Leftrightarrow c < 1$, or $\alpha > \hbar^2/2ma$.

Conclusion: One bound state if $\alpha \leq \hbar^2/2ma$; two if $\alpha > \hbar^2/2ma$.



$$\alpha = \frac{\hbar^2}{ma} \Rightarrow c = \frac{1}{2} \cdot \begin{cases} \text{Even: } e^{-z} = \frac{1}{2}z - 1 \Rightarrow z = 2.21772, \\ \text{Odd: } e^{-z} = 1 - \frac{1}{2}z \Rightarrow z = 1.59362. \end{cases}$$

$$\boxed{E = -0.615(\hbar^2/ma^2); \quad E = -0.317(\hbar^2/ma^2).}$$

$$\alpha = \frac{\hbar^2}{4ma} \Rightarrow c = 2. \text{ Only even: } e^{-z} = 2z - 1 \Rightarrow z = 0.738835; \quad \boxed{E = -0.0682(\hbar^2/ma^2).}$$

Problem 2.28

$$\psi = \begin{cases} Ae^{ikx} + Be^{-ikx} & (x < -a) \\ Ce^{ikx} + De^{-ikx} & (-a < x < a) \\ Fe^{ikx} & (x > a) \end{cases}. \quad \text{Impose boundary conditions:}$$

- (1) Continuity at $-a$: $Ae^{-ika} + Be^{ika} = Ce^{-ika} + De^{ika} \Rightarrow \beta A + B = \beta C + D$, where $\beta \equiv e^{-2ika}$.
 (2) Continuity at $+a$: $Ce^{ika} + De^{-ika} = Fe^{ika} \Rightarrow F = C + \beta D$.
 (3) Discontinuity in ψ' at $-a$: $ik(Ce^{-ika} - De^{ika}) - ik(Ae^{-ika} - Be^{ika}) = -\frac{2m\alpha}{\hbar^2}(Ae^{-ika} + Be^{ika})$
 $\Rightarrow \beta C - D = \beta(\gamma + 1)A + B(\gamma - 1)$, where $\gamma \equiv i2m\alpha/\hbar^2 k$.
 (4) Discontinuity in ψ' at $+a$: $ikFe^{ika} - ik(Ce^{ika} - De^{-ika}) = -\frac{2m\alpha}{\hbar^2}(Fe^{ika})$
 $\Rightarrow C - \beta D = (1 - \gamma)F$.

To solve for C and D , $\begin{cases} \text{add (2) and (4): } 2C = F + (1 - \gamma)F \Rightarrow 2C = (2 - \gamma)F. \\ \text{subtract (2) and (4): } 2\beta D = F - (1 - \gamma)F \Rightarrow 2D = (\gamma/\beta)F. \end{cases}$

$\begin{cases} \text{add (1) and (3): } 2\beta C = \beta A + B + \beta(\gamma + 1)A + B(\gamma - 1) \Rightarrow 2C = (\gamma + 2)A + (\gamma/\beta)B. \\ \text{subtract (1) and (3): } 2D = \beta A + B - \beta(\gamma + 1)A - B(\gamma - 1) \Rightarrow 2D = -\gamma\beta A + (2 - \gamma)B. \end{cases}$

$\begin{cases} \text{Equate the two expressions for } 2C: (2 - \gamma)F = (\gamma + 2)A + (\gamma/\beta)B. \\ \text{Equate the two expressions for } 2D: (\gamma/\beta)F = -\gamma\beta A + (2 - \gamma)B. \end{cases}$

Solve these for F and B , in terms of A . Multiply the first by $\beta(2 - \gamma)$, the second by γ , and subtract:

$$[\beta(2 - \gamma)^2 F = \beta(4 - \gamma^2)A + \gamma(2 - \gamma)B]; \quad [(\gamma^2/\beta)F = -\beta\gamma^2 A + \gamma(2 - \gamma)B].$$

$$\Rightarrow [\beta(2 - \gamma)^2 - \gamma^2/\beta] F = \beta[4 - \gamma^2 + \gamma^2] A = 4\beta A \Rightarrow \frac{F}{A} = \frac{4}{(2 - \gamma)^2 - \gamma^2/\beta^2}.$$

Let $g \equiv i/\gamma = \frac{\hbar^2 k}{2m\alpha}$; $\phi \equiv 4ka$, so $\gamma = \frac{i}{g}$, $\beta^2 = e^{-i\phi}$. Then: $\frac{F}{A} = \frac{4g^2}{(2g - i)^2 + e^{i\phi}}.$

Denominator: $4g^2 - 4ig - 1 + \cos \phi + i \sin \phi = (4g^2 - 1 + \cos \phi) + i(\sin \phi - 4g).$

$$\begin{aligned} |\text{Denominator}|^2 &= (4g^2 - 1 + \cos \phi)^2 + (\sin \phi - 4g)^2 \\ &= 16g^4 + 1 + \cos^2 \phi - 8g^2 - 2\cos \phi + 8g^2 \cos \phi + \sin^2 \phi - 8g \sin \phi + 16g^2 \\ &= 16g^4 + 8g^2 + 2 + (8g^2 - 2)\cos \phi - 8g \sin \phi. \end{aligned}$$

$$T = \left| \frac{F}{A} \right|^2 = \frac{8g^4}{(8g^4 + 4g^2 + 1) + (4g^2 - 1)\cos \phi - 4g \sin \phi}, \quad \text{where } g \equiv \frac{\hbar^2 k}{2m\alpha} \text{ and } \phi \equiv 4ka.$$

Problem 2.29

In place of Eq. 2.151, we have: $\psi(x) = \begin{cases} Fe^{-\kappa x} & (x > a) \\ D \sin(lx) & (0 < x < a) \\ -\psi(-x) & (x < 0) \end{cases}.$


Continuity of ψ : $Fe^{-\kappa a} = D \sin(la)$; continuity of ψ' : $-F\kappa e^{-\kappa a} = Dl \cos(la)$.

Divide: $-\kappa = l \cot(la)$, or $-\kappa a = la \cot(la) \Rightarrow \sqrt{z_0^2 - z^2} = -z \cot z$, or $-\cot z = \sqrt{(z_0/z)^2 - 1}.$

Wide, deep well: Intersections are at $\pi, 2\pi, 3\pi$, etc. Same as Eq. 2.157, but now for n even. This fills in the rest of the states for the infinite square well.

Shallow, narrow well: If $z_0 < \pi/2$, there is *no* odd bound state. The corresponding condition on V_0 is

$$V_0 < \frac{\pi^2 \hbar^2}{8ma^2} \Rightarrow \text{no odd bound state.}$$



Problem 2.30

$$\begin{aligned} 1 &= 2 \int_0^\infty |\psi|^2 dx = 2 \left(|D|^2 \int_0^a \cos^2 lx dx + |F|^2 \int_a^\infty e^{-2\kappa x} dx \right) \\ &= 2 \left[|D|^2 \left(\frac{x}{2} + \frac{1}{4l} \sin 2lx \right) \Big|_0^a + |F|^2 \left(-\frac{1}{2\kappa} e^{-2\kappa x} \right) \Big|_a^\infty \right] = 2 \left[|D|^2 \left(\frac{a}{2} + \frac{\sin 2la}{4l} \right) + |F|^2 \frac{e^{-2\kappa a}}{2\kappa} \right]. \end{aligned}$$

But $F = De^{\kappa a} \cos la$ (Eq. 2.152), so $1 = |D|^2 \left(a + \frac{\sin(2la)}{2l} + \frac{\cos^2(la)}{\kappa} \right)$.

Furthermore $\kappa = l \tan(la)$ (Eq. 2.154), so

$$\begin{aligned} 1 &= |D|^2 \left(a + \frac{2 \sin la \cos la}{2l} + \frac{\cos^3 la}{l \sin la} \right) = |D|^2 \left[a + \frac{\cos la}{l \sin la} (\sin^2 la + \cos^2 la) \right] \\ &= |D|^2 \left(a + \frac{1}{l \tan la} \right) = |D|^2 \left(a + \frac{1}{\kappa} \right). \end{aligned} \quad \boxed{D = \frac{1}{\sqrt{a + 1/\kappa}}}, \quad \boxed{F = \frac{e^{\kappa a} \cos la}{\sqrt{a + 1/\kappa}}}.$$

Problem 2.31

Equation 2.155 $\Rightarrow z_0 = \frac{a}{\hbar} \sqrt{2mV_0}$. We want $\alpha = \text{area of potential} = 2aV_0$ held constant as $a \rightarrow 0$. Therefore $V_0 = \frac{\alpha}{2a}$; $z_0 = \frac{a}{\hbar} \sqrt{2m \frac{\alpha}{2a}} = \frac{1}{\hbar} \sqrt{m\alpha a} \rightarrow 0$. So z_0 is *small*, and the intersection in Fig. 2.18 occurs at very small z . Solve Eq. 2.156 for very small z , by expanding $\tan z$:

$$\tan z \cong z = \sqrt{(z_0/z)^2 - 1} = (1/z) \sqrt{z_0^2 - z^2}.$$

Now (from Eqs. 2.146, 2.148 and 2.155) $z_0^2 - z^2 = \kappa^2 a^2$, so $z^2 = \kappa a$. But $z_0^2 - z^2 = z^4 \ll 1 \Rightarrow z \cong z_0$, so $\kappa a \cong z_0^2$. But we found that $z_0 \cong \frac{1}{\hbar} \sqrt{m\alpha a}$ here, so $\kappa a = \frac{1}{\hbar^2} m\alpha a$, or $\kappa = \frac{m\alpha}{\hbar^2}$. (At this point the a 's have canceled, and we can go to the limit $a \rightarrow 0$.)

$$\frac{\sqrt{-2mE}}{\hbar} = \frac{m\alpha}{\hbar^2} \Rightarrow -2mE = \frac{m^2 \alpha^2}{\hbar^2}. \quad \boxed{E = -\frac{m\alpha^2}{2\hbar^2}} \quad (\text{which agrees with Eq. 2.129}).$$

In Eq. 2.169, $V_0 \gg E \Rightarrow T^{-1} \cong 1 + \frac{V_0^2}{4EV_0} \sin^2 \left(\frac{2a}{\hbar} \sqrt{2mV_0} \right)$. But $V_0 = \frac{\alpha}{2a}$, so the argument of the sine is small, and we can replace $\sin \epsilon$ by ϵ : $T^{-1} \cong 1 + \frac{V_0}{4E} \left(\frac{2a}{\hbar} \right)^2 2mV_0 = 1 + (2aV_0)^2 \frac{m}{2\hbar^2 E}$. But $2aV_0 = \alpha$, so $T^{-1} = 1 + \frac{m\alpha^2}{2\hbar^2 E}$, in agreement with Eq. 2.141.

Problem 2.32

Multiply Eq. 2.165 by $\sin la$, Eq. 2.166 by $\frac{1}{l} \cos la$, and add:

$$\left. \begin{aligned} C \sin^2 la + D \sin la \cos la &= F e^{ika} \sin la \\ C \cos^2 la - D \sin la \cos la &= \frac{ik}{l} F e^{ika} \cos la \end{aligned} \right\} \quad C = F e^{ika} \left[\sin la + \frac{ik}{l} \cos la \right].$$

Multiply Eq. 2.165 by $\cos la$, Eq. 2.166 by $\frac{1}{l} \sin la$, and subtract:

$$\left. \begin{aligned} C \sin la \cos la + D \cos^2 la &= F e^{ika} \cos la \\ C \sin la \cos la - D \sin^2 la &= \frac{ik}{l} F e^{ika} \sin la \end{aligned} \right\} \quad D = F e^{ika} \left[\cos la - \frac{ik}{l} \sin la \right].$$

Put these into Eq. 2.163:

$$\begin{aligned} (1) \quad A e^{-ika} + B e^{ika} &= -F e^{ika} \left[\sin la + \frac{ik}{l} \cos la \right] \sin la + F e^{ika} \left[\cos la - \frac{ik}{l} \sin la \right] \cos la \\ &= F e^{ika} \left[\cos^2 la - \frac{ik}{l} \sin la \cos la - \sin^2 la - \frac{ik}{l} \sin la \cos la \right] \\ &= F e^{ika} \left[\cos(2la) - \frac{ik}{l} \sin(2la) \right]. \end{aligned}$$

Likewise, from Eq. 2.164:

$$\begin{aligned} (2) \quad A e^{-ika} - B e^{ika} &= -\frac{il}{k} F e^{ika} \left[\left(\sin la + \frac{ik}{l} \cos la \right) \cos la + \left(\cos la - \frac{ik}{l} \sin la \right) \sin la \right] \\ &= -\frac{il}{k} F e^{ika} \left[\sin la \cos la + \frac{ik}{l} \cos^2 la + \sin la \cos la - \frac{ik}{l} \sin^2 la \right] \\ &= -\frac{il}{k} F e^{ika} \left[\sin(2la) + \frac{ik}{l} \cos(2la) \right] = F e^{ika} \left[\cos(2la) - \frac{il}{k} \sin(2la) \right]. \end{aligned}$$

Add (1) and (2): $2A e^{-ika} = F e^{ika} \left[2 \cos(2la) - i \left(\frac{k}{l} + \frac{l}{k} \right) \sin(2la) \right]$, or:

$$F = \frac{e^{-2ika} A}{\cos(2la) - i \frac{\sin(2la)}{2kl} (k^2 + l^2)} \quad (\text{confirming Eq. 2.168}). \quad \text{Now subtract (2) from (1):}$$

$$2B e^{ika} = F e^{ika} \left[i \left(\frac{l}{k} - \frac{k}{l} \right) \sin(2la) \right] \Rightarrow B = i \frac{\sin(2la)}{2kl} (l^2 - k^2) F \quad (\text{confirming Eq. 2.167}).$$

$$T^{-1} = \left| \frac{A}{F} \right|^2 = \left| \cos(2la) - i \frac{\sin(2la)}{2kl} (k^2 + l^2) \right|^2 = \cos^2(2la) + \frac{\sin^2(2la)}{(2kl)^2} (k^2 + l^2)^2.$$

But $\cos^2(2la) = 1 - \sin^2(2la)$, so

$$T^{-1} = 1 + \sin^2(2la) \left[\underbrace{\frac{(k^2 + l^2)^2}{(2kl)^2} - 1}_{\frac{1}{(2kl)^2} [k^4 + 2k^2 l^2 + l^4 - 4k^2 l^2]} \right] = 1 + \frac{(k^2 - l^2)^2}{(2kl)^2} \sin^2(2la).$$

$$\frac{1}{(2kl)^2} [k^4 + 2k^2 l^2 + l^4 - 4k^2 l^2] = \frac{1}{(2kl)^2} [k^4 - 2k^2 l^2 + l^4] = \frac{(k^2 - l^2)^2}{(2kl)^2}.$$

But $k = \frac{\sqrt{2mE}}{\hbar}$, $l = \frac{\sqrt{2m(E+V_0)}}{\hbar}$; so $(2la) = \frac{2a}{\hbar} \sqrt{2m(E+V_0)}$; $k^2 - l^2 = -\frac{2mV_0}{\hbar^2}$, and

$$\frac{(k^2 - l^2)^2}{(2kl)^2} = \frac{\left(\frac{2m}{\hbar^2}\right)^2 V_0^2}{4 \left(\frac{2m}{\hbar^2}\right)^2 E(E+V_0)} = \frac{V_0^2}{4E(E+V_0)}.$$

$$\therefore T^{-1} = 1 + \frac{V_0^2}{4E(E+V_0)} \sin^2 \left(\frac{2a}{\hbar} \sqrt{2m(E+V_0)} \right), \text{ confirming Eq. 2.169.}$$

Problem 2.33

$$\underline{E < V_0}. \quad \psi = \begin{cases} Ae^{ikx} + Be^{-ikx} & (x < -a) \\ Ce^{\kappa x} + De^{-\kappa x} & (-a < x < a) \\ Fe^{ikx} & (x > a) \end{cases} \quad k = \frac{\sqrt{2mE}}{\hbar}; \quad \kappa = \frac{\sqrt{2m(V_0 - E)}}{\hbar}.$$

(1) Continuity of ψ at $-a$: $Ae^{-ika} + Be^{ika} = Ce^{-\kappa a} + De^{\kappa a}$.

(2) Continuity of ψ' at $-a$: $ik(Ae^{-ika} - Be^{ika}) = \kappa(Ce^{-\kappa a} - De^{\kappa a})$.

$$\Rightarrow 2Ae^{-ika} = \left(1 - i\frac{\kappa}{k}\right) Ce^{-\kappa a} + \left(1 + i\frac{\kappa}{k}\right) De^{\kappa a}.$$

(3) Continuity of ψ at $+a$: $Ce^{\kappa a} + De^{-\kappa a} = Fe^{ika}$.

(4) Continuity of ψ' at $+a$: $\kappa(Ce^{\kappa a} - De^{-\kappa a}) = ikFe^{ika}$.

$$\Rightarrow 2Ce^{\kappa a} = \left(1 + \frac{ik}{\kappa}\right) Fe^{ika}; \quad 2De^{-\kappa a} = \left(1 - \frac{ik}{\kappa}\right) Fe^{ika}.$$

$$\begin{aligned} 2Ae^{-ika} &= \left(1 - \frac{i\kappa}{k}\right) \left(1 + \frac{ik}{\kappa}\right) Fe^{ika} \frac{e^{-2\kappa a}}{2} + \left(1 + \frac{i\kappa}{k}\right) \left(1 - \frac{ik}{\kappa}\right) Fe^{ika} \frac{e^{2\kappa a}}{2} \\ &= \frac{Fe^{ika}}{2} \left\{ \left[1 + i\left(\frac{k}{\kappa} - \frac{\kappa}{k}\right) + 1\right] e^{-2\kappa a} + \left[1 + i\left(\frac{\kappa}{k} - \frac{k}{\kappa}\right) + 1\right] e^{2\kappa a} \right\} \\ &= \frac{Fe^{ika}}{2} \left[2(e^{-2\kappa a} + e^{2\kappa a}) + i\frac{(\kappa^2 - k^2)}{k\kappa} (e^{2\kappa a} - e^{-2\kappa a}) \right]. \end{aligned}$$

$$\begin{aligned} \text{But } \sinh x &\equiv \frac{e^x - e^{-x}}{2}, \quad \cosh x \equiv \frac{e^x + e^{-x}}{2}, \text{ so} \\ &= \frac{Fe^{ika}}{2} \left[4 \cosh(2\kappa a) + i\frac{(\kappa^2 - k^2)}{k\kappa} 2 \sinh(2\kappa a) \right] \\ &= 2Fe^{ika} \left[\cosh(2\kappa a) + i\frac{(\kappa^2 - k^2)}{2k\kappa} \sinh(2\kappa a) \right]. \end{aligned}$$

$$T^{-1} = \left| \frac{A}{F} \right|^2 = \cosh^2(2\kappa a) + \frac{(\kappa^2 - k^2)^2}{(2\kappa k)^2} \sinh^2(2\kappa a). \quad \text{But } \cosh^2 = 1 + \sinh^2, \text{ so}$$

$$T^{-1} = 1 + \underbrace{\left[1 + \frac{(\kappa^2 - k^2)^2}{(2\kappa k)^2} \right]}_{\star} \sinh^2(2\kappa a) = \boxed{1 + \frac{V_0^2}{4E(V_0 - E)} \sinh^2 \left(\frac{2a}{\hbar} \sqrt{2m(V_0 - E)} \right)},$$

$$\text{where } \star = \frac{4\kappa^2 k^2 + k^4 + \kappa^4 - 2\kappa^2 k^2}{(2\kappa k)^2} = \frac{(\kappa^2 + k^2)^2}{(2\kappa k)^2} = \frac{\left(\frac{2mE}{\hbar^2} + \frac{2m(V_0 - E)}{\hbar^2}\right)^2}{4 \frac{2mE}{\hbar^2} \frac{2m(V_0 - E)}{\hbar^2}} = \frac{V_0^2}{4E(V_0 - E)}.$$

(You can also get this from Eq. 2.169 by switching the sign of V_0 and using $\sin(i\theta) = i \sinh \theta$.)

$$\underline{\mathbf{E} = \mathbf{V}_0.} \quad \psi = \begin{cases} Ae^{ikx} + Be^{-ikx} & (x < -a) \\ C + Dx & (-a < x < a) \\ Fe^{ikx} & (x > a) \end{cases}$$

$$(\text{In central region } -\frac{\hbar^2}{2m} \frac{d^2\psi}{dx^2} + V_0\psi = E\psi \Rightarrow \frac{d^2\psi}{dx^2} = 0, \text{ so } \psi = C + Dx.)$$

$$(1) \text{ Continuous } \psi \text{ at } -a : Ae^{-ika} + Be^{ika} = C - Da.$$

$$(2) \text{ Continuous } \psi \text{ at } +a : Fe^{ika} = C + Da.$$

$$\Rightarrow (2.5) \quad 2Da = Fe^{ika} - Ae^{-ika} - Be^{ika}.$$

$$(3) \text{ Continuous } \psi' \text{ at } -a : ik(Ae^{-ika} - Be^{ika}) = D.$$

$$(4) \text{ Continuous } \psi' \text{ at } +a : ikFe^{ika} = D.$$

$$\Rightarrow (4.5) \quad Ae^{-2ika} - B = F.$$

Use (4) to eliminate D in (2.5): $Ae^{-2ika} + B = F - 2aikF = (1 - 2iak)F$, and add to (4.5):

$$2Ae^{-2ika} = 2F(1 - ika), \text{ so } T^{-1} = \left| \frac{A}{F} \right|^2 = 1 + (ka)^2 = \boxed{1 + \frac{2mE}{\hbar^2} a^2}.$$

(You can also get this from Eq. 2.169 by changing the sign of V_0 and taking the limit $E \rightarrow V_0$, using $\sin \epsilon \cong \epsilon$.)

$\mathbf{E} > \mathbf{V}_0$. This case is identical to the one in the book, only with $V_0 \rightarrow -V_0$. So

$$\boxed{T^{-1} = 1 + \frac{V_0^2}{4E(E - V_0)} \sin^2 \left(\frac{2a}{\hbar} \sqrt{2m(E - V_0)} \right)}.$$

Problem 2.34

(a)

$$\psi = \begin{cases} Ae^{ikx} + Be^{-ikx} & (x < 0) \\ Fe^{-\kappa x} & (x > 0) \end{cases} \text{ where } k = \frac{\sqrt{2mE}}{\hbar}; \quad \kappa = \frac{\sqrt{2m(V_0 - E)}}{\hbar}.$$

$$(1) \text{ Continuity of } \psi : A + B = F.$$

$$(2) \text{ Continuity of } \psi' : ik(A - B) = -\kappa F.$$

$$\Rightarrow A + B = -\frac{ik}{\kappa}(A - B) \Rightarrow A \left(1 + \frac{ik}{\kappa}\right) = -B \left(1 - \frac{ik}{\kappa}\right).$$

$$R = \left| \frac{B}{A} \right|^2 = \frac{|(1 + ik/\kappa)|^2}{|(1 - ik/\kappa)|^2} = \frac{1 + (k/\kappa)^2}{1 + (k/\kappa)^2} = \boxed{1}.$$

Although the wave function penetrates into the barrier, it is eventually all reflected.

(b)

$$\psi = \begin{cases} Ae^{ikx} + Be^{-ikx} & (x < 0) \\ Fe^{ilx} & (x > 0) \end{cases} \text{ where } k = \frac{\sqrt{2mE}}{\hbar}; \quad l = \frac{\sqrt{2m(E - V_0)}}{\hbar}.$$

(1) Continuity of ψ : $A + B = F$.

(2) Continuity of ψ' : $ik(A - B) = ilF$.

$$\Rightarrow A + B = \frac{k}{l}(A - B); \quad A \left(1 - \frac{k}{l}\right) = -B \left(1 + \frac{k}{l}\right).$$

$$R = \left| \frac{B}{A} \right|^2 = \frac{(1 - k/l)^2}{(1 + k/l)^2} = \frac{(k - l)^2}{(k + l)^2} = \frac{(k - l)^4}{(k^2 - l^2)^2}.$$

$$\text{Now } k^2 - l^2 = \frac{2m}{\hbar^2}(E - E + V_0) = \left(\frac{2m}{\hbar^2}\right)V_0; \quad k - l = \frac{\sqrt{2m}}{\hbar}[\sqrt{E} - \sqrt{E - V_0}], \text{ so}$$

$$R = \frac{(\sqrt{E} - \sqrt{E - V_0})^4}{V_0^2}.$$

(c)



From the diagram, $T = P_t/P_i = |F|^2 v_t / |A|^2 v_i$, where P_i is the probability of finding the incident particle in the box corresponding to the time interval dt , and P_t is the probability of finding the transmitted particle in the associated box to the *right* of the barrier.

But $\frac{v_t}{v_i} = \frac{\sqrt{E - V_0}}{\sqrt{E}}$ (from Eq. 2.98). So $T = \sqrt{\frac{E - V_0}{E}} \left| \frac{F}{A} \right|^2$. Alternatively, from Problem 2.19:

$$J_i = \frac{\hbar k}{m} |A|^2; \quad J_t = \frac{\hbar l}{m} |F|^2; \quad T = \frac{J_t}{J_i} = \left| \frac{F}{A} \right|^2 \frac{l}{k} = \left| \frac{F}{A} \right|^2 \sqrt{\frac{E - V_0}{E}}.$$

For $E < V_0$, of course, $T = 0$.

(d)

$$\text{For } E > V_0, \quad F = A + B = A + A \left(\frac{\frac{k}{l} - 1}{\frac{k}{l} + 1} \right) = A \frac{2k/l}{\left(\frac{k}{l} + 1 \right)} = \frac{2k}{k + l} A.$$

$$T = \left| \frac{F}{A} \right|^2 \frac{l}{k} = \left(\frac{2k}{k + l} \right)^2 \frac{l}{k} = \frac{4kl}{(k + l)^2} = \frac{4kl(k - l)^2}{(k^2 - l^2)^2} = \frac{4\sqrt{E}\sqrt{E - V_0}(\sqrt{E} - \sqrt{E - V_0})^2}{V_0^2}.$$

$$T + R = \frac{4kl}{(k + l)^2} + \frac{(k - l)^2}{(k + l)^2} = \frac{4kl + k^2 - 2kl + l^2}{(k + l)^2} = \frac{k^2 + 2kl + l^2}{(k + l)^2} = \frac{(k + l)^2}{(k + l)^2} = 1. \quad \checkmark$$

Problem 2.35

(a)


$$\psi(x) = \begin{cases} Ae^{ikx} + Be^{-ikx} & (x < 0) \\ Fe^{ilx} & (x > 0) \end{cases} \text{ where } k \equiv \frac{\sqrt{2mE}}{\hbar}, \quad l \equiv \frac{\sqrt{2m(E+V_0)}}{\hbar}.$$

$$\left. \begin{array}{l} \text{Continuity of } \psi \Rightarrow A + B = F \\ \text{Continuity of } \psi' \Rightarrow ik(A - B) = ilF \end{array} \right\} \Rightarrow$$

$$A + B = \frac{k}{l}(A - B); \quad A \left(1 - \frac{k}{l}\right) = -B \left(1 + \frac{k}{l}\right); \quad \frac{B}{A} = -\left(\frac{1 - k/l}{1 + k/l}\right).$$

$$\begin{aligned} R &= \left|\frac{B}{A}\right|^2 = \left(\frac{l - k}{l + k}\right)^2 = \left(\frac{\sqrt{E+V_0} - \sqrt{E}}{\sqrt{E+V_0} + \sqrt{E}}\right)^2 \\ &= \left(\frac{\sqrt{1+V_0/E} - 1}{\sqrt{1+V_0/E} + 1}\right)^2 = \left(\frac{\sqrt{1+3} - 1}{\sqrt{1+3} + 1}\right)^2 = \left(\frac{2-1}{2+1}\right)^2 = \boxed{\frac{1}{9}}. \end{aligned}$$

- (b) The cliff is *two-dimensional*, and even if we pretend the car drops straight down, the potential *as a function of distance along the* (crooked, but now one-dimensional) *path* is $-mgx$ (with x the vertical coordinate), as shown.



- (c) Here $V_0/E = 12/4 = 3$, the same as in part (a), so $R = 1/9$, and hence $T = \boxed{8/9 = 0.8889}$.

Problem 2.36

Start with Eq. 2.22: $\psi(x) = A \sin kx + B \cos kx$. This time the boundary conditions are $\psi(a) = \psi(-a) = 0$:

$$A \sin ka + B \cos ka = 0; \quad -A \sin ka + B \cos ka = 0.$$

$$\begin{cases} \text{Subtract :} & A \sin ka = 0 \Rightarrow ka = j\pi \text{ or } A = 0, \\ \text{Add :} & B \cos ka = 0 \Rightarrow ka = (j - \frac{1}{2})\pi \text{ or } B = 0, \end{cases}$$

(where $j = 1, 2, 3, \dots$).

If $B = 0$ (so $A \neq 0$), $k = j\pi/a$. In this case let $n \equiv 2j$ (so n is an *even* integer); then $k = n\pi/2a$, $\psi = A \sin(n\pi x/2a)$. Normalizing: $1 = |A|^2 \int_{-a}^a \sin^2(n\pi x/2a) dx = |A|^2 a \Rightarrow A = 1/\sqrt{a}$.


If $A = 0$ (so $B \neq 0$), $k = (j - \frac{1}{2})\pi/a$. In this case let $n \equiv 2j - 1$ (n is an *odd* integer); again $k = n\pi/2a$, $\psi = B \cos(n\pi x/2a)$. Normalizing: $1 = |B|^2 \int_{-a}^a \cos^2(n\pi x/2a) dx = |B|^2 a \Rightarrow B = 1/\sqrt{a}$.

In either case Eq. 2.21 yields $E = \frac{\hbar^2 k^2}{2m} = \frac{n^2 \pi^2 \hbar^2}{2m(2a)^2}$ (in agreement with Eq. 2.27 for a well of width $2a$).

The substitution $x \rightarrow (x + a)/2$ takes Eq. 2.28 to

$$\sqrt{\frac{2}{a}} \sin\left(\frac{n\pi}{a} \frac{(x+a)}{2}\right) = \sqrt{\frac{2}{a}} \sin\left(\frac{n\pi x}{2a} + \frac{n\pi}{2}\right) = \begin{cases} (-1)^{n/2} \sqrt{\frac{2}{a}} \sin\left(\frac{n\pi x}{2a}\right) & (n \text{ even}), \\ (-1)^{(n-1)/2} \sqrt{\frac{2}{a}} \cos\left(\frac{n\pi x}{2a}\right) & (n \text{ odd}). \end{cases}$$

So (apart from normalization) we recover the results above. The graphs are the same as Figure 2.2, except that some are upside down (different normalization).



Problem 2.37

Use the trig identity $\sin 3\theta = 3 \sin \theta - 4 \sin^3 \theta$ to write

$$\sin^3\left(\frac{\pi x}{a}\right) = \frac{3}{4} \sin\left(\frac{\pi x}{a}\right) - \frac{1}{4} \sin\left(\frac{3\pi x}{a}\right). \quad \text{So (Eq. 2.28): } \Psi(x, 0) = A \sqrt{\frac{a}{2}} \left[\frac{3}{4} \psi_1(x) - \frac{1}{4} \psi_3(x) \right].$$

Normalize using Eq. 2.38: $|A|^2 \frac{a}{2} \left(\frac{9}{16} + \frac{1}{16} \right) = \frac{5}{16} a |A|^2 = 1 \Rightarrow \boxed{A = \frac{4}{\sqrt{5a}}}.$

So $\Psi(x, 0) = \frac{1}{\sqrt{10}} [3\psi_1(x) - \psi_3(x)]$, and hence (Eq. 2.17)

$$\boxed{\Psi(x, t) = \frac{1}{\sqrt{10}} \left[3\psi_1(x)e^{-iE_1 t/\hbar} - \psi_3(x)e^{-iE_3 t/\hbar} \right].}$$

$$|\Psi(x, t)|^2 = \frac{1}{10} \left[9\psi_1^2 + \psi_3^2 - 6\psi_1\psi_3 \cos\left(\frac{E_3 - E_1}{\hbar}t\right) \right]; \quad \text{so}$$

$$\langle x \rangle = \int_0^a x |\Psi(x, t)|^2 dx = \frac{9}{10} \langle x \rangle_1 + \frac{1}{10} \langle x \rangle_3 - \frac{3}{5} \cos\left(\frac{E_3 - E_1}{\hbar}t\right) \int_0^a x \psi_1(x) \psi_3(x) dx,$$

where $\langle x \rangle_n = a/2$ is the expectation value of x in the n th stationary state. The remaining integral is

$$\begin{aligned} \frac{2}{a} \int_0^a x \sin\left(\frac{\pi x}{a}\right) \sin\left(\frac{3\pi x}{a}\right) dx &= \frac{1}{a} \int_0^a x \left[\cos\left(\frac{2\pi x}{a}\right) - \cos\left(\frac{4\pi x}{a}\right) \right] dx \\ &= \frac{1}{a} \left[\left(\frac{a}{2\pi}\right)^2 \cos\left(\frac{2\pi x}{a}\right) + \left(\frac{xa}{2\pi}\right) \sin\left(\frac{2\pi x}{a}\right) - \left(\frac{a}{4\pi}\right)^2 \cos\left(\frac{4\pi x}{a}\right) - \left(\frac{xa}{4\pi}\right) \sin\left(\frac{4\pi x}{a}\right) \right]_0^a = 0. \end{aligned}$$

Evidently then,

$$\langle x \rangle = \frac{9}{10} \left(\frac{a}{2}\right) + \frac{1}{10} \left(\frac{a}{2}\right) = \boxed{\frac{a}{2}}.$$

Using Eq. 2.39,

$$\langle H \rangle = p_1 E_1 + p_3 E_3 = \left(\frac{9}{10}\right) \frac{\pi^2 \hbar^2}{2ma^2} + \left(\frac{1}{10}\right) \frac{9\pi^2 \hbar^2}{2ma^2} = \boxed{\frac{9\pi^2 \hbar^2}{10ma^2}}.$$

Problem 2.38

(a) New allowed energies: $E_n = \frac{n^2 \pi^2 \hbar^2}{2m(2a)^2}$; $\Psi(x, 0) = \sqrt{\frac{2}{a}} \sin\left(\frac{\pi}{a}x\right)$, $\psi_n(x) = \sqrt{\frac{2}{2a}} \sin\left(\frac{n\pi}{2a}x\right)$.

$$\begin{aligned} c_n &= \frac{\sqrt{2}}{a} \int_0^a \sin\left(\frac{\pi}{a}x\right) \sin\left(\frac{n\pi}{2a}x\right) dx = \frac{\sqrt{2}}{2a} \int_0^a \left\{ \cos\left[\left(\frac{n}{2} - 1\right) \frac{\pi x}{a}\right] - \cos\left[\left(\frac{n}{2} + 1\right) \frac{\pi x}{a}\right] \right\} dx \\ &= \frac{1}{\sqrt{2}a} \left\{ \frac{\sin\left[\left(\frac{n}{2} - 1\right) \frac{\pi x}{a}\right]}{\left(\frac{n}{2} - 1\right) \frac{\pi}{a}} - \frac{\sin\left[\left(\frac{n}{2} + 1\right) \frac{\pi x}{a}\right]}{\left(\frac{n}{2} + 1\right) \frac{\pi}{a}} \right\} \Big|_0^a \quad (\text{for } n \neq 2) \\ &= \frac{1}{\sqrt{2}\pi} \left\{ \frac{\sin\left[\left(\frac{n}{2} - 1\right) \pi\right]}{\left(\frac{n}{2} - 1\right)} - \frac{\sin\left[\left(\frac{n}{2} + 1\right) \pi\right]}{\left(\frac{n}{2} + 1\right)} \right\} = \frac{\sin\left[\left(\frac{n}{2} + 1\right) \pi\right]}{\sqrt{2}\pi} \left[\frac{1}{\left(\frac{n}{2} - 1\right)} - \frac{1}{\left(\frac{n}{2} + 1\right)} \right] \\ &= \frac{4\sqrt{2}}{\pi} \frac{\sin\left[\left(\frac{n}{2} + 1\right) \pi\right]}{(n^2 - 4)} = \begin{cases} 0, & \text{if } n \text{ is even} \\ \pm \frac{4\sqrt{2}}{\pi(n^2 - 4)}, & \text{if } n \text{ is odd} \end{cases}. \end{aligned}$$

$$c_2 = \frac{\sqrt{2}}{a} \int_0^a \sin^2\left(\frac{\pi}{a}x\right) dx = \frac{\sqrt{2}}{a} \int_0^a \frac{1}{2} dx = \frac{1}{\sqrt{2}}. \quad \text{So the probability of getting } E_n \text{ is}$$

$$P_n = |c_n|^2 = \begin{cases} \frac{1}{2}, & \text{if } n = 2 \\ \frac{32}{\pi^2(n^2 - 4)^2}, & \text{if } n \text{ is odd} \\ 0, & \text{otherwise} \end{cases}.$$

Most probable: $E_2 = \boxed{\frac{\pi^2 \hbar^2}{2ma^2}}$ (same as before). Probability: $P_2 = \boxed{1/2}$.

(b) Next most probable: $E_1 = \boxed{\frac{\pi^2 \hbar^2}{8ma^2}}$, with probability $P_1 = \boxed{\frac{32}{9\pi^2} = 0.36025}$.

(c) $\langle H \rangle = \int \Psi^* H \Psi dx = \frac{2}{a} \int_0^a \sin\left(\frac{\pi}{a}x\right) \left(-\frac{\hbar^2}{2m} \frac{d^2}{dx^2}\right) \sin\left(\frac{\pi}{a}x\right) dx$, but this is exactly the same as before the wall moved – for which we know the answer: $\boxed{\frac{\pi^2 \hbar^2}{2ma^2}}$.

Problem 2.39

(a) According to Eq. 2.36, the most general solution to the time-dependent Schrödinger equation for the infinite square well is

$$\Psi(x, t) = \sum_{n=1}^{\infty} c_n \psi_n(x) e^{-i(n^2 \pi^2 \hbar / 2ma^2)t}.$$

Now $\frac{n^2 \pi^2 \hbar}{2ma^2} T = \frac{n^2 \pi^2 \hbar}{2ma^2} \frac{4ma^2}{\pi \hbar} = 2\pi n^2$, so $e^{-i(n^2 \pi^2 \hbar / 2ma^2)(t+T)} = e^{-i(n^2 \pi^2 \hbar / 2ma^2)t} e^{-i2\pi n^2}$, and since n^2 is an integer, $e^{-i2\pi n^2} = 1$. Therefore $\Psi(x, t+T) = \Psi(x, t)$. QED

(b) The classical revival time is the time it takes the particle to go down and back: $T_c = 2a/v$, with the velocity given by

$$E = \frac{1}{2}mv^2 \Rightarrow v = \sqrt{\frac{2E}{m}} \Rightarrow T_c = a\sqrt{\frac{2m}{E}}.$$

(c) The two revival times are equal if

$$\frac{4ma^2}{\pi\hbar} = a\sqrt{\frac{2m}{E}}, \quad \text{or} \quad \boxed{E = \frac{\pi^2\hbar^2}{8ma^2} = \frac{E_1}{4}}.$$

Problem 2.40

(a) Let $V_0 \equiv 32\hbar^2/ma^2$. This is just like the *odd* bound states for the finite square well, since they are the ones that go to zero at the origin. Referring to the solution to Problem 2.29, the wave function is

$$\psi(x) = \begin{cases} D \sin lx, & l \equiv \sqrt{2m(E + V_0)}/\hbar \quad (0 < x < a), \\ F e^{-\kappa x}, & \kappa \equiv \sqrt{-2mE}/\hbar \quad (x > a), \end{cases}$$

and the boundary conditions at $x = a$ yield

$$-\cot z = \sqrt{(z_0/z)^2 - 1}$$

with

$$z_0 = \frac{\sqrt{2mV_0}}{\hbar}a = \frac{\sqrt{2m(32\hbar^2/ma^2)}}{\hbar}a = 8.$$

Referring to the figure (Problem 2.29), and noting that $(5/2)\pi = 7.85 < z_0 < 3\pi = 9.42$, we see that there are three bound states.

(b) Let

$$\begin{aligned} I_1 &\equiv \int_0^a |\psi|^2 dx = |D|^2 \int_0^a \sin^2 lx \, dx = |D|^2 \left[\frac{x}{2} - \frac{1}{2l} \sin lx \cos lx \right]_0^a = |D|^2 \left[\frac{a}{2} - \frac{1}{2l} \sin la \cos la \right]; \\ I_2 &\equiv \int_a^\infty |\psi|^2 dx = |F|^2 \int_a^\infty e^{-2\kappa x} dx = |F|^2 \left[-\frac{e^{-2\kappa x}}{2\kappa} \right]_a^\infty = |F|^2 \frac{e^{-2\kappa a}}{2\kappa}. \end{aligned}$$

But continuity at $x = a \Rightarrow F e^{-\kappa a} = D \sin la$, so $I_2 = |D|^2 \frac{\sin^2 la}{2\kappa}$.

Normalizing:

$$1 = I_1 + I_2 = |D|^2 \left[\frac{a}{2} - \frac{1}{2l} \sin la \cos la + \frac{\sin^2 la}{2\kappa} \right] = \frac{1}{2\kappa} |D|^2 \left[\kappa a - \frac{\kappa}{l} \sin la \cos la + \sin^2 la \right]$$

But (referring again to Problem 2.29) $\kappa/l = -\cot la$, so

$$= \frac{1}{2\kappa} |D|^2 [\kappa a + \cot la \sin la \cos la + \sin^2 la] = |D|^2 \frac{(1 + \kappa a)}{2\kappa}.$$

So $|D|^2 = 2\kappa/(1 + \kappa a)$, and the probability of finding the particle outside the well is

$$P = I_2 = \frac{2\kappa}{1 + \kappa a} \frac{\sin^2 la}{2\kappa} = \frac{\sin^2 la}{1 + \kappa a}.$$

We can express this in terms of $z \equiv la$ and z_0 : $\kappa a = \sqrt{z_0^2 - z^2}$ (page 80),

$$\sin^2 la = \sin^2 z = \frac{1}{1 + \cot^2 z} = \frac{1}{1 + (z_0/z)^2 - 1} = \left(\frac{z}{z_0} \right)^2 \Rightarrow P = \frac{z^2}{z_0^2(1 + \sqrt{z_0^2 - z^2})}.$$

So far, this is correct for *any* bound state. In the present case $z_0 = 8$ and z is the third solution to $-\cot z = \sqrt{(8/z)^2 - 1}$, which occurs somewhere in the interval $7.85 < z < 8$. Mathematica gives $z = 7.9573$ and $P = 0.54204$.

```
FindRoot[Cot[z] == -Sqrt(8/z)^2 - 1, {z, 7.9}]
{z -> 7.95732}

z^2 / (64 (1 + Sqrt(64 - z^2)))

z^2
-----
64 (1 + Sqrt(64 - z^2))

*/. z -> 7.957321523328964`
0.542041
```

Problem 2.41

- (a) In the standard notation $\xi \equiv \sqrt{m\omega/\hbar}x$, $\alpha \equiv (m\omega/\pi\hbar)^{1/4}$,

$$\Psi(x, 0) = A(1 - 2\xi)^2 e^{-\xi^2/2} = A(1 - 4\xi + 4\xi^2)e^{-\xi^2/2}.$$

It can be expressed as a linear combination of the first three stationary states (Eq. 2.59 and 2.62, and Problem 2.10):

$$\psi_0(x) = \alpha e^{-\xi^2/2}, \quad \psi_1(x) = \sqrt{2}\alpha\xi e^{-\xi^2/2}, \quad \psi_2(x) = \frac{\alpha}{\sqrt{2}}(2\xi^2 - 1)e^{-\xi^2/2}.$$

So $\Psi(x, 0) = c_0\psi_0 + c_1\psi_1 + c_2\psi_2 = \alpha(c_0 + \sqrt{2}\xi c_1 + \sqrt{2}\xi^2 c_2 - \frac{1}{\sqrt{2}}c_2)e^{-\xi^2/2}$ with (equating like powers)

$$\begin{cases} \alpha\sqrt{2}c_2 = 4A & \Rightarrow c_2 = 2\sqrt{2}A/\alpha, \\ \alpha\sqrt{2}c_1 = -4A & \Rightarrow c_1 = -2\sqrt{2}A/\alpha, \\ \alpha(c_0 - c_2/\sqrt{2}) = A & \Rightarrow c_0 = (A/\alpha) + c_2/\sqrt{2} = (1 + 2)A/\alpha = 3A/\alpha. \end{cases}$$

Normalizing: $1 = |c_0|^2 + |c_1|^2 + |c_2|^2 = (8 + 8 + 9)(A/\alpha)^2 = 25(A/\alpha)^2 \Rightarrow A = \alpha/5$.

$$c_0 = \frac{3}{5}, \quad c_1 = -\frac{2\sqrt{2}}{5}, \quad c_2 = \frac{2\sqrt{2}}{5}.$$

$$\langle H \rangle = \sum |c_n|^2 \left(n + \frac{1}{2}\right) \hbar\omega = \frac{9}{25} \left(\frac{1}{2}\hbar\omega\right) + \frac{8}{25} \left(\frac{3}{2}\hbar\omega\right) + \frac{8}{25} \left(\frac{5}{2}\hbar\omega\right) = \frac{\hbar\omega}{50} (9 + 24 + 40) = \boxed{\frac{73}{50}\hbar\omega}.$$

- (b)

$$\Psi(x, t) = \frac{3}{5}\psi_0 e^{-i\omega t/2} - \frac{2\sqrt{2}}{5}\psi_1 e^{-3i\omega t/2} + \frac{2\sqrt{2}}{5}\psi_2 e^{-5i\omega t/2} = e^{-i\omega t/2} \left[\frac{3}{5}\psi_0 - \frac{2\sqrt{2}}{5}\psi_1 e^{-i\omega t} + \frac{2\sqrt{2}}{5}\psi_2 e^{-2i\omega t} \right].$$

To change the sign of the middle term we need $e^{-i\omega T} = -1$ (then $e^{-2i\omega T} = 1$); evidently $\omega T = \pi$, or

$$\boxed{T = \pi/\omega}.$$

Problem 2.42

Everything in Section 2.3.2 still applies, except that there is an additional boundary condition: $\psi(0) = 0$. This eliminates all the *even* solutions ($n = 0, 2, 4, \dots$), leaving only the odd solutions. So

$$E_n = \left(n + \frac{1}{2}\right) \hbar\omega, \quad n = 1, 3, 5, \dots$$

Problem 2.43

(a) Normalization is the same as before: $A = \left(\frac{2a}{\pi}\right)^{1/4}$.

(b) Equation 2.103 says

$$\phi(k) = \frac{1}{\sqrt{2\pi}} \left(\frac{2a}{\pi}\right)^{1/4} \int_{-\infty}^{\infty} e^{-ax^2} e^{ilx} e^{-ikx} dx \quad [\text{same as before, only } k \rightarrow k - l] = \frac{1}{(2\pi a)^{1/4}} e^{-(k-l)^2/4a}.$$

$$\begin{aligned} \Psi(x, t) &= \frac{1}{\sqrt{2\pi}} \frac{1}{(2\pi a)^{1/4}} \int_{-\infty}^{\infty} \underbrace{e^{-(k-l)^2/4a} e^{i(kx - \hbar k^2 t/2m)}}_{e^{-l^2/4a} e^{-[(\frac{1}{4a} + i\frac{\hbar t}{2m})k^2 - (ix + \frac{l}{2a})k]}} dk \\ &= \frac{1}{\sqrt{2\pi}} \frac{1}{(2\pi a)^{1/4}} e^{-l^2/4a} \sqrt{\frac{\pi}{(\frac{1}{4a} + i\frac{\hbar t}{2m})}} e^{(ix + l/2a)^2/[4(1/4a + i\hbar t/2m)]} \\ &= \left(\frac{2a}{\pi}\right)^{1/4} \frac{1}{\sqrt{1 + 2i\hbar at/m}} e^{-l^2/4a} e^{a(ix + l/2a)^2/(1 + 2i\hbar at/m)}. \end{aligned}$$

(c) Let $\theta \equiv 2\hbar at/m$, as before: $|\Psi|^2 = \sqrt{\frac{2a}{\pi}} \frac{1}{\sqrt{1 + \theta^2}} e^{-l^2/2a} e^{a \left[\frac{(ix + l/2a)^2}{(1 + i\theta)} + \frac{(-ix + l/2a)^2}{(1 - i\theta)} \right]}$. Expand the term in square brackets:

$$\begin{aligned} [] &= \frac{1}{1 + \theta^2} \left[(1 - i\theta) \left(ix + \frac{l}{2a} \right)^2 + (1 + i\theta) \left(-ix + \frac{l}{2a} \right)^2 \right] \\ &= \frac{1}{1 + \theta^2} \left[\left(-x^2 + \frac{ixl}{a} + \frac{l^2}{4a^2} \right) + \left(-x^2 - \frac{ixl}{a} + \frac{l^2}{4a^2} \right) \right. \\ &\quad \left. + i\theta \left(x^2 - \frac{ixl}{a} - \frac{l^2}{4a^2} \right) + i\theta \left(-x^2 - \frac{ixl}{a} + \frac{l^2}{4a^2} \right) \right] \\ &= \frac{1}{1 + \theta^2} \left[-2x^2 + \frac{l^2}{2a^2} + 2\theta \frac{xl}{a} \right] = \frac{1}{1 + \theta^2} \left[-2x^2 + 2\theta \frac{xl}{a} - \frac{\theta^2 l^2}{2a^2} + \frac{\theta^2 l^2}{2a^2} + \frac{l^2}{2a^2} \right] \\ &= \frac{-2}{1 + \theta^2} \left(x - \frac{\theta l}{2a} \right)^2 + \frac{l^2}{2a^2}. \end{aligned}$$

$$|\Psi(x, t)|^2 = \sqrt{\frac{2}{\pi}} \sqrt{\frac{a}{1 + \theta^2}} e^{-l^2/2a} e^{-\frac{2a}{1 + \theta^2} (x - \theta l/2a)^2} e^{l^2/2a} = \sqrt{\frac{2}{\pi}} w e^{-2w^2 (x - \theta l/2a)^2},$$

where $w \equiv \sqrt{a/(1 + \theta^2)}$. The result is the same as before, except $x \rightarrow (x - \frac{\theta l}{2a}) = (x - \frac{\hbar l}{m} t)$, so $|\Psi|^2$ has the same (flattening Gaussian) shape – only this time the center moves at constant speed $v = \hbar l/m$.

(d)

$$\langle x \rangle = \int_{-\infty}^{\infty} x |\Psi(x, t)|^2 dx. \quad \text{Let } y \equiv x - \theta l / 2a = x - vt, \text{ so } x = y + vt.$$

$$= \int_{-\infty}^{\infty} (y + vt) \sqrt{\frac{2}{\pi}} w e^{-2w^2 y^2} dy = vt.$$

(The first integral is trivially zero; the second is 1 by normalization.)

$$= \boxed{\frac{\hbar l}{m} t}; \quad \langle p \rangle = m \frac{d\langle x \rangle}{dt} = \boxed{\hbar l}.$$

$$\langle x^2 \rangle = \int_{-\infty}^{\infty} (y + vt)^2 \sqrt{\frac{2}{\pi}} w e^{-2w^2 y^2} dy = \frac{1}{4w^2} + 0 + (vt)^2 \text{ (the first integral is same as before).}$$

$$\boxed{\langle x^2 \rangle = \frac{1}{4w^2} + \left(\frac{\hbar l t}{m} \right)^2} \quad \langle p^2 \rangle = -\hbar^2 \int_{-\infty}^{\infty} \Psi^* \frac{d^2 \Psi}{dx^2} dx.$$

$$\Psi = \left(\frac{2a}{\pi} \right)^{1/4} \frac{1}{\sqrt{1+i\theta}} e^{-l^2/4a} e^{a(ix+l/2a)^2/(1+i\theta)}, \text{ so } \frac{d\Psi}{dx} = \frac{2ia(ix + \frac{l}{2a})}{(1+i\theta)} \Psi;$$

$$\frac{d^2 \Psi}{dx^2} = \left[\frac{2ia(ix + l/2a)}{1+i\theta} \right] \frac{d\Psi}{dx} + \frac{2i^2 a}{1+i\theta} \Psi = \left[\frac{-4a^2(ix + l/2a)^2}{(1+i\theta)^2} - \frac{2a}{1+i\theta} \right] \Psi.$$

$$\begin{aligned} \langle p^2 \rangle &= \frac{4a^2 \hbar^2}{(1+i\theta)^2} \int_{-\infty}^{\infty} \left[\left(ix + \frac{l}{2a} \right)^2 + \frac{(1+i\theta)}{2a} \right] |\Psi|^2 dx \\ &= \frac{4a^2 \hbar^2}{(1+i\theta)^2} \int_{-\infty}^{\infty} \left[-\left(y + vt - \frac{il}{2a} \right)^2 + \frac{(1+i\theta)}{2a} \right] |\Psi|^2 dy \\ &= \frac{4a^2 \hbar^2}{(1+i\theta)^2} \left\{ -\int_{-\infty}^{\infty} y^2 |\Psi|^2 dy - 2 \left(vt - \frac{il}{2a} \right) \int_{-\infty}^{\infty} y |\Psi|^2 dy \right. \\ &\quad \left. + \left[-\left(vt - \frac{il}{2a} \right)^2 + \frac{(1+i\theta)}{2a} \right] \int_{-\infty}^{\infty} |\Psi|^2 dy \right\} \\ &= \frac{4a^2 \hbar^2}{(1+i\theta)^2} \left[-\frac{1}{4w^2} + 0 - \left(vt - \frac{il}{2a} \right)^2 + \frac{(1+i\theta)}{2a} \right] \\ &= \frac{4a^2 \hbar^2}{(1+i\theta)^2} \left\{ -\frac{1+\theta^2}{4a} - \left[\left(\frac{-il}{2a} \right) (1+i\theta) \right]^2 + \frac{(1+i\theta)}{2a} \right\} \\ &= \frac{a\hbar^2}{1+i\theta} \left[-(1-i\theta) + \frac{l^2}{a}(1+i\theta) + 2 \right] = \frac{a\hbar^2}{1+i\theta} \left[(1+i\theta) \left(1 + \frac{l^2}{a} \right) \right] = \boxed{\hbar^2(a + l^2)}. \end{aligned}$$

$$\sigma_x^2 = \langle x^2 \rangle - \langle x \rangle^2 = \frac{1}{4w^2} + \left(\frac{\hbar l t}{m} \right)^2 - \left(\frac{\hbar l t}{m} \right)^2 = \frac{1}{4w^2} \Rightarrow \boxed{\sigma_x = \frac{1}{2w}};$$

$$\sigma_p^2 = \langle p^2 \rangle - \langle p \rangle^2 = \hbar^2 a + \hbar^2 l^2 - \hbar^2 l^2 = \hbar^2 a, \text{ so } \boxed{\sigma_p = \hbar \sqrt{a}}.$$

(e) σ_x and σ_p are same as before, so the uncertainty principle still holds.


Problem 2.44

Equation 2.22 $\Rightarrow \psi(x) = A \sin kx + B \cos kx$, $0 \leq x \leq a$, with $k = \sqrt{2mE}/\hbar$.

Even solutions: $\psi(x) = \psi(-x) = A \sin(-kx) + B \cos(-kx) = -A \sin kx + B \cos kx$ ($-a \leq x \leq 0$).

$$\text{Boundary conditions} \begin{cases} \psi \text{ continuous at } 0 : B = B \text{ (no new condition).} \\ \psi' \text{ discontinuous (Eq. 2.125 with sign of } \alpha \text{ switched): } Ak + Ak = \frac{2m\alpha}{\hbar^2} B \Rightarrow B = \frac{\hbar^2 k}{m\alpha} A. \\ \psi \rightarrow 0 \text{ at } x = a : A \sin(ka) + \frac{\hbar^2 k}{m\alpha} A \cos(ka) = 0 \Rightarrow \tan(ka) = -\frac{\hbar^2 k}{m\alpha}. \end{cases}$$

$$\psi(x) = A \left(\sin kx + \frac{\hbar^2 k}{m\alpha} \cos kx \right) \quad (0 \leq x \leq a); \quad \psi(-x) = \psi(x).$$



From the graph, the allowed energies are slightly above

$$ka = \frac{n\pi}{2} \quad (n = 1, 3, 5, \dots) \quad \text{so} \quad E_n \gtrsim \frac{n^2 \pi^2 \hbar^2}{2m(2a)^2} \quad (n = 1, 3, 5, \dots).$$

These energies are somewhat higher than the corresponding energies for the infinite square well (Eq. 2.27, with $a \rightarrow 2a$). As $\alpha \rightarrow 0$, the straight line ($-\hbar^2 k/m\alpha$) gets steeper and steeper, and the intersections get closer to $n\pi/2$; the energies then reduce to those of the ordinary infinite well. As $\alpha \rightarrow \infty$, the straight line approaches horizontal, and the intersections are at $n\pi$ ($n = 1, 2, 3, \dots$), so $E_n \rightarrow \frac{n^2 \pi^2 \hbar^2}{2ma^2}$ – these are the allowed energies for the infinite square well of width a . At this point the barrier is impenetrable, and we have two isolated infinite square wells.

Odd solutions: $\psi(x) = -\psi(-x) = -A \sin(-kx) - B \cos(-kx) = A \sin(kx) - B \cos(kx)$ ($-a \leq x \leq 0$).

$$\text{Boundary conditions} \begin{cases} \psi \text{ continuous at } 0 : B = -B \Rightarrow B = 0. \\ \psi' \text{ discontinuous: } Ak - Ak = \frac{2m\alpha}{\hbar^2} (0) \text{ (no new condition).} \\ \psi(a) = 0 \Rightarrow A \sin(ka) = 0 \Rightarrow ka = \frac{n\pi}{2} \quad (n = 2, 4, 6, \dots). \end{cases}$$

$$\psi(x) = A \sin(kx), \quad (-a < x < a); \quad E_n = \frac{n^2 \pi^2 \hbar^2}{2m(2a)^2} \quad (n = 2, 4, 6, \dots).$$

These are the *exact* (even n) energies (and wave functions) for the infinite square well (of width $2a$). The point is that the *odd* solutions (even n) are *zero* at the origin, so they never “feel” the delta function at all.

Problem 2.45

$$\left. \begin{aligned} -\frac{\hbar^2}{2m} \frac{d^2 \psi_1}{dx^2} + V \psi_1 &= E \psi_1 \Rightarrow -\frac{\hbar^2}{2m} \psi_2 \frac{d^2 \psi_1}{dx^2} + V \psi_1 \psi_2 = E \psi_1 \psi_2 \\ -\frac{\hbar^2}{2m} \frac{d^2 \psi_2}{dx^2} + V \psi_2 &= E \psi_2 \Rightarrow -\frac{\hbar^2}{2m} \psi_1 \frac{d^2 \psi_2}{dx^2} + V \psi_1 \psi_2 = E \psi_1 \psi_2 \end{aligned} \right\} \Rightarrow -\frac{\hbar^2}{2m} \left[\psi_2 \frac{d^2 \psi_1}{dx^2} - \psi_1 \frac{d^2 \psi_2}{dx^2} \right] = 0.$$

But $\frac{d}{dx} \left[\psi_2 \frac{d\psi_1}{dx} - \psi_1 \frac{d\psi_2}{dx} \right] = \frac{d\psi_2}{dx} \frac{d\psi_1}{dx} + \psi_2 \frac{d^2\psi_1}{dx^2} - \frac{d\psi_1}{dx} \frac{d\psi_2}{dx} - \psi_1 \frac{d^2\psi_2}{dx^2} = \psi_2 \frac{d^2\psi_1}{dx^2} - \psi_1 \frac{d^2\psi_2}{dx^2}$. Since this is zero, it follows that $\psi_2 \frac{d\psi_1}{dx} - \psi_1 \frac{d\psi_2}{dx} = K$ (a constant). But $\psi \rightarrow 0$ at ∞ so the constant must be zero. Thus $\psi_2 \frac{d\psi_1}{dx} = \psi_1 \frac{d\psi_2}{dx}$, or $\frac{1}{\psi_1} \frac{d\psi_1}{dx} = \frac{1}{\psi_2} \frac{d\psi_2}{dx}$, so $\ln \psi_1 = \ln \psi_2 + \text{constant}$, or $\psi_1 = (\text{constant})\psi_2$. QED

Problem 2.46

$-\frac{\hbar^2}{2m} \frac{d^2\psi}{dx^2} = E\psi$ (where x is measured around the circumference), or $\frac{d^2\psi}{dx^2} = -k^2\psi$, with $k \equiv \frac{\sqrt{2mE}}{\hbar}$, so

$$\psi(x) = Ae^{ikx} + Be^{-ikx}.$$

But $\psi(x+L) = \psi(x)$, since $x+L$ is the same point as x , so

$$Ae^{ikx}e^{ikL} + Be^{-ikx}e^{-ikL} = Ae^{ikx} + Be^{-ikx},$$

and this is true for *all* x . In particular, for $x=0$:

$$(1) \quad Ae^{ikL} + Be^{-ikL} = A + B. \quad \text{And for } x = \frac{\pi}{2k} :$$

$$Ae^{i\pi/2}e^{ikL} + Be^{-i\pi/2}e^{-ikL} = Ae^{i\pi/2} + Be^{-i\pi/2}, \text{ or } iAe^{ikL} - iBe^{-ikL} = iA - iB, \text{ so}$$

$$(2) \quad Ae^{ikL} - Be^{-ikL} = A - B. \quad \text{Add (1) and (2): } 2Ae^{ikL} = 2A.$$


Either $A=0$, or else $e^{ikL}=1$, in which case $kL=2n\pi$ ($n=0, \pm 1, \pm 2, \dots$). But if $A=0$, then $Be^{-ikL}=B$, leading to the same conclusion. So for every positive n there are *two* solutions: $\psi_n^+(x) = Ae^{i(2n\pi x/L)}$ and $\psi_n^-(x) = Be^{-i(2n\pi x/L)}$ ($n=0$ is ok too, but in that case there is just *one* solution). Normalizing: $\int_0^L |\psi_{\pm}|^2 dx = 1 \Rightarrow A=B=1/\sqrt{L}$. Any *other* solution (with the same energy) is a linear combination of these.

$$\psi_n^{\pm}(x) = \frac{1}{\sqrt{L}} e^{\pm i(2n\pi x/L)}; \quad E_n = \frac{2n^2\pi^2\hbar^2}{mL^2} \quad (n=0, 1, 2, 3, \dots).$$


The theorem fails because here ψ does *not* go to zero at ∞ ; x is restricted to a finite range, and we are unable to determine the constant K (in Problem 2.45).

Problem 2.47


- (a) (i) $b=0 \Rightarrow$ ordinary finite square well. Exponential decay outside; sinusoidal inside (cos for ψ_1 , sin for ψ_2). No nodes for ψ_1 , one node for ψ_2 .



- (ii) Ground state is *even*. Exponential decay outside, sinusoidal inside the wells, hyperbolic cosine in barrier. First excited state is *odd* – hyperbolic sine in barrier. No nodes for ψ_1 , one node for ψ_2 .



- (iii) For $b \gg a$, same as (ii), but wave function very small in barrier region. Essentially two isolated finite square wells; ψ_1 and ψ_2 are degenerate (in energy); they are even and odd linear combinations of the ground states of the two separate wells.




- (b) From Eq. 2.157 we know that for $b = 0$ the energies fall slightly below

$$\left. \begin{aligned} E_1 + V_0 &\approx \frac{\pi^2 \hbar^2}{2m(2a)^2} = \frac{h}{4} \\ E_2 + V_0 &\approx \frac{4\pi^2 \hbar^2}{2m(2a)^2} = h \end{aligned} \right\} \text{ where } h \equiv \frac{\pi^2 \hbar^2}{2ma^2}.$$

For $b \gg a$, the width of each (isolated) well is a , so

$$E_1 + V_0 \approx E_2 + V_0 \approx \frac{\pi^2 \hbar^2}{2ma^2} = h \text{ (again, slightly below this).}$$



[Within each well, $\frac{d^2\psi}{dx^2} = -\frac{2m}{\hbar^2}(V_0 + E)\psi$, so the more *curved* the wave function, the higher the energy.]

- (c) In the (even) ground state the energy is *lowest* in configuration (i), with $b \rightarrow 0$, so the electron tends to draw the nuclei together, promoting *bonding* of the atoms. In the (odd) first excited state, by contrast, the electron drives the nuclei apart.

Problem 2.48

(a)

$$\frac{d\Psi}{dx} = \frac{2\sqrt{3}}{a\sqrt{a}} \cdot \begin{cases} 1, & (0 < x < a/2) \\ -1, & (a/2 < x < a) \end{cases} = \boxed{\frac{2\sqrt{3}}{a\sqrt{a}} \left[1 - 2\theta\left(x - \frac{a}{2}\right) \right]}.$$

(b)

$$\frac{d^2\Psi}{dx^2} = \frac{2\sqrt{3}}{a\sqrt{a}} \left[-2\delta\left(x - \frac{a}{2}\right) \right] = \boxed{-\frac{4\sqrt{3}}{a\sqrt{a}}\delta\left(x - \frac{a}{2}\right)}.$$

(c)

$$\langle H \rangle = -\frac{\hbar^2}{2m} \left(-\frac{4\sqrt{3}}{a\sqrt{a}} \right) \int \Psi^* \delta\left(x - \frac{a}{2}\right) dx = \frac{2\sqrt{3}\hbar^2}{ma\sqrt{a}} \underbrace{\Psi^*\left(\frac{a}{2}\right)}_{\sqrt{3/a}} = \frac{2 \cdot 3 \cdot \hbar^2}{m \cdot a \cdot a} = \boxed{\frac{6\hbar^2}{ma^2}}. \quad \checkmark$$

Problem 2.49

(a)

$$\frac{\partial\Psi}{\partial t} = \left(-\frac{m\omega}{2\hbar} \right) \left[\frac{a^2}{2} (-2i\omega e^{-2i\omega t}) + \frac{i\hbar}{m} - 2ax(-i\omega)e^{-i\omega t} \right] \Psi, \text{ so}$$

$$i\hbar \frac{\partial\Psi}{\partial t} = \left[-\frac{1}{2}ma^2\omega^2 e^{-2i\omega t} + \frac{1}{2}\hbar\omega + max\omega^2 e^{-i\omega t} \right] \Psi.$$

$$\frac{\partial\Psi}{\partial x} = \left[\left(-\frac{m\omega}{2\hbar} \right) (2x - 2ae^{-i\omega t}) \right] \Psi = -\frac{m\omega}{\hbar} (x - ae^{-i\omega t}) \Psi;$$

$$\frac{\partial^2\Psi}{\partial x^2} = -\frac{m\omega}{\hbar} \Psi - \frac{m\omega}{\hbar} (x - ae^{-i\omega t}) \frac{\partial\Psi}{\partial x} = \left[-\frac{m\omega}{\hbar} + \left(\frac{m\omega}{\hbar} \right)^2 (x - ae^{-i\omega t})^2 \right] \Psi.$$

$$\begin{aligned} -\frac{\hbar^2}{2m} \frac{\partial^2\Psi}{\partial x^2} + \frac{1}{2}m\omega^2 x^2 \Psi &= -\frac{\hbar^2}{2m} \left[-\frac{m\omega}{\hbar} + \left(\frac{m\omega}{\hbar} \right)^2 (x - ae^{-i\omega t})^2 \right] \Psi + \frac{1}{2}m\omega^2 x^2 \Psi \\ &= \left[\frac{1}{2}\hbar\omega - \frac{1}{2}m\omega^2 (x^2 - 2axe^{-i\omega t} + a^2e^{-2i\omega t}) + \frac{1}{2}m\omega^2 x^2 \right] \Psi \\ &= \left[\frac{1}{2}\hbar\omega + max\omega^2 e^{-i\omega t} - \frac{1}{2}m\omega^2 a^2 e^{-2i\omega t} \right] \Psi \\ &= i\hbar \frac{\partial\Psi}{\partial t} \text{ (comparing second line above). } \quad \checkmark \end{aligned}$$

(b)

$$\begin{aligned}
 |\Psi|^2 &= \sqrt{\frac{m\omega}{\pi\hbar}} e^{-\frac{m\omega}{2\hbar} \left[\left(x^2 + \frac{a^2}{2} (1 + e^{2i\omega t}) - \frac{i\hbar t}{m} - 2ax e^{i\omega t} \right) + \left(x^2 + \frac{a^2}{2} (1 + e^{-2i\omega t}) + \frac{i\hbar t}{m} - 2ax e^{-i\omega t} \right) \right]} \\
 &= \sqrt{\frac{m\omega}{\pi\hbar}} e^{-\frac{m\omega}{2\hbar} [2x^2 + a^2 + a^2 \cos(2\omega t) - 4ax \cos(\omega t)]}. \quad \text{But } a^2[1 + \cos(2\omega t)] = 2a^2 \cos^2 \omega t, \text{ so} \\
 &= \sqrt{\frac{m\omega}{\pi\hbar}} e^{-\frac{m\omega}{\hbar} [x^2 - 2ax \cos(\omega t) + a^2 \cos^2(\omega t)]} = \boxed{\sqrt{\frac{m\omega}{\pi\hbar}} e^{-\frac{m\omega}{\hbar} (x - a \cos \omega t)^2}}.
 \end{aligned}$$

The wave packet is a *Gaussian* of fixed shape, whose *center* oscillates back and forth sinusoidally, with amplitude a and angular frequency ω .

(c) Note that this wave function *is* correctly normalized (compare Eq. 2.59). Let $y \equiv x - a \cos \omega t$:

$$\begin{aligned}
 \langle x \rangle &= \int x |\Psi|^2 dx = \int (y + a \cos \omega t) |\Psi|^2 dy = 0 + a \cos \omega t \int |\Psi|^2 dy = \boxed{a \cos \omega t}. \\
 \langle p \rangle &= m \frac{d\langle x \rangle}{dt} = \boxed{-m a \omega \sin \omega t}. \quad \frac{d\langle p \rangle}{dt} = -m a \omega^2 \cos \omega t. \quad V = \frac{1}{2} m \omega^2 x^2 \implies \frac{dV}{dx} = m \omega^2 x. \\
 \langle -\frac{dV}{dx} \rangle &= -m \omega^2 \langle x \rangle = -m \omega^2 a \cos \omega t = \frac{d\langle p \rangle}{dt}, \text{ so Ehrenfest's theorem is satisfied.}
 \end{aligned}$$

Problem 2.50

(a)

$$\frac{\partial \Psi}{\partial t} = \left[-\frac{m\alpha}{\hbar^2} \frac{\partial}{\partial t} |x - vt| - i \frac{(E + \frac{1}{2} m v^2)}{\hbar} \right] \Psi; \quad \frac{\partial}{\partial t} |x - vt| = \begin{cases} -v, & \text{if } x - vt > 0 \\ v, & \text{if } x - vt < 0 \end{cases}.$$

We can write this in terms of the θ -function (Eq. 2.143):

$$2\theta(z) - 1 = \begin{cases} 1, & \text{if } z > 0 \\ -1, & \text{if } z < 0 \end{cases}, \text{ so } \frac{\partial}{\partial t} |x - vt| = -v[2\theta(x - vt) - 1].$$

$$i\hbar \frac{\partial \Psi}{\partial t} = \left\{ i \frac{m\alpha v}{\hbar} [2\theta(x - vt) - 1] + E + \frac{1}{2} m v^2 \right\} \Psi. \quad [\star]$$

$$\begin{aligned}
 \frac{\partial \Psi}{\partial x} &= \left[-\frac{m\alpha}{\hbar^2} \frac{\partial}{\partial x} |x - vt| + \frac{imv}{\hbar} \right] \Psi \\
 \frac{\partial}{\partial x} |x - vt| &= \{1, \text{ if } x > vt; -1, \text{ if } x < vt\} = 2\theta(x - vt) - 1. \\
 &= \left\{ -\frac{m\alpha}{\hbar^2} [2\theta(x - vt) - 1] + \frac{imv}{\hbar} \right\} \Psi.
 \end{aligned}$$

$$\frac{\partial^2 \Psi}{\partial x^2} = \left\{ -\frac{m\alpha}{\hbar^2} [2\theta(x - vt) - 1] + \frac{imv}{\hbar} \right\}^2 \Psi - \frac{2m\alpha}{\hbar^2} \left[\frac{\partial}{\partial x} \theta(x - vt) \right] \Psi.$$

But (from Problem 2.24(b)) $\frac{\partial}{\partial x}\theta(x - vt) = \delta(x - vt)$, so

$$\begin{aligned} & -\frac{\hbar^2}{2m} \frac{\partial^2 \Psi}{\partial x^2} - \alpha \delta(x - vt) \Psi \\ &= \left(-\frac{\hbar^2}{2m} \left\{ -\frac{m\alpha}{\hbar^2} [2\theta(x - vt) - 1] + \frac{imv}{\hbar} \right\}^2 + \alpha \delta(x - vt) - \alpha \delta(x - vt) \right) \Psi \\ &= -\frac{\hbar^2}{2m} \left\{ \frac{m^2 \alpha^2}{\hbar^4} \underbrace{[2\theta(x - vt) - 1]^2}_1 - \frac{m^2 v^2}{\hbar^2} - 2i \frac{mv}{\hbar} \frac{m\alpha}{\hbar^2} [2\theta(x - vt) - 1] \right\} \Psi \\ &= \left\{ -\frac{m\alpha^2}{2\hbar^2} + \frac{1}{2}mv^2 + i \frac{mv\alpha}{\hbar} [2\theta(x - vt) - 1] \right\} \Psi = i\hbar \frac{\partial \Psi}{\partial t} \quad (\text{compare } [\star]). \quad \checkmark \end{aligned}$$

(b)

$$|\Psi|^2 = \frac{m\alpha}{\hbar^2} e^{-2m\alpha|y|/\hbar^2} \quad (y \equiv x - vt).$$


$$\text{Check normalization: } 2 \frac{m\alpha}{\hbar^2} \int_0^\infty e^{-2m\alpha y/\hbar^2} dy = \frac{2m\alpha}{\hbar^2} \frac{\hbar^2}{2m\alpha} = 1. \quad \checkmark$$

$$\begin{aligned} \langle H \rangle &= \int_{-\infty}^\infty \Psi^* H \Psi dx. \quad \text{But } H\Psi = i\hbar \frac{\partial \Psi}{\partial t}, \text{ which we calculated above } [\star]. \\ &= \int \left\{ \frac{im\alpha v}{\hbar} [2\theta(y) - 1] + E + \frac{1}{2}mv^2 \right\} |\Psi|^2 dy = \boxed{E + \frac{1}{2}mv^2}. \end{aligned}$$

(Note that $[2\theta(y) - 1]$ is an *odd* function of y .) *Interpretation:* The wave packet is dragged along (at speed v) with the delta-function. The total energy is the energy it *would* have in a stationary delta-function (E), plus *kinetic* energy due to the motion ($\frac{1}{2}mv^2$).

Problem 2.51

(a)




(b) $\frac{d\psi_0}{dx} = -Aa \operatorname{sech}(ax) \tanh(ax); \quad \frac{d^2\psi_0}{dx^2} = -Aa^2 [-\operatorname{sech}(ax) \tanh^2(ax) + \operatorname{sech}(ax) \operatorname{sech}^2(ax)].$

$$\begin{aligned}
 H\psi_0 &= -\frac{\hbar^2}{2m} \frac{d^2\psi_0}{dx^2} - \frac{\hbar^2 a^2}{m} \operatorname{sech}^2(ax) \psi_0 \\
 &= \frac{\hbar^2}{2m} A a^2 [-\operatorname{sech}(ax) \tanh^2(ax) + \operatorname{sech}^3(ax)] - \frac{\hbar^2 a^2}{m} A \operatorname{sech}^3(ax) \\
 &= \frac{\hbar^2 a^2 A}{2m} [-\operatorname{sech}(ax) \tanh^2(ax) + \operatorname{sech}^3(ax) - 2 \operatorname{sech}^3(ax)] \\
 &= -\frac{\hbar^2 a^2}{2m} A \operatorname{sech}(ax) [\tanh^2(ax) + \operatorname{sech}^2(ax)].
 \end{aligned}$$

But $(\tanh^2 \theta + \operatorname{sech}^2 \theta) = \frac{\sinh^2 \theta}{\cosh^2 \theta} + \frac{1}{\cosh^2 \theta} = \frac{\sinh^2 \theta + 1}{\cosh^2 \theta} = 1$, so

$$= -\frac{\hbar^2 a^2}{2m} \psi_0. \quad \text{QED} \quad \text{Evidently } \boxed{E = -\frac{\hbar^2 a^2}{2m}}.$$

$$1 = |A|^2 \int_{-\infty}^{\infty} \operatorname{sech}^2(ax) dx = |A|^2 \frac{1}{a} \tanh(ax) \Big|_{-\infty}^{\infty} = \frac{2}{a} |A|^2 \implies \boxed{A = \sqrt{\frac{a}{2}}}.$$



(c)

$$\begin{aligned}
 \frac{d\psi_k}{dx} &= \frac{A}{ik+a} [(ik - a \tanh ax) ik - a^2 \operatorname{sech}^2 ax] e^{ikx}. \\
 \frac{d^2\psi_k}{dx^2} &= \frac{A}{ik+a} \{ ik [(ik - a \tanh ax) ik - a^2 \operatorname{sech}^2 ax] - a^2 ik \operatorname{sech}^2 ax + 2a^3 \operatorname{sech}^2 ax \tanh ax \} e^{ikx}. \\
 -\frac{\hbar^2}{2m} \frac{d^2\psi_k}{dx^2} + V\psi_k &= \frac{A}{ik+a} \left\{ \frac{-\hbar^2 ik}{2m} [-k^2 - iak \tanh ax - a^2 \operatorname{sech}^2 ax] + \frac{\hbar^2 a^2}{2m} ik \operatorname{sech}^2 ax \right. \\
 &\quad \left. - \frac{\hbar^2 a^3}{m} \operatorname{sech}^2 ax \tanh ax - \frac{\hbar^2 a^2}{m} \operatorname{sech}^2 ax (ik - a \tanh ax) \right\} e^{ikx} \\
 &= \frac{A e^{ikx}}{ik+a} \frac{\hbar^2}{2m} (ik^3 - ak^2 \tanh ax + ia^2 k \operatorname{sech}^2 ax + ia^2 k \operatorname{sech}^2 ax \\
 &\quad - 2a^3 \operatorname{sech}^2 ax \tanh ax - 2ia^2 k \operatorname{sech}^2 ax + 2a^3 \operatorname{sech}^2 ax \tanh ax) \\
 &= \frac{A e^{ikx}}{ik+a} \frac{\hbar^2}{2m} k^2 (ik - a \tanh ax) = \frac{\hbar^2 k^2}{2m} \psi_k = E\psi_k. \quad \text{QED}
 \end{aligned}$$

As $x \rightarrow +\infty$, $\tanh ax \rightarrow +1$, so $\boxed{\psi_k(x) \rightarrow A \left(\frac{ik-a}{ik+a} \right) e^{ikx}}$, which represents a transmitted wave.

$$\boxed{R=0.} \quad T = \left| \frac{ik-a}{ik+a} \right|^2 = \left(\frac{-ik-a}{-ik+a} \right) \left(\frac{ik-a}{ik+a} \right) = \boxed{1.}$$

Problem 2.52

(a) (1) From Eq. 2.133: $F + G = A + B$.

(2) From Eq. 2.135: $F - G = (1 + 2i\beta)A - (1 - 2i\beta)B$, where $\beta = m\alpha/\hbar^2 k$.

Subtract: $2G = -2i\beta A + 2(1 - i\beta)B \Rightarrow B = \frac{1}{1 - i\beta}(i\beta A + G)$. Multiply (1) by $(1 - 2i\beta)$ and add:

$$2(1 - i\beta)F - 2i\beta G = 2A \Rightarrow F = \frac{1}{1 - i\beta}(A + i\beta G). \quad \boxed{S = \frac{1}{1 - i\beta} \begin{pmatrix} i\beta & 1 \\ 1 & i\beta \end{pmatrix}}.$$

(b) For an *even* potential, $V(-x) = V(x)$, scattering from the right is the same as scattering from the left, with $x \leftrightarrow -x$, $A \leftrightarrow G$, $B \leftrightarrow F$ (see Fig. 2.22): $F = S_{11}G + S_{12}A$, $B = S_{21}G + S_{22}A$. So $S_{11} = S_{22}$, $S_{21} = S_{12}$. (Note that the delta-well S matrix in (a) has this property.) In the case of the finite square well, Eqs. 2.167 and 2.168 give

$$S_{21} = \frac{e^{-2ika}}{\cos 2la - i \frac{(k^2 + l^2)}{2kl} \sin 2la}; \quad S_{11} = \frac{i \frac{(l^2 - k^2)}{2kl} \sin 2la e^{-2ika}}{\cos 2la - i \frac{(k^2 + l^2)}{2kl} \sin 2la}. \quad \text{So}$$

$$\boxed{S = \frac{e^{-2ika}}{\cos 2la - i \frac{(k^2 + l^2)}{2kl} \sin 2la} \begin{pmatrix} i \frac{(l^2 - k^2)}{2kl} \sin 2la & 1 \\ 1 & i \frac{(l^2 - k^2)}{2kl} \sin 2la \end{pmatrix}}.$$

Problem 2.53

(a)

$$B = S_{11}A + S_{12}G \Rightarrow G = \frac{1}{S_{12}}(B - S_{11}A) = M_{21}A + M_{22}B \Rightarrow M_{21} = -\frac{S_{11}}{S_{12}}, \quad M_{22} = \frac{1}{S_{12}}.$$

$$F = S_{21}A + S_{22}G = S_{21}A + \frac{S_{22}}{S_{12}}(B - S_{11}A) = -\frac{(S_{11}S_{22} - S_{12}S_{21})}{S_{12}}A + \frac{S_{22}}{S_{12}}B = M_{11}A + M_{12}B.$$

$$\Rightarrow M_{11} = -\frac{\det S}{S_{12}}, \quad M_{12} = \frac{S_{22}}{S_{12}}. \quad \boxed{M = \frac{1}{S_{12}} \begin{pmatrix} -\det(S) & S_{22} \\ -S_{11} & 1 \end{pmatrix}}. \quad \text{Conversely:}$$

$$G = M_{21}A + M_{22}B \Rightarrow B = \frac{1}{M_{22}}(G - M_{21}A) = S_{11}A + S_{12}G \Rightarrow S_{11} = -\frac{M_{21}}{M_{22}}; \quad S_{12} = \frac{1}{M_{22}}.$$


$$F = M_{11}A + M_{12}B = M_{11}A + \frac{M_{12}}{M_{22}}(G - M_{21}A) = \frac{(M_{11}M_{22} - M_{12}M_{21})}{M_{22}}A + \frac{M_{12}}{M_{22}}G = S_{21}A + S_{22}G.$$

$$\Rightarrow S_{21} = \frac{\det M}{M_{22}}; \quad S_{22} = \frac{M_{12}}{M_{22}}. \quad \boxed{S = \frac{1}{M_{22}} \begin{pmatrix} -M_{21} & 1 \\ \det(M) & M_{12} \end{pmatrix}}.$$

[It happens that the time-reversal invariance of the Schrödinger equation, plus conservation of probability, requires $M_{22} = M_{11}^*$, $M_{21} = M_{12}^*$, and $\det(M) = 1$, but I won't use this here. See Merzbacher's *Quantum Mechanics*. Similarly, for *even* potentials $S_{11} = S_{22}$, $S_{12} = S_{21}$ (Problem 2.52).]

$$R_l = |S_{11}|^2 = \left| \frac{M_{21}}{M_{22}} \right|^2, \quad T_l = |S_{21}|^2 = \left| \frac{\det(M)}{M_{22}} \right|^2, \quad R_r = |S_{22}|^2 = \left| \frac{M_{12}}{M_{22}} \right|^2, \quad T_r = |S_{12}|^2 = \left| \frac{1}{M_{22}} \right|^2.$$

(b)



$$\begin{pmatrix} F \\ G \end{pmatrix} = M_2 \begin{pmatrix} C \\ D \end{pmatrix}, \quad \begin{pmatrix} C \\ D \end{pmatrix} = M_1 \begin{pmatrix} A \\ B \end{pmatrix}, \quad \text{so} \quad \begin{pmatrix} F \\ G \end{pmatrix} = M_2 M_1 \begin{pmatrix} A \\ B \end{pmatrix} = M \begin{pmatrix} A \\ B \end{pmatrix}, \quad \text{with } M = M_2 M_1. \quad \text{QED}$$

(c)

$$\psi(x) = \begin{cases} Ae^{ikx} + Be^{-ikx} & (x < a) \\ Fe^{ikx} + Ge^{-ikx} & (x > a) \end{cases}.$$

$$\begin{cases} \text{Continuity of } \psi : & Ae^{ika} + Be^{-ika} = Fe^{ika} + Ge^{-ika} \\ \text{Discontinuity of } \psi' : & ik(Fe^{ika} - Ge^{-ika}) - ik(Ae^{ika} - Be^{-ika}) = -\frac{2m\alpha}{\hbar^2}\psi(a) = -\frac{2m\alpha}{\hbar^2}(Ae^{ika} + Be^{-ika}). \end{cases}$$

$$(1) \quad Fe^{2ika} + G = Ae^{2ika} + B.$$

$$(2) \quad Fe^{2ika} - G = Ae^{2ika} - B + i\frac{2m\alpha}{\hbar^2 k}(Ae^{2ika} + B).$$

Add (1) and (2):

$$2Fe^{2ika} = 2Ae^{2ika} + i\frac{2m\alpha}{\hbar^2 k}(Ae^{2ika} + B) \Rightarrow F = \left(1 + i\frac{m\alpha}{\hbar^2 k}\right)A + i\frac{m\alpha}{\hbar^2 k}e^{-2ika}B = M_{11}A + M_{12}B.$$

$$\text{So } M_{11} = (1 + i\beta); \quad M_{12} = i\beta e^{-2ika}; \quad \beta \equiv \frac{m\alpha}{\hbar^2 k}.$$

Subtract (2) from (1):

$$2G = 2B - 2i\beta e^{2ika}A - 2i\beta B \Rightarrow G = (1 - i\beta)B - i\beta e^{2ika}A = M_{21}A + M_{22}B.$$

$$\text{So } M_{21} = -i\beta e^{2ika}; \quad M_{22} = (1 - i\beta).$$

$$M = \begin{pmatrix} (1 + i\beta) & i\beta e^{-2ika} \\ -i\beta e^{2ika} & (1 - i\beta) \end{pmatrix}.$$

(d)

$$M_2 = \begin{pmatrix} (1 + i\beta) & i\beta e^{-2ika} \\ -i\beta e^{2ika} & (1 - i\beta) \end{pmatrix}; \quad \text{to get } M_1, \text{ just switch the sign of } a: \quad M_1 = \begin{pmatrix} (1 + i\beta) & i\beta e^{2ika} \\ -i\beta e^{-2ika} & (1 - i\beta) \end{pmatrix}.$$

$$M = M_2 M_1 = \begin{pmatrix} [1 + 2i\beta + \beta^2(e^{-4ika} - 1)] & 2i\beta[\cos 2ka - \beta \sin 2ka] \\ -2i\beta[\cos 2ka - \beta \sin 2ka] & [1 - 2i\beta + \beta^2(e^{4ika} - 1)] \end{pmatrix}.$$

$$T = T_l = T_r = \frac{1}{|M_{22}|^2} \Rightarrow$$


$$\begin{aligned} T^{-1} &= [1 + 2i\beta + \beta^2(e^{-4ika} - 1)][1 - 2i\beta + \beta^2(e^{4ika} - 1)] \\ &= 1 - 2i\beta + \beta^2 e^{4ika} - \beta^2 + 2i\beta + 4\beta^2 + 2i\beta^3 e^{4ika} - 2i\beta^3 + \beta^2 e^{-4ika} \\ &\quad - \beta^2 - 2i\beta^3 e^{-4ika} + 2i\beta^3 + \beta^4(1 - e^{-4ika} - e^{4ika} + 1) \\ &= 1 + 2\beta^2 + \beta^2(e^{-4ika} + e^{4ika}) - 2i\beta^3(e^{-4ika} - e^{4ika}) + 2\beta^4 - \beta^4(e^{-4ika} + e^{4ika}) \\ &= 1 + 2\beta^2 + 2\beta^2 \cos 4ka + 2i\beta^3 2i \sin 4ka + 2\beta^4 - 2\beta^4 \cos 4ka \\ &= 1 + 2\beta^2(1 + \cos 4ka) - 4\beta^3 \sin 4ka + 2\beta^4(1 - \cos 4ka) \\ &= 1 + 4\beta^2 \cos^2 2ka - 8\beta^3 \sin 2ka \cos 2ka - 4\beta^4 \sin^2 2ka \end{aligned}$$

$$T = \frac{1}{1 + 4\beta^2(\cos 2ka - \beta \sin 2ka)^2}$$


Problem 2.54

I'll just show the first two graphs, and the last two. Evidently K lies between 0.9999 and 1.0001


```
Plot[Evaluate[u[x] /.
  NDSolve[{u'[x] - (x^2 - 0.9)*u[x] == 0, u[0] == 1,
    u'[0] == 0}, u[x], {x, 10^-8, 10},
  MaxSteps -> 10000}], {x, 0, 10},
PlotRange -> {-10, 10}];
```




```
Plot[Evaluate[u[x] /.
  NDSolve[{u'[x] - (x^2 - 1.1)*u[x] == 0, u[0] == 1,
    u'[0] == 0}, u[x], {x, 10^-8, 10},
  MaxSteps -> 10000}], {x, 0, 10},
PlotRange -> {-10, 10}];
```



```
Plot[Evaluate[u[x] /.
  NDSolve[{u'[x] - (x^2 - 0.9999)*u[x] == 0, u[0] == 1,
    u'[0] == 0}, u[x], {x, 10^-8, 10}, MaxSteps -> 10000]],
{x, 4, 5.5}, PlotRange -> {-1, 10}];
```




```
Plot[Evaluate[u[x] /.
  NDSolve[{u'[x] - (x^2 - 1.0001)*u[x] == 0, u[0] == 1,
    u'[0] == 0}, u[x], {x, 10^-8, 10}, MaxSteps -> 10000]],
{x, 4, 5.5}, PlotRange -> {-10, 1}];
```




Problem 2.55

The *correct* values (in Eq. 2.72) are $K = 2n + 1$ (corresponding to $E_n = (n + \frac{1}{2})\hbar\omega$). I'll start by “guessing” 2.9, 4.9, and 6.9, and tweaking the number until I've got 5 reliable significant digits. The results (see below) are 3.0000, 5.0000, 7.0000. (The actual *energies* are these numbers multiplied by $\frac{1}{2}\hbar\omega$.)


```
Plot[Evaluate[u[x] /.
  NDSolve[{u'[x] - (x^2 - 2.9)*u[x] == 0, u[0] == 0,
    u'[0] == 1}, u[x], {x, 10^-8, 10},
  MaxSteps -> 10000]], {x, 0, 5},
  PlotRange -> {-1, 5}];
```




```
Plot[Evaluate[u[x] /.
  NDSolve[{u'[x] - (x^2 - 2.99999)*u[x] == 0,
    u[0] == 0, u'[0] == 1}, u[x], {x, 10^-8, 10},
  MaxSteps -> 10000]], {x, 0, 5.5},
  PlotRange -> {-1, 7}];
```




```
Plot[Evaluate[u[x] /.
  NDSolve[{u'[x] - (x^2 - 4.99999)*u[x] == 0,
    u[0] == 1, u'[0] == 0}, u[x], {x, 10^-8, 10},
  MaxSteps -> 10000]], {x, 0, 6},
  PlotRange -> {-1.5, 1.2}];
```




```
Plot[Evaluate[u[x] /.
  NDSolve[{u'[x] - (x^2 - 5.00001)*u[x] == 0,
    u[0] == 1, u'[0] == 0}, u[x], {x, 10^-8, 10},
  MaxSteps -> 10000]], {x, 0, 6},
  PlotRange -> {-1.5, 1.2}];
```




```
Plot[Evaluate[u[x] /.
  NDSolve[{u'[x] - (x^2 - 3.00001)*u[x] == 0,
    u[0] == 0, u'[0] == 1}, u[x], {x, 10^-8, 10},
  MaxSteps -> 10000]], {x, 0, 5.5},
  PlotRange -> {-0.5, 7}];
```




```
Plot[Evaluate[u[x] /.
  NDSolve[{u'[x] - (x^2 - 4.9)*u[x] == 0, u[0] == 1,
    u'[0] == 0}, u[x], {x, 10^-8, 10},
  MaxSteps -> 10000]], {x, 0, 4},
  PlotRange -> {-1.5, 1.2}];
```




```
Plot[Evaluate[u[x] /.
  NDSolve[{u'[x] - (x^2 - 6.9)*u[x] == 0, u[0] == 0,
    u'[0] == 1}, u[x], {x, 10^-8, 10},
  MaxSteps -> 10000]], {x, 0, 4.5},
  PlotRange -> {-1, .5}];
```



```
Plot[Evaluate[u[x] /.
  NDSolve[{u'[x] - (x^2 - 6.99999)*u[x] == 0,
    u[0] == 0, u'[0] == 1}, u[x], {x, 10^-8, 10},
  MaxSteps -> 10000]], {x, 0, 6.5},
  PlotRange -> {-1, .5}];
```




```
Plot[Evaluate[u[x] /.
  NDSolve[{u'[x] - (x^2 - 7.00001)*u[x] == 0,
    u[0] == 0, u'[0] == 1}, u[x], {x, 10^-8, 10},
  MaxSteps -> 10000]], {x, 0, 6.5},
  PlotRange -> {-1, .5}];
```




Problem 2.56

The Schrödinger equation says $-\frac{\hbar^2}{2m}\psi'' = E\psi$, or, with the *correct* energies (Eq. 2.27) and $a = 1$, $\psi'' + (n\pi)^2\psi = 0$. I'll start with a "guess" using 9 in place of π^2 (that is, I'll use 9 for the ground state, 36 for the first excited state, 81 for the next, and finally 144). Then I'll tweak the parameter until the graph crosses the axis right at $x = 1$. The results (see below) are, to five significant digits: 9.8696, 39.478, 88.826, 157.91. (The actual *energies* are these numbers multiplied by $\hbar^2/2ma^2$.)


```
Plot[Evaluate[u[x] /.
  NDSolve[{u'[x] + (9)*u[x] == 0, u[0] == 0, u'[0] == 1},
    u[x], {x, 10^-8, 1.5}, MaxSteps -> 10000]],
  {x, 0, 1.2}, PlotRange -> {-0.5, .5}];
```




```
Plot[Evaluate[u[x] /.
  NDSolve[{u'[x] + (9.86959)*u[x] == 0, u[0] == 0,
    u'[0] == 1}, u[x], {x, 10^-8, 1.005},
  MaxSteps -> 10000]], {x, 0.99999, 1.00001},
  PlotRange -> {-0.00001, .00001}];
```




```
Plot[Evaluate[u[x] /.
  NDSolve[{u'[x] + (36)*u[x] == 0, u[0] == 0, u'[0] == 1},
    u[x], {x, 10^-8, 1.5}, MaxSteps -> 10000]],
  {x, 0, 1.2}, PlotRange -> {-0.5, .5}];
```




```
Plot[Evaluate[u[x] /.
  NDSolve[{u'[x] + (39.47803)*u[x] == 0, u[0] == 0,
    u'[0] == 1}, u[x], {x, 10^-8, 1.005},
  MaxSteps -> 10000]], {x, 0.99999, 1.00001},
  PlotRange -> {-0.00001, .00001}];
```




```
Plot[Evaluate[u[x] /.
  NDSolve[{u'[x] + (81) * u[x] == 0, u[0] == 0, u'[0] == 1},
    u[x], {x, 10-8, 1.5}, MaxSteps -> 10000]],
  {x, 0, 1.2}, PlotRange -> {-0.15, 0.15}];
```



```
Plot[Evaluate[u[x] /.
  NDSolve[{u'[x] + (88.82630) * u[x] == 0, u[0] == 0,
    u'[0] == 1}, u[x], {x, 10-8, 1.005},
  MaxSteps -> 10000]], {x, 0.99999, 1.00001},
  PlotRange -> {-0.00001, 0.00001}];
```



```
Plot[Evaluate[u[x] /.
  NDSolve[{u'[x] + (144) * u[x] == 0, u[0] == 0, u'[0] == 1},
    u[x], {x, 10-8, 1.5}, MaxSteps -> 10000]],
  {x, 0, 1.2}, PlotRange -> {-0.1, 0.1}];
```



```
Plot[Evaluate[u[x] /.
  NDSolve[{u'[x] + (157.9129) * u[x] == 0, u[0] == 0,
    u'[0] == 1}, u[x], {x, 10-8, 1.005},
  MaxSteps -> 10000]], {x, 0.99999, 1.00001},
  PlotRange -> {-0.00001, 0.00001}];
```

