


Solutions for Introduction to Modern Astrophysics 2nd Edition by Carroll

[CLICK HERE TO ACCESS COMPLETE Solutions](#)



Solutions

Chapter 2

Celestial Mechanics

2.1 From Fig. 2.4, note that

$$r^2 = (x - ae)^2 + y^2 \quad \text{and} \quad r'^2 = (x + ae)^2 + y^2.$$

Substituting Eq. (2.1) into the second expression gives

$$r = 2a - \sqrt{(x + ae)^2 + y^2}$$

which is now substituted into the first expression. After some rearrangement,

$$\frac{x^2}{a^2} + \frac{y^2}{a^2(1 - e^2)} = 1.$$

Finally, from Eq. (2.2),

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1.$$

2.2 The area integral in Cartesian coordinates is given by

$$A = \int_{-a}^a \int_{-b\sqrt{1-x^2/a^2}}^{b\sqrt{1-x^2/a^2}} dy dx = \frac{2b}{a} \int_{-a}^a \sqrt{a^2 - x^2} dx = \pi ab.$$

2.3 (a) From Eq. (2.3) the radial velocity is given by

$$v_r = \frac{dr}{dt} = \frac{a(1 - e^2)}{(1 + e \cos \theta)^2} e \sin \theta \frac{d\theta}{dt}. \quad (\text{S2.1})$$

Using Eqs. (2.31) and (2.32)

$$\frac{d\theta}{dt} = \frac{2}{r^2} \frac{dA}{dt} = \frac{L}{\mu r^2}.$$

The angular momentum can be written in terms of the orbital period by integrating Kepler's second law. If we further substitute $A = \pi ab$ and $b = a(1 - e^2)^{1/2}$ then

$$L = 2\mu\pi a^2(1 - e^2)^{1/2}/P.$$

Substituting L and r into the expression for $d\theta/dt$ gives

$$\frac{d\theta}{dt} = \frac{2\pi(1 + e \cos \theta)^2}{P(1 - e^2)^{3/2}}.$$

This can now be used in Eq. (S2.1), which simplifies to

$$v_r = \frac{2\pi a e \sin \theta}{P(1 - e^2)^{1/2}}.$$

Similarly, for the transverse velocity

$$v_\theta = r \frac{d\theta}{dt} = \frac{2\pi a(1 + e \cos \theta)}{(1 - e^2)^{1/2} P}.$$

(b) Equation (2.36) follows directly from $v^2 = v_r^2 + v_\theta^2$, Eq. (2.37) (Kepler's third law), and Eq. (2.3).

2.4 The total energy of the orbiting bodies is given by

$$E = \frac{1}{2}m_1v_1^2 + \frac{1}{2}m_2v_2^2 - G\frac{m_1m_2}{r}$$

where $r = |\mathbf{r}_2 - \mathbf{r}_1|$. Now,

$$v_1 = \dot{\mathbf{r}}_1 = -\frac{m_2}{m_1 + m_2}\dot{\mathbf{r}} \quad \text{and} \quad v_2 = \dot{\mathbf{r}}_2 = \frac{m_1}{m_1 + m_2}\dot{\mathbf{r}}.$$

Finally, using $M = m_1 + m_2$, $\mu = m_1m_2/(m_1 + m_2)$, and $m_1m_2 = \mu M$, we obtain Eq. (2.25).

2.5 Following a procedure similar to Problem 2.4,

$$\begin{aligned} \mathbf{L} &= m_1\mathbf{r}_1 \times \mathbf{v}_1 + m_2\mathbf{r}_2 \times \mathbf{v}_2 \\ &= m_1 \left[-\frac{m_2}{m_1 + m_2} \right] \mathbf{r} \times \left[-\frac{m_2}{m_1 + m_2} \right] \mathbf{v} \\ &\quad + m_2 \left[\frac{m_1}{m_1 + m_2} \right] \mathbf{r} \times \left[\frac{m_1}{m_1 + m_2} \right] \mathbf{v} \\ &= \mu \mathbf{r} \times \mathbf{v} = \mathbf{r} \times \mathbf{p} \end{aligned}$$

2.6 (a) The total orbital angular momentum of the Sun–Jupiter system is given by Eq. (2.30). Referring to the data in Appendices A and C, $M_\odot = 1.989 \times 10^{30}$ kg, $M_J = 1.899 \times 10^{27}$ kg, $M = M_J + M_\odot = 1.991 \times 10^{30}$ kg, and $\mu = M_J M_\odot / (M_J + M_\odot) = 1.897 \times 10^{27}$ kg. Furthermore, $e = 0.0489$, $a = 5.2044$ AU = 7.786×10^{11} m. Substituting,

$$L_{\text{total orbit}} = \mu \sqrt{GMa(1 - e^2)} = 1.927 \times 10^{43} \text{ kg m}^2 \text{ s}^{-1}.$$

(b) The distance of the Sun from the center of mass is $a_\odot = \mu a / M_\odot = 7.426 \times 10^8$ m. The Sun's orbital speed is $v_\odot = 2\pi a_\odot / P_J = 12.46 \text{ m s}^{-1}$, where $P_J = 3.743 \times 10^8$ s is the system's orbital period. Thus, for an assumed circular orbit,

$$L_{\text{Sun orbit}} = M_\odot a_\odot v_\odot = 1.840 \times 10^{40} \text{ kg m}^2 \text{ s}^{-1}.$$

(c) The distance of Jupiter from the center of mass is $a_J = \mu a / M_J = 7.778 \times 10^{11}$ m, and its orbital speed is $v_J = 2\pi a_J / P_J = 1.306 \times 10^4 \text{ m s}^{-1}$. Again assuming a circular orbit,

$$L_{\text{Jupiter orbit}} = M_J a_J v_J = 1.929 \times 10^{43} \text{ kg m}^2 \text{ s}^{-1}.$$

This is in good agreement with

$$L_{\text{total orbit}} - L_{\text{Sun orbit}} = 1.925 \times 10^{43} \text{ kg m}^2 \text{ s}^{-1}.$$

- (d) The moment of inertia of the Sun is approximately

$$I_{\odot} \sim \frac{2}{5} M_{\odot} R_{\odot}^2 \sim 3.85 \times 10^{47} \text{ kg m}^2$$

and the moment of inertia of Jupiter is approximately

$$I_J \sim \frac{2}{5} M_J R_J^2 \sim 3.62 \times 10^{42} \text{ kg m}^2.$$

(Note: Since the Sun and Jupiter are centrally condensed, these values are overestimates; see Section 23.2.) Using $\omega = 2\pi/P$,

$$L_{\text{Sun rotate}} = 1.078 \times 10^{42} \text{ kg m}^2 \text{ s}^{-1}$$

$$L_{\text{Jupiter rotate}} = 6.312 \times 10^{38} \text{ kg m}^2 \text{ s}^{-1}.$$

- (e) Jupiter's orbital angular momentum.

2.7 (a) $v_{\text{esc}} = \sqrt{2GM_J/R_J} = 60.6 \text{ km s}^{-1}$

(b) $v_{\text{esc}} = \sqrt{2GM_{\odot}/1 \text{ AU}} = 42.1 \text{ km s}^{-1}.$

2.8 (a) From Kepler's third law (Eq. 2.37) with $a = R_{\oplus} + h = 6.99 \times 10^6 \text{ m}$, $P = 5820 \text{ s} = 96.9 \text{ min}.$

(b) The orbital period of a geosynchronous satellite is the same as Earth's sidereal rotation period, or $P = 8.614 \times 10^4 \text{ s}$. From Eq. (2.37), $a = 4.22 \times 10^7 \text{ m}$, implying an altitude of $h = a - R_{\oplus} = 3.58 \times 10^7 \text{ m} = 5.6 R_{\oplus}.$

(c) A geosynchronous satellite must be “parked” over the equator and orbiting in the direction of Earth's rotation. This is because the center of the satellite's orbit is the center of mass of the Earth–satellite system (essentially Earth's center).

- 2.9 The integral average of the potential energy is given by

$$\langle U \rangle = \frac{1}{P} \int_0^P U(t) dt = -\frac{1}{P} \int_0^P \frac{GM\mu}{r(t)} dt.$$

Using Eqs. (2.31) and (2.32) to solve for dt in terms of $d\theta$, and making the appropriate changes in the limits of integration,

$$\langle U \rangle = -\frac{1}{P} \int_0^{2\pi} \frac{GM\mu^2 r}{L} d\theta.$$

Writing r in terms of θ via Eq. (2.3) leads to

$$\begin{aligned} \langle U \rangle &= -\frac{GM\mu^2 a(1-e^2)}{PL} \int_0^{2\pi} \frac{d\theta}{1+e \cos \theta} \\ &= -\frac{2\pi GM\mu^2 a(1-e^2)^{1/2}}{PL}. \end{aligned}$$

Using Eq. (2.30) to eliminate the total orbital angular momentum L , and Kepler's third law (Eq. 2.37) to replace the orbital period P , we arrive at

$$\langle U \rangle = -G \frac{M\mu}{a}.$$

2.10 Using the integral average from Problem 2.9

$$\langle r \rangle = \frac{1}{P} \int_0^P r(t) dt.$$

Using substitutions similar to the solution of Problem 2.9 we eventually arrive at

$$\langle r \rangle = \frac{a}{2\pi} (1 - e^2)^{5/2} \int_0^{2\pi} \frac{d\theta}{(1 + e \cos \theta)^3}. \quad (\text{S2.2})$$

It is evident that for $e = 0$, $\langle r \rangle = a$, as expected for perfectly circular motion. However, $\langle r \rangle$ deviates from a for other values of e . This function is most easily evaluated numerically. Employing a simple trapezoid method with 10^6 intervals, gives the results shown in Table S2.1.

Table S2.1: Results of the numerical evaluation of Eq. (S2.2) for Problem 2.10.

e	$\langle r \rangle / a$
0.00000	1.000000
0.10000	1.005000
0.20000	1.020000
0.30000	1.045000
0.40000	1.080000
0.50000	1.125000
0.60000	1.180000
0.70000	1.245000
0.80000	1.320000
0.90000	1.405000
0.95000	1.451250
0.99000	1.490050
0.99900	1.499001
0.99990	1.499900
0.99999	1.499990
1.00000	0.000000

- 2.11 Since planetary orbits are very nearly circular (except Mercury and Pluto), the assumption of perfectly circular motion was a good approximation. Furthermore, since a geocentric model maintains circular motion, it was very difficult to make any observational distinction between geocentric and heliocentric universes. (Parallax effects are far too small to be noticeable with the naked eye.)
- 2.12 (a) The graph of $\log_{10} P$ vs. $\log_{10} a$ for the Galilean moons is given in Fig. S2.1.
 (b) Using the data for Io and Callisto, we find a slope of 1.5.
 (c) Assuming that the mass of Jupiter is much greater than the masses of any of the Galilean moons, Kepler's third law can be written as

$$\log M + 2 \log P = \log \left(\frac{4\pi^2}{G} \right) + 3 \log a,$$

or

$$\begin{aligned} \log P &= \frac{3}{2} \log a + \frac{1}{2} \log \left(\frac{4\pi^2}{G} \right) - \frac{1}{2} \log M. \\ &= m \log a + b \end{aligned}$$

where the y-intercept is

$$b = \frac{1}{2} \log \left(\frac{4\pi^2}{G} \right) - \frac{1}{2} \log M.$$

Solving for $\log M$ we have

$$\log M = \log \left(\frac{4\pi^2}{G} \right) - 2b.$$

Taking the slope as $m = 3/2$ and using the data for any of the Galilean moons we find $b = -7.753$ (in SI units). Solving gives $M = 1.900 \times 10^{27}$ kg, in good agreement with the value given in Appendix C and Problem 2.6.




Figure S2.1: $\log_{10} P$ vs. $\log_{10} a$ for the Galilean moons.

- 2.13 (a) Since the velocity and position vectors are perpendicular at perihelion and aphelion, conservation of angular momentum leads to $r_p v_p = r_a v_a$. Thus

$$\frac{v_p}{v_a} = \frac{r_a}{r_p} = \frac{1+e}{1-e},$$

where the last relation is obtained from Eqs. (2.5) and (2.6).

- (b) Conservation of energy at perihelion and aphelion gives

$$\frac{1}{2} \mu v_a^2 - G \frac{M\mu}{r_a} = \frac{1}{2} \mu v_p^2 - G \frac{M\mu}{r_p}.$$

Making use of Eqs. (2.5) and (2.6), and using the result of part (a) to replace v_p leads to

$$\frac{1}{2} v_a^2 - \frac{GM}{a(1+e)} = \frac{1}{2} v_a^2 \left(\frac{1+e}{1-e} \right)^2 - \frac{GM}{1(1-e)}.$$

After some manipulation, we obtain Eq. (2.34); Eq. (2.33) follows immediately.

- (c) The orbital angular momentum can now be obtained from

$$L = \mu r_p v_p = \mu a(1-e) \sqrt{\frac{GM}{a} \left(\frac{1+e}{1-e} \right)}.$$

Equation (2.30) follows directly.

- 2.14 (a) From Kepler's third law in the form $P^2 = a^3$ (P in years and a in AU), $a = 17.9$ AU.
 (b) Since $m_{\text{comet}} \ll M_{\odot}$, Kepler's third law in the form of Eq. (2.37) gives

$$M_{\odot} \simeq \frac{4\pi^2 a^3}{GP^2} = 1.98 \times 10^{30} \text{ kg}.$$

- (c) From Example 2.1.1, at perihelion $r_p = a(1 - e) = 0.585$ AU and at aphelion $r_a = a(1 + e) = 35.2$ AU.
 (d) At perihelion, Eq. (2.33) gives $v_p = 55 \text{ km s}^{-1}$, and at aphelion, Eq. (2.34) gives $v_a = 0.91 \text{ km s}^{-1}$.
 When the comet is on the semiminor axis $r = a$, and Eq. (2.36) gives

$$v = \sqrt{\frac{GM_{\odot}}{a}} = 7.0 \text{ km s}^{-1}.$$

(e) $K_p/K_a = (v_p/v_a)^2 = 3650.$

- 2.15 Using 50,000 time steps, $r = \sqrt{x^2 + y^2} \simeq 1$ AU when $t \simeq 0.105$ yr. (Note: **Orbit** can be downloaded from the companion web site at <http://www.cambridge.org/astrophysics>.)
 2.16 The data are plotted in Fig. S2.2. (Note: **Orbit** can be downloaded from the companion web site at <http://www.cambridge.org/astrophysics>.)




Figure S2.2: Results for Problem 2.16.

- 2.17 (Note: **Orbit** can be downloaded from the companion web site at <http://www.cambridge.org/astrophysics>.)
 (a) See Fig. S2.3.
 (b) See Fig. S2.3.
 (c) Figure S2.3 shows that the orbit of Mars is very close to a perfect circle, with the center of the circle slightly offset from the focal point of the ellipse. Kepler's early attempts at developing a model of the solar system based on perfect circles were not far off.
 2.18 A modified Fortran 95 version of **Orbit** that works for this problem is given below.




Figure S2.3: Results for Problem 2.17. The dots designate the elliptical orbit of Mars, and the principal focus of the ellipse is indicated by the circle at $(x, y) = (0, 0)$. The dashed line is for a perfect circle of radius $r = a = 1.5237$ AU centered at $x = -ae = -0.1423$ AU (marked by the \times).

- (a) The orbits generated by the modified **Orbit** are shown in Fig. S2.4.
- (b) The calculation indicates $S = 2.205$ yr.
- (c) Eq. (1.1) yields a value of $S = 2.135$ yr. The results do not agree exactly because the derivation of Eq. (1.1) assumes constant speeds throughout the orbits.
- (d) No, because the relative speeds during the partially-completed orbits are different.
- (e) Since the orbits are not circular, Mars is at different distances from Earth during different oppositions. The closest opposition occurs when Earth is at aphelion and Mars is at perihelion, as in the start of this calculation.




Figure S2.4: The orbits of Earth and Mars including correct eccentricities. The positions of two successive oppositions are shown. The first opposition occurs when Earth is at aphelion and Mars is at perihelion (the positions of closest approach).

```

PROGRAM Orbit
!
!   General Description:
!   =====
!   Orbit computes the orbit of a small mass about a much larger mass,
!   or it can be considered as computing the motion of the reduced mass
!   about the center of mass.
!
!   "An Introduction to Modern Astrophysics", Appendix J
!   Bradley W. Carroll and Dale A. Ostlie
!   Second Edition
!
!   Weber State University
!   Ogden, UT
!   modastro@weber.edu
!-----
!
!   *****This version has been modified for Problem 2.17*****
!
!   USE Constants, ONLY      :   i1, dp, G, AU, M_Sun, pi, two_pi, yr, &
!                               radians_to_degrees, eps_dp
!
!   IMPLICIT NONE
!   REAL(dp)                :: t, dt, LoM_E, LoM_M, P_E, P_M
!   REAL(dp)                :: Mstar, theta_E, dtheta_E, theta_M, dtheta_M, r_E, r_M
!   INTEGER                 :: n, k, kmax
!   INTEGER(i1)             :: ios      !I/O error flag
!   REAL(dp)               :: delta    !error range at end of period
!   CHARACTER               :: xpause
!
!   REAL(dp), PARAMETER    :: a_E = AU, a_M = 1.5236*AU, e_E = 0.0167, e_M = 0.0935
!
!   Open the output file
!   OPEN (UNIT = 10, FILE = "Orbit.txt", STATUS = 'REPLACE', ACTION = 'WRITE', &
!         IOSTAT = ios)
!   IF (ios /= 0) THEN
!     WRITE (*, '( " Unable to open Orbit.txt. --- Terminating calculation")')
!     STOP
!   END IF
!
!   Convert entered values to conventional SI units
!   Mstar = M_Sun
!
!   Calculate the orbital period of Earth in seconds using Kepler's Third Law (Eq. 2.37)
!   To be used to determine time steps
!   P_E = SQRT(4*pi**2*a_E**3/(G*Mstar))
!
!   Enter the number of time steps and the time interval to be printed
!   n = 100000
!   n = n + 1 !increment to include t=0 (initial) point
!   kmax = 100
!
!   Print header information for output file
!   WRITE (10, '( "t, theta_E, r_E, x_E, y_E, theta_M, r_M, x_M, y_M")')
!
!   Initialize print counter, angle, elapsed time, and time step.
!   k = 1 !printer counter
!   theta_E = 0 !angle from direction to perihelion (radians)
!   theta_M = 0
!
!   t = 0 !elapsed time (s)
!   dt = P_E/(n-1) !time step (s)
!   delta = eps_dp !allowable error at end of period
!
!   Start main time step loop
!   DO
!     Calculate the distance from the principal focus using Eq. (2.3); Kepler's First Law.
!     r_E = a_E*(1 - e_E**2)/(1 - e_E*COS(theta_E)) !Earth starts at aphelion
!     r_M = a_M*(1 - e_M**2)/(1 + e_M*COS(theta_M)) !Mars starts at perihelion
!
!     If time to print, convert to cartesian coordinates. Be sure to print last point also.
!     IF (k == 1 .OR. (theta_E - theta_M)/two_pi > 1 + delta) &

```

```

        WRITE (10, '(9F10.4)') t/yr, theta_E*radians_to_degrees, r_E/AU, &
            r_E*COS(theta_E)/AU, r_E*SIN(theta_E)/AU, &
            theta_M*radians_to_degrees, r_M/AU, &
            r_M*COS(theta_M)/AU, r_M*SIN(theta_M)/AU

!      Exit the loop if Earth laps Mars.
      IF ((theta_E - theta_M)/two_pi > 1 + delta) EXIT

!      Prepare for the next time step: Update the elapsed time.
      t = t + dt

!      Calculate the angular momentum per unit mass, L/m (Eq. 2.30).
      LoM_E = SQRT(G*Mstar*a_E*(1 - e_E**2))
      LoM_M = SQRT(G*Mstar*a_M*(1 - e_M**2))

!      Compute the next value for theta using the fixed time step by combining
!      Eq. (2.31) with Eq. (2.32), which is Kepler's Second Law.
      dtheta_E = LoM_E/r_E**2*dt
      theta_E = theta_E + dtheta_E

      dtheta_M = LoM_M/r_M**2*dt
      theta_M = theta_M + dtheta_M

!      Reset the print counter if necessary
      k = k + 1
      IF (k > kmax) k = 1

END DO

WRITE (*,'(/,"The calculation is finished and the data are in Orbit.txt"')')

WRITE (*,'(//,"Enter any character and press <enter> to exit:  ")', ADVANCE = 'NO')
READ (*,*) xpause
END PROGRAM Orbit

```

Chapter 3

The Continuous Spectrum of Light

- 3.1 (a) The parallax angle is $p = 33.6''/2 = 8.14 \times 10^{-5}$ rad. For a baseline $B = R_{\oplus}$ (Fig. 3.1),

$$d = \frac{B}{\tan p} = 7.83 \times 10^{10} \text{ m} = 0.523 \text{ AU}.$$

- (b) Suppose that the clocks of the two observers are out of sync by an amount Δt , causing the baseline for their distance measurement to be $2B + v_{\text{rel}}\Delta t$ instead of $2B$. Here, v_{rel} is the relative velocity of Mars and the Earth at the time of opposition,

$$v_{\text{rel}} = 29.79 \text{ km s}^{-1} - 24.13 \text{ km s}^{-1} = 5.66 \text{ km s}^{-1}.$$

The calculated value (which is wrong) of the distance between Earth and Mars is

$$d_{\text{calc}} = \frac{R_{\oplus}}{\tan p},$$

while the actual value is

$$d_{\text{act}} = \frac{R_{\oplus} + 0.5v_{\text{rel}}\Delta t}{\tan p}.$$

If the distance to Mars is to be measured to within 10 percent, then

$$\frac{d_{\text{act}} - d_{\text{calc}}}{d_{\text{act}}} = 0.1.$$

Inserting the expressions for d_{act} and d_{calc} and solving for Δt results in $\Delta t = 250$ s.

- 3.2 From Example 3.2.1, the solar constant is 1365 W m^{-2} . Using the inverse square law, Eq. (3.2), with $L = 100 \text{ W}$ gives $r = 0.0764 \text{ m} = 7.64 \text{ cm}$.

- 3.3 (a) From Eq. (3.1),

$$d = \frac{1}{p''} \text{ pc} = 2.64 \text{ pc} = 8.61 \text{ ly} = 5.44 \times 10^5 \text{ AU} = 8.14 \times 10^{16} \text{ m}.$$

- (b) The distance modulus for Sirius is, from Eq. (3.6),

$$m - M = 5 \log_{10}(d/10 \text{ pc}) = -2.89.$$