
Test Bank for Social Problems and the Quality of Life 14th Edition by Lauer

CLICK HERE TO ACCESS COMPLETE Test Bank



Test Bank

Social Problems and the Quality of Life, 14e (Lauer) Chapter 2 Alcohol and Other Drugs

1) Repeated use of a drug or alcohol to the point of periodic or chronic intoxication that is detrimental to the user or society is called
A) abuse
B) habit
C) addiction
D) social deviance
Answer: C
Accessibility: Keyboard Navigation
2) The term <i>aqua vitae</i> means
A) water of life
B) fresh water
C) living water
D) sea of life
Answer: A
Accessibility: Keyboard Navigation
3) The improper use of drugs or alcohol to the degree that the consequences are defined as detrimental to the user or to society is called A) illegal consumption B) immorality C) abuse D) illegal sales
Answer: C Accessibility: Keyboard Navigation
 4) Addiction has been called a because continued abuse of a drug causes changes in brain function that drive the addict to compulsive seeking and use the drug. A) brain retardation B) mental retardation C) mental illness D) brain disease
Answer: D Accessibility: Keyboard Navigation

5) The nation's most serious health problem is A) breast cancer B) AIDS C) ovarian cancer D) abuse of alcohol
Answer: D Accessibility: Keyboard Navigation
6) A blood alcohol level of about can make the individual feel a sense of release from tensions and inhibitions. A) 0.05 percent B) 0.10 percent C) 0.20 percent D) 0.30 percent E) 0.40 percent
Answer: A Accessibility: Keyboard Navigation
7) A blood alcohol level of about affects the individual's motor control and his hands, arms and legs become clumsy. A) 0.05 percent B) 0.10 percent C) 0.20 percent D) 0.30 percent E) 0.40 percent
Answer: B Accessibility: Keyboard Navigation
8) A blood alcohol level of about causes the motor and emotional functions of the brain to be impaired as the individual staggers and becomes intensely emotional. A) 0.05 percent B) 0.10 percent C) 0.20 percent D) 0.30 percent E) 0.40 percent
Answer: C Accessibility: Keyboard Navigation

to death. A) 0.05 percent B) 0.10 percent C) 0.20 percent D) 0.30 percent E) 0.40 percent	and may lead
Answer: E Accessibility: Keyboard Navigation	
10) The damaging effects of alcohol abuse are most obvious in a(n)	the individual
Answer: B Accessibility: Keyboard Navigation	
11) Alcoholism is defined in terms of four symptoms. Which of the following is those symptoms?A) a craving or compulsion to drinkB) loss of control to limit drinking on any particular occasionC) drinking beer instead of drinking liquorD) physical dependence on alcohol	NOT one of
Answer: C Accessibility: Keyboard Navigation	
 12) Compared to the general population, American Indians A) begin drinking at an early age B) drink less frequently C) refrain from drinking in greater amounts D) have a lesser alcohol-related death rate 	
Answer: A Accessibility: Keyboard Navigation	

- 13) According to a national survey, what percentage of Americans aged 12 or older are current drinkers?
- A) 6.7 percent
- B) 23.1 percent
- C) 52.7 percent
- D) 99.9 percent

Answer: C

Accessibility: Keyboard Navigation

- 14) Which of the following groups has the highest incidence of use and abuse of alcohol?
- A) American Indians
- B) African Americans
- C) Latinos
- D) whites

Answer: A

Accessibility: Keyboard Navigation

- 15) Which of the following statements is true about binge drinkers?
- A) They are more likely than others to use protection while having sex.
- B) They are more likely than others to be involved in property damage.
- C) They are less likely than others to fall behind in school.
- D) They are less likely than others to drive after drinking.

Answer: B

Accessibility: Keyboard Navigation

- 16) Which sex is more likely to abuse alcohol?
- A) males
- B) females
- C) both are equally likely to abuse alcohol
- D) in fact, neither sex abuses alcohol

Answer: A

Accessibility: Keyboard Navigation

- 17) Among male veterans, the death rate for alcoholics was how much higher compared to that of nonalcoholics?
- A) 1.5 times
- B) 2.5 times
- C) 3.5 times
- D) 4.5 times

Answer: B

Accessibility: Keyboard Navigation

18) In terms of cognitive ability, alcoholism costs the user about years of life. A) 2 B) 4 C) 6 D) 10
Answer: D Accessibility: Keyboard Navigation
19) The amount of alcoholics in Russia is that of the United States. A) the same as B) triple C) less than D) more than double
Answer: D Accessibility: Keyboard Navigation
20) Compared to those from nonalcoholic homes, the children of alcohol abusers are more likely to have A) lower rates of anxiety B) lower rates of depression C) higher rates of drug use D) higher rates of marriage
Answer: C Accessibility: Keyboard Navigation
21) The leading cause of death among men aged 36 to 60 in Hungary is A) heart disease B) cirrhosis of the liver C) cancer D) automobile accidents
Answer: B Accessibility: Keyboard Navigation
22) Any substance having no physiological effect that is given to a subject who believes it to be a drug that does have an effect is called A) an independent variable B) a dependent variable C) a placebo D) a test drug
Answer: C Accessibility: Keyboard Navigation

23) Which of the following drugs is the most widely used? A) cocaine B) crack C) alcohol D) marijuana
Answer: C Accessibility: Keyboard Navigation
24) In a national survey of those aged 12 or older, what percent of 18- to 25-year-olds used any illicit drug in the last 30 days? A) 10.1 percent B) 22 percent C) 18.5 percent D) 51.8 percent
Answer: B Accessibility: Keyboard Navigation
 25) In both rural and urban areas, children begin experimenting with drugs as early as which grade? A) third B) seventh C) freshman in high school D) senior in high school
Answer: A Accessibility: Keyboard Navigation
26) Worldwide, tobacco dependence is responsible for about deaths per year. A) 40,000 B) 400,000 C) 6 million D) 4 billion
Answer: C Accessibility: Keyboard Navigation
27) In the United States, the use of causes more physiological damage than any nonalcoholic drug and is the leading cause of preventable death. A) methamphetamine B) opium C) marijuana D) tobacco
Answer: D Accessibility: Keyboard Navigation

A) impaired psychomotor performance B) interference with the normal pattern of sex hormones C) adverse effects on the respiratory system D) reduced sperm count in males
Answer: A Accessibility: Keyboard Navigation
29) In 2012, federal funding for various drug control programs was over A) \$3 billion B) \$26 billion C) \$15 billion D) \$100 billion
Answer: B Accessibility: Keyboard Navigation
30) Supervised withdrawal from dependence on a drug is called A) a placebo B) detoxification C) remission D) freedom
Answer: B Accessibility: Keyboard Navigation
31) The American colonists generally viewed alcoholic beverages as A) the "work of the devil" B) an evil to be diligently avoided C) one of God's gifts D) the reason for succeeding from England
Answer: C Accessibility: Keyboard Navigation
32) Until the beginning of the 19 th century, the use of what substance was less offensive to Americans than was the smoking of cigarettes? A) LSD B) opium C) tranquilizers D) marijuana
Answer: B Accessibility: Keyboard Navigation

CLICK HERE TO ACCESS THE COMPLETE Test Bank

33) The damaging e who is addicted to a	effects of alcohol abuse are most obviousleshol.	ous in a(n)	—the individual
Answer: alcoholic Accessibility: Key	board Navigation		
34) Thes	ex is far more likely to abuse alcohol	than is the	sex.
Answer: male; fen Accessibility: Key			
35) A general loss of from alcoholism is	of intellectual abilities, impaired mem called	ory, and possible pers	sonality change
Answer: dementia Accessibility: Key			
36) Americans drin	k about 9.44 liters of alcohol a year. F	Russians drink about _	·
Answer: 15.8 liter Accessibility: Key			
37) The leading cau	se of death in Hungary is		
Answer: cirrhosis Accessibility: Key			
38) is the	most widely used drug.		
Answer: Alcohol Accessibility: Key	board Navigation		
39) At 19.6 percent days.	was the drug used the mo	st by people aged 18-	25 in the last 30
Answer: marijuana Accessibility: Key			
40) The term aqua	vitae means		
Answer: water of I Accessibility: Key			

CLICK HERE TO ACCESS THE COMPLETE Test Bank

Answer: Addiction Accessibility: Keyboard Navigation 42) Improper use of drugs or alcohol to the degree that the consequences are defined as detrimental to the user or society is known as Answer: abuse Accessibility: Keyboard Navigation 43) Any substance having no physiological effect that is given to a subject who believes it to be a drug that does have an effect is called a(n) Answer: placebo Accessibility: Keyboard Navigation 44) The American colonists generally viewed alcoholic beverages as Answer: one of God's gifts Accessibility: Keyboard Navigation 45) are prescription drugs—pain relievers, tranquilizers, stimulants, and sedatives—hat are used for their intoxication effects rather than for medical purposes. Answer: Psychotherapeutic drugs Accessibility: Keyboard Navigation 46) is defined as having five or more drinks on the same occasion. Answer: Binge drinking Accessibility: Keyboard Navigation 47) Alcohol abuse can lead to early death. Answer: TRUE Accessibility: Keyboard Navigation
Answer: abuse Accessibility: Keyboard Navigation 43) Any substance having no physiological effect that is given to a subject who believes it to be a drug that does have an effect is called a(n) Answer: placebo Accessibility: Keyboard Navigation 44) The American colonists generally viewed alcoholic beverages as Answer: one of God's gifts Accessibility: Keyboard Navigation 45) are prescription drugs—pain relievers, tranquilizers, stimulants, and sedatives—hat are used for their intoxication effects rather than for medical purposes. Answer: Psychotherapeutic drugs Accessibility: Keyboard Navigation 46) is defined as having five or more drinks on the same occasion. Answer: Binge drinking Accessibility: Keyboard Navigation 47) Alcohol abuse can lead to early death. Answer: TRUE
Accessibility: Keyboard Navigation 43) Any substance having no physiological effect that is given to a subject who believes it to be a frug that does have an effect is called a(n) Answer: placebo Accessibility: Keyboard Navigation 44) The American colonists generally viewed alcoholic beverages as Answer: one of God's gifts Accessibility: Keyboard Navigation 45) are prescription drugs—pain relievers, tranquilizers, stimulants, and sedatives—hat are used for their intoxication effects rather than for medical purposes. Answer: Psychotherapeutic drugs Accessibility: Keyboard Navigation 46) is defined as having five or more drinks on the same occasion. Answer: Binge drinking Accessibility: Keyboard Navigation 47) Alcohol abuse can lead to early death. Answer: TRUE
Answer: placebo Accessibility: Keyboard Navigation 44) The American colonists generally viewed alcoholic beverages as Answer: one of God's gifts Accessibility: Keyboard Navigation 45) are prescription drugs—pain relievers, tranquilizers, stimulants, and sedatives—hat are used for their intoxication effects rather than for medical purposes. Answer: Psychotherapeutic drugs Accessibility: Keyboard Navigation 46) is defined as having five or more drinks on the same occasion. Answer: Binge drinking Accessibility: Keyboard Navigation 47) Alcohol abuse can lead to early death. Answer: TRUE
Accessibility: Keyboard Navigation 44) The American colonists generally viewed alcoholic beverages as Answer: one of God's gifts Accessibility: Keyboard Navigation 45) are prescription drugs—pain relievers, tranquilizers, stimulants, and sedatives—hat are used for their intoxication effects rather than for medical purposes. Answer: Psychotherapeutic drugs Accessibility: Keyboard Navigation 46) is defined as having five or more drinks on the same occasion. Answer: Binge drinking Accessibility: Keyboard Navigation 47) Alcohol abuse can lead to early death. Answer: TRUE
Answer: one of God's gifts Accessibility: Keyboard Navigation 45) are prescription drugs—pain relievers, tranquilizers, stimulants, and sedatives—hat are used for their intoxication effects rather than for medical purposes. Answer: Psychotherapeutic drugs Accessibility: Keyboard Navigation 46) is defined as having five or more drinks on the same occasion. Answer: Binge drinking Accessibility: Keyboard Navigation 47) Alcohol abuse can lead to early death. Answer: TRUE
Accessibility: Keyboard Navigation 45) are prescription drugs—pain relievers, tranquilizers, stimulants, and sedatives—hat are used for their intoxication effects rather than for medical purposes. Answer: Psychotherapeutic drugs Accessibility: Keyboard Navigation 46) is defined as having five or more drinks on the same occasion. Answer: Binge drinking Accessibility: Keyboard Navigation 47) Alcohol abuse can lead to early death. Answer: TRUE
Answer: Psychotherapeutic drugs Accessibility: Keyboard Navigation 46) is defined as having five or more drinks on the same occasion. Answer: Binge drinking Accessibility: Keyboard Navigation 47) Alcohol abuse can lead to early death. Answer: TRUE
Accessibility: Keyboard Navigation 46) is defined as having five or more drinks on the same occasion. Answer: Binge drinking Accessibility: Keyboard Navigation 47) Alcohol abuse can lead to early death. Answer: TRUE
Answer: Binge drinking Accessibility: Keyboard Navigation 47) Alcohol abuse can lead to early death. Answer: TRUE
Accessibility: Keyboard Navigation 47) Alcohol abuse can lead to early death. Answer: TRUE
Answer: TRUE
48) Heavy drinking allows the person to "loosen up" and "release inhibitions," therefore making sexual activity more enjoyable for both parties.
Answer: FALSE Accessibility: Keyboard Navigation

49) Americans drink far more than Russians do.

Answer: FALSE

Accessibility: Keyboard Navigation

50) Early use of alcohol by children is likely to lead to early sexual activity.

Answer: TRUE

Accessibility: Keyboard Navigation

51) Homes in which one or both spouses are alcoholics have lower rates of both verbal and physical abuse.

Answer: FALSE

Accessibility: Keyboard Navigation

52) Both Jews and Italians use alcohol as part of a traditional way of life and therefore alcoholism is extremely high in these communities.

Answer: FALSE

Accessibility: Keyboard Navigation

53) Most researchers have found there is a tendency for multiple use when individuals begin using one drug.

Answer: TRUE

Accessibility: Keyboard Navigation

54) Smokers are likely to drink alcohol.

Answer: TRUE

Accessibility: Keyboard Navigation

55) We cannot say that the use of one drug causes the individual to experiment with another drug. To do so would be the *fallacy of non sequitur*.

Answer: TRUE

Accessibility: Keyboard Navigation

56) Nonsmokers who are exposed to a smoking environment are more likely than those not in such an environment to develop coronary heart disease.

Answer: TRUE

Accessibility: Keyboard Navigation

57) Drug abusers are less likely to come from broken homes than are nonabusers.

Answer: FALSE

Accessibility: Keyboard Navigation

58) Nicotine-dependent individuals are more likely to have higher self-esteem than nonusers.

Answer: FALSE

Accessibility: Keyboard Navigation

59) Whereas moderate alcohol consumption protects the heart, heavy consumption increases the risk of strokes.

Answer: TRUE

Accessibility: Keyboard Navigation

60) Being integrated into a group in which drug use is approved is one of the strongest factors associated with drug use at all ages.

Answer: TRUE

Accessibility: Keyboard Navigation

61) Marijuana users experience more cohesion and harmony in their relationships and have less relationship conflict than nonusers.

Answer: FALSE

Accessibility: Keyboard Navigation